

# Accessing the nucleon ~~tensor~~ <sup>transverse</sup> structure in inclusive deep inelastic scattering

Alberto Accardi<sup>a</sup>, Alessandro Bacchetta<sup>b</sup>  
<sup>a</sup>Hampton University, Hampton, VA 23668, USA,  
 and Jefferson Lab, Newport News, VA 23606, USA

<sup>b</sup>Dipartimento di Fisica, Università degli Studi di Pavia, and INFN, Sez. di Pavia, 27100 Pavia, Italy

(Dated: Thursday, May 4th, 2017)

We revisit the standard analysis of inclusive Deep Inelastic Scattering (DIS) off nucleons taking into account the fact that on-shell quarks cannot be present in the final state, but they rather decay into hadrons - a process that can be described in terms of suitable "jet" correlators. As a consequence, a spin-flip term associated with the invariant mass of the produced hadrons is generated non perturbatively, and couples to the target's transversity distribution function. In inclusive cross sections, this provides an hitherto neglected and large contribution to the twist-3 part of the  $g_2$  structure function, that can explain the discrepancy between recent calculations and fits of this quantity. It also provides an extension of the Burkardt-Cottingham sum rule, putting constraints on the small- $x$  behavior of the transversity function, as well as an extension of the Efremov-Teryaev-Leader sum rule, suggesting a novel way to measure the tensor charge of the proton.

## I. INTRODUCTION

The tensor charge is a fundamental property of the nucleon, that is at present poorly constrained but of fundamental importance, not the least because its knowledge can be also used to put constraints on the search of physics beyond the Standard Model [1–3]. The tensor charge has been estimated in lattice QCD (see, e.g., [4–8]), but only limited information is available from direct measurements. Its experimental extraction requires first of all the measurement of the so-called transversity parton distribution function, denoted by  $h_1^q(x)$  (see Ref. [9] for a review on transversity and Refs. [10–12] for the most recent extractions). Secondly, one needs to perform a flavor decomposition and evaluate its integral, that corresponds to the contribution of flavor  $q$  to the tensor charge.

The transversity distribution is notoriously difficult to access because it is a chiral-odd function and needs to be combined with a spin-flip mechanism to appear in a scattering process [13]. Usually, this spin flip is provided by another nonperturbative distribution or fragmentation function, accessible in Drell-Yan or semi-inclusive DIS [14–17]. The only other known way to attain spin-flip terms in QED and QCD is taking into account mass corrections. In fact, it is well known that the transversity distribution gives a contribution to the structure function  $g_2$  in inclusive DIS (see, e.g., [18] and references therein), and in particular to the violation of the so-called Wandzura-Wilczek relation for  $g_2$  [19]. However, this contribution is proportional to the current quark mass and can be expected to be negligibly small. This implies that transversity cannot be accessed in inclusive DIS, but only in more complex processes such as semi-inclusive DIS or dilepton production in hadron-hadron collisions.

In this paper, we discuss a novel way of accessing the transversity parton distribution function (PDF) and measuring the proton's tensor charge in totally inclusive Deep Inelastic Scattering. We revisit the standard analysis of inclusive DIS taking into account the fact that on-shell quarks cannot be present in the final state, but they rather decay into hadrons, typically forming mini-jets of hadrons. This is sufficient to modify the structure of the DIS cut diagram, even if none of the hadrons is detected in the final state. For a proper description of this effect, we include "jet correlators" into the analysis, and pay particular attention to ensuring that our results are gauge invariant. We observe that jet correlators introduce a new contribution to the inclusive  $g_2$  structure function. This term is proportional to the transversity distribution function multiplied by a nonperturbative "jet mass" parameter, much larger than the mass of light quarks, and has the interesting features that: (a) it violates the Wandzura-Wilczek relation; (b) it extends the Burkhardt-Cottingham sum rule, providing a useful bound on the small- $x$  behavior of the transversity distribution; (c) it also extends the Efremov-Teryaev-Leader sum rule, and provides a novel way to measure the proton's tensor charge. We provide estimates of this new jet-mass-induced contribution based on a recent extraction of transversity, and show that it can indeed be very large.

## II. THE JET CORRELATOR

Motivated by large- $x$  mass corrections to inclusive DIS structure functions, Accardi and Qiu [20] have introduced in the LO handbag diagram a "jet correlator", also called "jet factor" by Collins and Rogers in Ref. [21], that accounts for invariant mass production in the current region, and ensures that leading twist calculations in collinear factorization are consistent with the requirement imposed by baryon number conservation that  $x_B < 1$  [20]. The jet correlator is

⊗ Similarly, the transversity couples to the current quark mass  $m_q$  in two-photon interference processes such as in transversity-dependent single spin asymmetries (Sivers effect, Schlegel), where the term is forbidden by T-parity invariance (Chiral-odd) and the spin transversity is not doubly suppressed.




FIG. 1: Diagrams contributing to inclusive DIS scattering up to twist-3, including a jet correlator in the top part. Note the gluon attaches to both the nucleon and jet correlators. The Hermitian conjugates of diagrams (b) and (c), i.e., with gluons attaching to the right of the cut, are not shown.

depicted in Figure 1(a) and is defined as

$$\Xi_{ij}(l, n_+) = \int \frac{d^4\eta}{(2\pi)^4} e^{il \cdot \eta} \langle 0 | \mathcal{U}_{(+\infty, \eta)}^{n_+} \psi_i(\eta) \bar{\psi}_j(0) \mathcal{U}_{(0, +\infty)}^{n_+} | 0 \rangle, \quad (1)$$

*here,*  
In this definition,  $l$  is the quark's four-momentum,  $\Psi$  the quark field operator (with quark flavor index omitted for simplicity), and  $|0\rangle$  is the non-perturbative vacuum state. Furthermore, we explicitly guarantee the correlator's gauge invariance by introducing two Wilson line operators  $\mathcal{U}^{n_+}$  running to infinity first along a light-cone plus direction determined by the vector  $n_+$ , then along the direction transverse to that vector, see [22] for details. This path choice for the Wilson line is required by QCD factorization theorems, and the vector is determined by the particular hard process to which the jet correlator contributes. For example, in the case of inclusive DIS discussed in this paper, *this is determined by the four momentum transfer  $q$  and the proton's momentum  $p$ .*

The correlator  $\Xi$  can be parametrized in terms of jet parton correlation functions [22, 23], using the vectors  $l$  and  $n_+$ :

$$\Xi(l, n_+) = \Lambda A_1(l^2) \mathbf{1} + A_2(l^2) l + \frac{\Lambda^2}{l \cdot n_+} \not{n}_+ B_1(l^2) + \frac{i\Lambda}{2l \cdot n_+} [l, \not{n}_+] B_2(l^2), \quad (2)$$

where  $\Lambda$  is an arbitrary scale, introduced for power counting purposes. In this parametrization, no terms proportional to  $\gamma_5$  enter because of parity invariance. Time reversal invariance in QCD requires  $B_2 = 0$ , while  $B_1$  contributes only at twist-4 order, and will not be considered further in this paper. We focus, instead, on the role of chiral odd terms in the  $g_2$  structure function up to twist 3. At this order,

$$\Xi(l, n_+) = \Lambda A_1(l^2) \mathbf{1} + A_2(l^2) l + O(\Lambda^2/Q^2) \quad (3)$$

is nothing else than the cut quark propagator; note however, that we consider here the full QCD vacuum rather than the perturbative one (or, in other words, the interacting rather than the free quark fields). The  $A_1$  and  $A_2$  terms can be nicely interpreted in terms of the spectral representation of the cut quark propagator (see, e.g., Sec. 6.3 of [23] and Sec. 2.7.2 of [24]),

$$\Xi(l) = \int d\mu^2 [J_1(\mu^2) \mu \mathbf{1} + J_2(\mu^2) l] \delta(l^2 - \mu^2), \quad (4)$$

where  $\mu^2$  can be interpreted as the invariant mass of the current jet, i.e., of the particles going through the cut in the top blob of Fig.1(a). The  $J_i$  are the spectral functions of the quark propagator, also called “jet functions” in [20], that can be interpreted as the current jet mass distributions. These satisfy [23, 24, 25]

$$J_2(\mu^2) \geq J_1(\mu^2) \geq 0 \quad \text{and} \quad \int d\mu^2 J_2(\mu^2) = 1. \quad (5)$$

From a comparison of Eqns.(2) and (4), one can see that

$$A_1(l^2) = \frac{\sqrt{l^2}}{\Lambda} J_1(l^2) \quad A_2(l^2) = J_2(l^2). \quad (6)$$

When inserting the jet correlator in the handbag diagram for inclusive DIS, the integration over  $dl^+$ , or equivalently  $dl^2/(2l^-)$ , is kinematically coupled to the other integrations, and induces corrections of order  $O(1/Q^2)$ , whose effect

on the  $F_2$  structure function has been studied in Ref. [20]. In this paper, where we limit our attention to effects of order  $O(1/Q)$ , we can neglect  $k^-$  compared to  $q^-$ . As a consequence, we can extend the integration over  $dl^2$  to infinity, the jet correlator decouples from the parton correlator  $\Phi$ , and the inclusive structure functions only depend on the integrated jet correlator

$$\Xi(l^-, l_T) \equiv \int \frac{dl^2}{2l^-} \Xi(l) = \frac{\Lambda}{2l^-} \xi_1 \mathbf{1} + \xi_2 \frac{h_-}{2} + \text{higher twists} . \quad (7)$$

where the terms up to twist-3 are independent of  $l_T$ , and the neglected higher twist terms only contribute to  $O(1/Q^2)$  to the cross section. Note that thanks to Eq. (5) we also obtain

$$\xi_1 = \int d\mu^2 \frac{\mu}{\Lambda} J_1(\mu^2) \equiv \frac{M_q}{\Lambda}, \quad \xi_2 = \int d\mu^2 J_2(\mu^2) = 1 . \quad (8)$$

Here,  $M_q$  can be interpreted as the average invariant mass produced in the spin-flip fragmentation processes of a quark of flavor  $q$ . It is important to notice that  $\xi_2 = 1$  exactly due to CPT invariance (see Sec. 10.7 of Ref. [25]), while  $0 < M_q < \int d\mu^2 \mu J_2(\mu^2)$  is dynamically determined. From the analytic properties of spectral functions we know that  $J_2(\mu^2) = Z\delta(\mu^2 - m_q) + \bar{J}_2(\mu^2)\theta(\mu^2 - m_\pi^2)$  with the continuum starting at  $m_\pi$  the mass of the pion due to color confinement effects [20]. Taking into account that  $J_1 < J_2$ , we may therefore estimate

$$M_q = O(100 \text{ MeV}) . \quad (9)$$

Although  $M_q$  is in general a nonperturbative quantity, it is interesting to notice that on the perturbative vacuum,

$$\Xi^{\text{pert}}(l) = (l + m_q \mathbf{1}) \delta(l^2 - m_q^2) + O(\alpha_s) ,$$

where  $m_q$  is the current quark mass; one recovers, therefore, the result of the calculation with the conventional handbag diagram. However, we are here considering non perturbative effects on the quark fragmentation and  $M_q > m_q$ . Therefore, differently from  $J_2$ , the  $J_1$  function leaves an imprint on the inclusive DIS cross section even in the asymptotic  $Q^2 \rightarrow \infty$  regime.

### III. TWIST-3 ANALYSIS

Extending the analysis to the calculation of twist-3 structure functions requires not only to consider the  $\xi_1$  term in the jet correlator, but also quark-gluon-quark correlators in both the proton and the vacuum as depicted in Figs.1(b) and (c), respectively.

In diagram (b), the  $\xi_1$  terms contribute to  $O(1/Q^2)$ , so that up to  $O(1/Q)$  these give the same contribution as in the conventional handbag calculation. The novel element in our analysis, instead, is the jet's quark-gluon-quark correlator  $\Xi_A^\mu(l, k)$  in diagram (c), defined as

$$(\Xi_A^\mu)_{ij} = \frac{1}{2} \int \frac{d\eta^+ d\eta_T}{(2\pi)^3} e^{ik \cdot \eta} \langle 0 | \mathcal{U}_{(+\infty, \eta)}^{n+} g A^\mu(\eta) \psi_i(\eta) | \bar{\psi}_j(0) \mathcal{U}_{(0, +\infty)}^{n+} | 0 \rangle \quad (10)$$

This diagram and its Hermitian conjugate are not only important to account for all contribution of order  $O(1/Q)$ , but also to restore gauge invariance, which is broken in diagram 1(a) due to the different mass of the incoming and outgoing quark lines, namely,  $m_q \neq M_q$ .

Rather than directly using the definition (10), it is convenient to calculate the inclusive cross section as an integral of the semi-inclusive one summed over all produced hadron flavors, then utilize the QCD equation of motions, sum over all hadron flavors, and take advantage of

$$\sum_h \int d^2 p_{hT} \frac{dp_h^-}{2p_h^-} \Delta^h(l, p_h) = \Xi(l) \quad (11)$$

where  $\Delta^h$  is the quark fragmentation correlator for production of a hadron of flavor  $h$  and momentum  $p_h$  [22]. In terms of the TMD fragmentation functions we are interested in, the sum rule (11) reads

$$\sum_h \int dz d^2 p_{hT} z D_1^h(z, p_{hT}) = \xi_2 = 1 \quad (12)$$

$$\sum_h \int dz d^2 p_{hT} E^h(z, p_{hT}) = \xi_1 = M_q/\Lambda , \quad (13)$$

where  $D_1^h(z, p_{hT})$  is the twist-2 quark fragmentation function, that depends on the hadron's collinear momentum fraction  $z$  and transverse momentum  $p_{hT}$ , and  $E^h(z, p_{hT})$  is a chiral-odd twist-3 function defined in [22].

The role of the  $\xi_1 = M_q/\Lambda$  term in inclusive DIS can be discussed by analyzing the following terms of the semi-inclusive hadronic tensor:

$$2\Lambda W^{\mu\nu} = i\frac{2\Lambda}{Q}\hat{t}^{[\mu}\epsilon_{\perp}^{\nu]\rho}S_{\perp\rho}\sum_q e_q^2\left[2x_B g_T^q(x_B)\sum_h\int dz d^2p_{hT} D_1^{q,h}(z, p_{hT}) + 2h_1^q(x_B)\sum_h\int dz d^2p_{hT} \tilde{E}^{q,h}(z, p_{hT})\right] + \dots, \quad (14)$$

where  $g_T^q$  and  $\tilde{E}^q$  are twist-3 TMDs originating, respectively, from the quark-quark and the quark-gluon-quark fragmentation correlators. For clarity, here we reintroduced the quark flavor  $q$ ,  $e_q$  being its electric charge. The first term can be easily integrated with the help of the sum rules (III) and (I). To integrate the second term, however, we first need make use of the relation  $\tilde{E}(z) = E(z) - (m_q/\Lambda)zD_1(z)$ , which is a consequence of the QCD equations of motion [22], then make use of the sum rule (11) *to obtain*  $1??$   $???$

$$\sum_h\int dz d^2p_{hT} \tilde{E}^{q,h}(z, p_{hT}) = \sum_h\int dz d^2p_{hT} \left[E^{q,h}(z, p_{hT}) - \frac{m_q}{\Lambda}zD_1^{q,h}(z, p_{hT})\right] = \xi_1 - \frac{m_q}{\Lambda}\xi_2 = \frac{M_q - m_q}{\Lambda}. \quad (15)$$

This formula is the single most important result of this paper, and provides a non perturbative generalization of the commonly used  $\int \tilde{E} = 0$  sum rule introduced in [13]. Indeed, calculating the jet correlator on the perturbative vacuum one would obtain, as already discussed,  $M_q = m_q$  and the integral would vanish.

Finally, the contraction of the hadronic tensor with the leptonic tensor leads to the following result for the inclusive DIS cross section up to order  $\Lambda/Q$  [22]:

$$\frac{d\sigma}{dx_B dy d\phi_S} = \frac{2\alpha^2}{x_B y Q^2} \frac{y^2}{2(1-\epsilon)} \left\{ F_T + \epsilon F_L + S_{\parallel} \lambda_e \sqrt{1-\epsilon^2} F_{LL} + |\mathbf{S}_{\perp}| \lambda_e \sqrt{2\epsilon(1-\epsilon)} \cos \phi_S F_{LT}^{\cos \phi_S} \right\}, \quad (16)$$

where  $\phi_S$  is the angle between the transverse component of the proton spin vector and the lepton plane,  $\epsilon$  is the ratio of the longitudinal and transverse photon fluxes, and  $\lambda_e$  is the electron's helicity. The structure functions on the right hand side *correspond to* *need*

$$F_{TT} = x_B \sum_q e_q^2 f_1^q(x_B), \quad (17)$$

$$F_{TL} = 0, \quad (18)$$

$$F_{LL} = x_B \sum_q e_q^2 g_1^q(x_B), \quad (19)$$

$$F_{TT}^{\sin \phi_S} = 0, \quad (20)$$

$$F_{LT}^{\cos \phi_S} = -x_B \sum_q e_q^2 \frac{2\Lambda}{Q} \left( x_B g_T^q(x_B) + \frac{M_q - m_q}{\Lambda} h_1^q(x_B) \right), \quad (21)$$

where  $f_1^q$ ,  $g_1^q$  and  $h_1^q$  are the unpolarized, polarized, and transversity PDFs, respectively. The second term in the last structure function is a new result from our analysis, *and, although proportional to the jet mass,* *it* is not suppressed as an inverse power of  $Q$  compared to the standard  $g_T$  term. Perturbatively,  $M_q = m_q$  and the new term vanishes. However, on the nonperturbative vacuum the jet mass  $M_q$  is larger than the quark's, originating a nonnegligible term to the twist-3 part of the target's  $g_2$  structure function, as we will discuss in the next section.

#### IV. THE $g_2$ STRUCTURE FUNCTION

The new term in Eq (21) only appears in the  $g_2$  structure function of the proton. Following the derivation in Ref. [18], one finds

$$g_2(x_B) = g_2^{WW}(x_B) + \frac{1}{2} \sum_q e_q^2 \left( \hat{g}_T^{q,*}(x_B) + \int_{x_B}^1 \frac{dy}{y} \hat{g}_T^q(y) + \frac{m_q}{\Lambda} \left( \frac{h_1^q}{x} \right)^*(x_B) + \frac{M_q - m_q}{\Lambda} \frac{h_1^q(x_B)}{x_B} \right), \quad (22)$$

where we defined  $\hat{f}^*(x) = -f(x) + \int_x^1 \frac{dy}{y} f(y)$ . The first four terms coincide with the result obtained in the conventional handbag approximation [18], while the fifth is new. Note that even if the relation is written for the sum over

quark flavors weighted by their charge squared, it is also valid flavor by flavor; in fact, the steps leading to such a decomposition are formulated at the correlator level.

The first term is also known as the Wandzura-Wilczek function,  $g_2^{WW} = -g_1^*$ , and contains all the twist-2 chiral-even contributions to the  $g_2$  structure coming from quark-quark correlators. The second and third terms contain all “pure twist-3” contributions, i.e., those coming from quark-gluon-quark correlators. The fourth and fifth terms contain chiral-odd twist-2 contributions and depend on the transversity distribution function,  $h_1$ . The fourth term is usually neglected for light quarks since it is proportional to  $m_q = O(1 \text{ MeV})$ . The last term, new in our analysis, is again proportional to the transversity distribution but multiplied by a constant of  $O(100 \text{ MeV})$ , and cannot be a priori neglected.

It is important to estimate the size of the various contributions to the non Wandzura-Wilczek part of  $g_2$ . We define the shorthand notation

$$\begin{aligned} g_2^{\text{tw3}} &= \frac{1}{2} \sum_q e_q^2 \left( \tilde{g}_T^{q*}(x_B) + \int_{x_B}^1 \frac{dy}{y} \tilde{g}_T^q(y) \right) \\ g_2^{\text{quark}} &= \frac{1}{2} \sum_q e_q^2 \frac{m_q}{\Lambda} (h_1^q)^*(x_B), \\ g_2^{\text{jet}} &= \frac{1}{2} \sum_q e_q^2 \frac{M_q - m_q}{\Lambda} \frac{h_1^q(x_B)}{x_B}. \end{aligned} \quad (23)$$

These terms are compared in Figure 2 to the  $g_2 - g_2^{WW}$  function obtained in the very recent JAM15 fit of polarized DIS asymmetries [26] that includes a large amount of precise data at large  $x_B$  and small  $Q^2$  from Jefferson Lab, and simultaneously fits the higher-twist components of  $g_1$  and  $g_2$  to the data. For the “pure twist-3” contribution,  $g_2^{\text{tw3}}$ , i.e., the contribution from quark-gluon-quark matrix elements, we show a recent light-front model calculation by Braun et al. [27]; (for  $\pi$ -nucleon bag model calculations, see [28, 29]) To estimate the contributions from quark ( $g_2^{\text{quark}}$ ) and jet mass ( $g_2^{\text{jet}}$ ) effects, that depend on chiral-odd quark-quark matrix elements, we use the recent Pavia15 fit of the transversity distribution from Ref. [10], which is comparable also to other extractions [12, 30]. Furthermore, we choose the values of the mass parameters to be  $m_q = 5 \text{ MeV}$  and  $M_q = 100 \text{ MeV}$ . **[AA: Did we really use 100 MeV in teh plots? I seem to remember 50 MeV.]**

As one can see, in the proton case the pure twist-3 contribution is quite smaller in magnitude, and opposite in sign, compared to the twist-3 term extracted in the JAM15 fit. The quark-mass contribution, as expected, is essentially negligible. For what concerns the jet-mass contribution, the uncertainties due to the  $h_1$  extraction are very large, especially at low  $x_B$ . In addition, there is an overall normalization uncertainty due to the choice of  $M_q$ , not shown in the plot. In any case, the jet-mass contribution is strikingly large. If we assume that the pure twist-3 contributions are of the order of the model calculation by Braun et al., the breaking of the Wandura-Wilczek relation can be used to constrain the extractions of the transversity distribution. This is in particular true at low  $x_B$ , where the pure twist-3 term is expected to vanish. Moreover, it is quite clear that the gap between the pure twist-3  $g_2^{\text{tw3}}$  function and the JAM15 fit can be explained by the new jet-mass contribution we discuss in this paper.

In the neutron case, the jet contribution is very negative at intermediate to large values of  $x_B$ . If one trusts the order of magnitude of the  $g_2^{\text{tw3}}$  calculation by Braun et al., one would conclude that the jet contribution should not be that large. However, this contribution depends strongly on the  $d$  quark’s transversity, whose fit suffers from large systematic uncertainties and saturates the negative Soffer bound. Recent data in  $p + p$  collisions indicate, in fact, that  $h_1^d$  might be less negative than in the Pavia15 fits [31]. Correspondingly the jet contribution to the proton at  $x_B \approx 0.1$  would become less positive, improving as well the agreement with the JAM15 fit.

## V. MOMENTS OF THE $g_2$ STRUCTURE FUNCTION

It is interesting to consider the moments of the non Wandzura-Wilczek contribution to  $g_2$ ,

$$d_N \equiv (N+1) \int_0^1 x^N \left( g_2(x) - g_2^{WW}(x) \right). \quad (24)$$

Keep in mind, however, that  $g_2 - g_2^{WW}$  was fitted assuming the 1st sum rule  $\int_0^1 dx (g_2 - g_2^{WW}) = 0$ , which however is not satisfied after inclusion of jet correlator contributions, as discussed in the next Section. Therefore, ~~the comparison of the fit that goes to 0 for the sum of the 3 other class terms should be done with caution.~~






FIG. 2: Different contributions to the non Wandzura-Wilczek part of the proton (left) and neutron (right)  $g_2$  structure function compared to the JAM15 fit of the  $g_2 - g_2^{\text{WW}}$  (solid black) [26]. The quark and jet contributions are shown with a dotted red and a dot-dashed green line respectively, with uncertainty bands coming from the Pavia15 fit of the transversity function [10]. The uncertainty in the choice  $m_q = 5$  GeV and  $M_q = 100$  GeV is not shown. The pure twist-3 contribution calculated by Braun et al. [27] is shown as a dashed blue line (no uncertainty estimate was provided in the original reference).

For a generic function  $f$ , let us define its  $N$ -th moment as  $f[N] = \int_0^1 dx x^N f(x)$ . It is then straightforward to verify that  $f^*[N] = \frac{1}{N} f[N] N/(N+1)$  and

$$d_N = (N+1)g_2[N] + Ng_1[N] \quad (25)$$

$$= \frac{1}{2} \sum_q e_q^2 \left[ N \tilde{g}_T^q[N] + \hat{g}_T^q[N] \right] \frac{m_q}{\Lambda} \left( \frac{h_1^q}{x} [N] + \frac{(N+1)M_q - m_q}{\Lambda} h_1^q[N-1] \right). \quad (26)$$

The zero-th moment,  $d_0$ , provides an interesting relationship between transversity and the ~~inclusive~~ structure function ~~g2~~.

$$\int dx g_2(x) = \sum_q e_q^2 \frac{M_q - m_q}{\Lambda} \int dx \frac{1}{x} h_1^q(x). \quad (27)$$

Here we used the fact  $f^*[0] = 0$  for any function  $f$ , which eliminates the contribution from the quark-mass and the twist-3  $\tilde{g}_T$  terms; instead,  $\hat{g}_T^q[0]$  vanishes identically due to the symmetry properties of the quark-gluon-quark correlators [18], or equivalently, as a consequence of the Lorentz invariance of QCD interactions, that entails  $\int_0^1 dx g_1^q(x) = \int_0^1 g_T^q(x)$ .

Note that the only surviving term on the right-hand side is the new jet contribution. Thus, our new sum rule (27) generalizes the Burkhardt-Cottingham (BC) sum rule [32], which states that  $\int_0^1 dx g_2(x) = 0$ , while we have shown that jet-mass corrections, and in particular from invariant mass generation in spin-flip processes, can directly violate this. In fact, the possibility of a violation of the BC sum rule due to contributions from spin-flip processes was already mentioned in the original derivation, but these do not show up in treatments that only consider free-field quark propagators for the struck quark [13]. Although we formulated (27) in terms of a sum over quark flavors in order to display a clear connection to the structure function  $g_2$ , we stress that this is valid also flavor by flavor, i.e. for each single flavor the only measurable nonzero contribution to the zeroth moment of the structure function  $g_2$  can come from the coupling between its jet mass and transversity function<sup>1</sup>.

<sup>1</sup> This conclusion is true even if the BC sum rule is broken by a  $J = 0$  fixed pole with non-polynomial residue [13], since this would appear as a  $\delta(x_B)$  contribution and would not be measurable.

One should notice that since  $h_1$  is slowly driven to 0 by QCD evolution as  $Q^2 \rightarrow \infty$ , the BC sum rule is still satisfied at least asymptotically. At finite scales, however, the only way to preserve the validity of the Burkhardt-Cottingham sum rule is if

$$\int dx \frac{1}{x} h_1^q(x) = 0. \quad (28)$$

Interestingly, one can show that this constraint, if valid at any given scale  $Q_0$  is conserved through QCD evolution. However, we think that this constraint is unlikely to be satisfied in general, since it is explicitly broken in perturbative QCD [33], as well as in model calculations [34–39]. Then, if we assume that the BC sum rule is broken by a finite amount, we obtain that  $h_1(x)/x$  must be integrable, implying a bound on the small  $x$  behavior of the transversity,

$$h_1^q(x) \propto x^\epsilon \quad \epsilon > 0. \quad (29)$$

This bound can be very useful, e.g., in transversity fits, where the data at small  $x$  is, as yet, very limited, and in general for proper extrapolations to  $x_B = 0$  when experimentally measuring moments.

The first moment,  $d_1$ , is the first one to display a contribution from the pure twist-3 part of  $g_2$ :

$$d_1 = \frac{1}{2} \sum_q e_q^2 \left( 2\tilde{g}_T^q[1] + \hat{g}_T^q[1] + \frac{2M_q - m_q}{\Lambda} h_1^q[0] \right) \quad (30)$$

where  $h_1^q[0] = \int_0^1 dx h_1^q(x)$  is the contribution of a quark  $q$  to the target's tensor charge. Here, we neglected the quark mass contribution, although its first moment is not identically equal to zero. The second moment,  $d_2$ , is also interesting because the pure twist-3 part can be related to quark-gluon-quark correlators, see [13], and interpreted as the average color force experienced by the struck quark as it exits the nucleon [40]; for experimental measurements of  $d_2$ , see, e.g., Refs. [41–45]. Neglecting the quark mass term contribution, one obtains

$$d_2 = \frac{1}{2} \sum_q e_q^2 \left( 3\tilde{g}_T^q[2] + \hat{g}_T^q[2] + \frac{3M_q - m_q}{\Lambda} h_1^q[1] \right) \quad (31)$$

For both the  $d_1$  and  $d_2$  moments, the transversity contribution is a background to the extraction of the pure twist-3 piece. Fortunately, it is a quantity that can be extracted from the lattice [4–8] **[AA: Check the  $q + \bar{q}$  vs.  $q - \bar{q}$  issue for lattice moments discussed in Oxford]** or fitted [10–12]. Furthermore, the new sum rule (28) and the bound (29) promise to improve future transversity fits. **[Finally, the  $M_q$  jet mass parameter can also be measured, e.g., in electron-positron collisions, as we will briefly discuss later.]** Therefore the pure twist-3 part can, in principle, be properly isolated and measured.

To conclude this section, we note that the jet contribution also leads to an explicit breaking of the Efremov-Teriyayev-Leader (ETL) sum rule [46], in which the pure twist-3 contribution to the first moment of  $g_2 - g_2^{WW}$  also disappears. To see this, let's define the valence contribution to a given structure function as  $f^V = \frac{1}{2} \sum_q e_q^2 (f^q - f^{\bar{q}})$ . Then, as shown in [46],  $\tilde{g}_T^V[1] + \hat{g}_T^V[1] = 0$ , and from Eq. (30) we obtain

$$d_1^V = \frac{1}{2} \sum_q e_q^2 \frac{2M_q - m_q}{\Lambda} (h_1^q[0] - h_1^{\bar{q}}[0]). \quad (32)$$

Assuming  $M_{\text{light}} \equiv M_u \approx M_d \gg m_u, m_d$  and isospin symmetry of the proton and neutron, we can also see that

$$d_1^V = \frac{M_{\text{light}}}{\Lambda} \delta_T(p), \quad (33)$$

This gives an alternative way to access the proton tensor charge,  $\delta_T(p) = \sum_q e_q^2 (h_1^q[0] - h_1^{\bar{q}}[0])$ , by measuring moments of the flavor separated  $g_2$  structure function. **[AA: I tried to build an observable for  $d_1^V$  such as  $p - n$  or  $p - \bar{p}$ , but I am messing up a bit.]**

## VI. MEASURING THE INTEGRATED JET FCFS

To experimentally measure jet functions, and in particular the jet mass parameter  $M_q$ , a promising avenue is through inclusive single hadron production,  $e^+e^- \rightarrow hX$ , and inclusive dihadron production from the same hemisphere,  $e^+e^- \rightarrow hhX$ , see Fig. 3. **[AA: Andrea suggested to remove teh figures, to avoid competitors stealing our**

idea and doing the actual analysis of the observables sensitive to  $M_q$ . I see his point, but am a bit torn about this.] In single-hadron production, the fragmentation functions  $\Delta^h$  play the role of PDFs in DIS, and couple to the jet functions in an analogous way. To access the spin-flip  $J_1$  function one needs to detect a polarized hadron, such as a  $\Delta$  baryon. In double hadron production, the enlarged number of Dirac structures of the dihadron fragmentation correlators [47, 48] allows one to access the jet function in novel ways, and in particular to isolate the contribution from the helicity-flip  $J_1$  term in combination with the chiral-odd fragmentation function  $H_1^\perp$ .

Studying and classifying all the possibilities offered by single and double hadron production in electron-positron annihilation events will open up a rich phenomenology, which will in turn be needed to extract pure twist-3 matrix elements from the  $g_2$  DIS structure function. While measurements in the asymptotically large  $Q^2$  regime will provide access to the integral of the  $J_1$  jet function, i.e., to the jet-mass parameter  $M_q$ , perhaps even more interesting is the possibility to experimentally measure at finite values of  $Q^2$  the momentum dependence of the jet functions  $J_1$  and  $J_2$ , that enter structure functions in a way analogous to that discussed in Ref. [20]. In other words, it will become possible to experimentally access the quark's spectral function.




FIG. 3: Single hadron (left) and double hadron (right) production in  $e^+e^-$  collisions at LO with jet and fragmentation correlators.

## VII. CONCLUSIONS

In this paper, we revisited the inclusive DIS analysis, including the effects due to the production of a system of final state hadrons in the current direction, which we conveniently referred to as a “jet.” We described this in terms of a jet correlator that corresponds, up to twist-4 contributions, to the nonperturbative quark cut propagator, or, equivalently, to the quark’s spectral function. We then carried out the analysis of the DIS cross section up to twist-3 level. The introduction of this jet correlator leads to a difference in the expression of the structure function  $g_2$  in inclusive DIS with respect to the standard analysis: a new term appears, proportional to a jet mass parameter and to the transversity distribution function. This new term contributes to the violation of the Wandzura-Wilczek relation, in addition to the standard pure twist-3 terms and quark mass corrections. Contrary to these standard terms, however, the new jet mass correction does not necessarily integrate to zero and so violates also the Burkhardt-Cottingham sum rule. This is yet another example of how surprising and rich the phenomenology of polarized inclusive DIS can be. Detailed measurements of the  $g_2$  structure function can be used to constrain the jet mass parameter, the transversity distribution function and the nucleon tensor charge, helping their extraction from other observables in, e.g., electron-positron annihilation and semi-inclusive DIS, respectively. The knowledge of jet mass parameter and transversity distribution will be eventually needed for a precise extraction of pure twist-3 terms from the structure function  $g_2$ . [AA: We should try and make the conclusions a little crisper.]

## Acknowledgments

We are grateful to A. Signori for a careful reading of a draft of this article, and to A. Mukherjee, M. Sattman, and W. Vogelsang for interesting discussions. This work was supported by the United States Department of Energy (DOE) contract No. DE-AC05-06OR23177, under which Jefferson Science Associates, LLC operates Jefferson Lab, by the DOE contract DE-SC008791, and by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No. 647981, 3DSPIN).

[AA: NOTE: the umlauts do not show up in the bibliography!! See, e.g., reference [5]. Need to fix



this.]

- 
- [1] V. Cirigliano, S. Gardner, and B. Holstein, *Prog. Part. Nucl. Phys.* **71**, 93 (2013).
  - [2] T. Bhattacharya, V. Cirigliano, R. Gupta, H.-W. Lin, and B. Yoon, *Phys. Rev. Lett.* **115**, 212002 (2015).
  - [3] A. Courtoy, S. Baeler, M. Gonzalez-Alonso, and S. Liuti, *Phys. Rev. Lett.* **115**, 162001 (2015).
  - [4] J. R. Green, J. W. Negele, A. V. Pochinsky, S. N. Syritsyn, M. Engelhardt, and S. Krieg, *Phys. Rev.* **D86**, 114509 (2012).
  - [5] G. S. Bali, S. Collins, B. Glssle, M. Gckeler, J. Najjar, R. H. Rdl, A. Schfer, R. W. Schiel, W. Sldner, and A. Sternbeck, *Phys. Rev.* **D91**, 054501 (2015).
  - [6] T. Bhattacharya, V. Cirigliano, S. Cohen, R. Gupta, A. Joseph, H.-W. Lin, and B. Yoon (PNDME), *Phys. Rev.* **D92**, 094511 (2015).
  - [7] A. Abdel-Rehim et al., *Phys. Rev.* **D92**, 114513 (2015), [Erratum: *Phys. Rev.* **D93**, no.3, 039904(2016)].
  - [8] T. Bhattacharya, V. Cirigliano, S. Cohen, R. Gupta, H.-W. Lin, and B. Yoon (2016), arXiv:1606.07049 [hep-lat].
  - [9] V. Barone, A. Drago, and P. G. Ratcliffe, *Phys. Rept.* **359**, 1 (2002).
  - [10] M. Radici, A. Courtoy, A. Bacchetta, and M. Guagnelli, *JHEP* **05**, 123 (2015).
  - [11] M. Anselmino, M. Boglione, U. D'Alesio, J. O. Gonzalez Hernandez, S. Melis, F. Murgia, and A. Prokudin, *Phys. Rev.* **D92**, 114023 (2015).
  - [12] Z.-B. Kang, A. Prokudin, P. Sun, and F. Yuan, *Phys. Rev.* **D93**, 014009 (2016).
  - [13] R. L. Jaffe (1996), arXiv:hep-ph/9602236 [hep-ph], [Lect. Notes Phys.496,178(1997)].
  - [14] J. P. Ralston and D. E. Soper, *Nucl. Phys.* **B152**, 109 (1979).
  - [15] R. L. Jaffe and X.-D. Ji, *Phys. Rev. Lett.* **67**, 552 (1991).
  - [16] R. L. Jaffe and X.-D. Ji, *Phys. Rev. Lett.* **71**, 2547 (1993).
  - [17] J. C. Collins, *Nucl. Phys.* **B396**, 161 (1993).
  - [18] A. Accardi, A. Bacchetta, W. Melnitchouk, and M. Schlegel, *JHEP* **11**, 093 (2009).
  - [19] S. Wandzura and F. Wilczek, *Phys. Lett.* **B72**, 195 (1977).
  - [20] A. Accardi and J.-W. Qiu, *JHEP* **07**, 090 (2008).
  - [21] J. C. Collins, T. C. Rogers, and A. M. Stasto, *Phys. Rev.* **D77**, 085009 (2008).
  - [22] A. Bacchetta, M. Diehl, K. Goeke, A. Metz, P. J. Mulders, and M. Schlegel, *JHEP* **02**, 093 (2007).
  - [23] E. D'Hoker, *Quantum Field Theory - Part 1* (2004), URL [http://www.pa.ucla.edu/sites/default/files/files/dhoker%20lecture%20notes/quantum\\_field\\_theory.pdf](http://www.pa.ucla.edu/sites/default/files/files/dhoker%20lecture%20notes/quantum_field_theory.pdf).
  - [24] J. C. Romão, *Advanced quantum field theory - Part 1* (2013), URL <http://porthos.ist.utl.pt/ftp/textos/tca.pdf>.
  - [25] S. Weinberg, *The Quantum theory of fields. Vol. 1: Foundations* (Cambridge University Press, 2005), ISBN 9780521670531, 9780511252044.
  - [26] N. Sato, W. Melnitchouk, S. E. Kuhn, J. J. Ethier, and A. Accardi (Jefferson Lab Angular Momentum), *Phys. Rev.* **D93**, 074005 (2016).
  - [27] V. M. Braun, T. Lautenschlager, A. N. Manashov, and B. Pirnay, *Phys. Rev.* **D83**, 094023 (2011).
  - [28] R. L. Jaffe and X.-D. Ji, *Phys. Rev.* **D43**, 724 (1991).
  - [29] M. Stratmann, *Z. Phys.* **C60**, 763 (1993).
  - [30] M. Anselmino, M. Boglione, U. D'Alesio, S. Melis, F. Murgia, and A. Prokudin, *Phys. Rev.* **D87**, 094019 (2013).
  - [31] M. Radici, A. M. Ricci, A. Bacchetta, and A. Mukherjee (2016), arXiv:1604.06585 [hep-ph].
  - [32] H. Burkhardt and W. N. Cottingham, *Annals Phys.* **56**, 453 (1970).
  - [33] R. Kundu and A. Metz, *Phys. Rev.* **D65**, 014009 (2002).
  - [34] P. Schweitzer, D. Urbano, M. V. Polyakov, C. Weiss, P. V. Pobylitsa, and K. Goeke, *Phys. Rev.* **D64**, 034013 (2001).
  - [35] M. Wakamatsu, *Phys. Lett.* **B653**, 398 (2007).
  - [36] B. Pasquini, M. Pincetti, and S. Boffi, *Phys. Rev.* **D72**, 094029 (2005).
  - [37] I. C. Cloet, W. Bentz, and A. W. Thomas, *Phys. Lett.* **B659**, 214 (2008).
  - [38] A. Bacchetta, F. Conti, and M. Radici, *Phys. Rev.* **D78**, 074010 (2008).
  - [39] C. Bourrely, F. Buccella, and J. Soffer, *Phys. Rev.* **D83**, 074008 (2011).
  - [40] M. Burkardt, *Phys. Rev.* **D88**, 014014 (2013).
  - [41] P. L. Anthony et al. (E155), *Phys. Lett.* **B553**, 18 (2003).
  - [42] K. Slifer et al. (Resonance Spin Structure), *Phys. Rev. Lett.* **105**, 101601 (2010).
  - [43] P. Solvignon et al. (E01-012), *Phys. Rev.* **C92**, 015208 (2015).
  - [44] M. Posik et al. (Jefferson Lab Hall A), *Phys. Rev. Lett.* **113**, 022002 (2014).
  - [45] D. Flay et al. (2016), arXiv:1603.03612 [nucl-ex].
  - [46] A. V. Efremov, O. V. Teryaev, and E. Leader, *Phys. Rev.* **D55**, 4307 (1997).
  - [47] A. Bacchetta and M. Radici, *Phys. Rev.* **D67**, 094002 (2003).
  - [48] A. Bacchetta and M. Radici, *Phys. Rev.* **D69**, 074026 (2004).