

## ECE 222 Homework #2

Scan or photograph and save as a single .pdf file. Name your file "Lastname\_HW02.pdf"

Do the following problems (in bold):

1) Concepts of line, surface, and volume charge density

**4.1, 4.2, 4.4, 4.5, 4.8(a), 4.9(a).** These help you become familiar with the concepts of line, surface, and volume charge density, as well as practice volume integrals in various coordinates, which you will regularly use in the rest of the class. *See pages 6 & 11*

2) Calculation of electric field based on Coulomb's law and the principle of superposition

**4.12, 4.16, 4.17, 4.19.** Each of these will require you to draw the distribution of charges that are the sources of electric field, the electric field vector that stems from each source, and add them up *See pages 11 & 12*

Useful integral for 4.16:

$$\int_{-L/2}^{L/2} \frac{dz_0}{(r^2 + z_0^2)^{3/2}} = \frac{1}{r^2} \frac{L}{\sqrt{r^2 + (L/2)^2}}$$

3) Calculation of electric field based on integral form of Gauss's law

**3.50, 4.22, 4.25, 4.26, 4.27, 4.29.** Coulomb's law is often not the best way to calculate the electric field stemming from a given charge distribution. In cases where the charge distribution has a clear symmetry (cylinder, sphere, plane, etc.), the use of Gauss's law (which, in its integral form, equates the surface integral of  $\vec{D}$  through an appropriately chosen surface with the volume integral of charge density  $\rho_v$ ) is the optimal route to calculate electric field everywhere. *See pages 12 & 13*

\*4.1 A cube 2 m on a side is located in the first octant in a Cartesian coordinate system with one of its corners at the origin. Find the total charge contained in the cube if the charge density is given by  $\rho_v = xy^2e^{-2z}$  (mC/m<sup>3</sup>).

4.2 Find the total charge contained in a cylindrical volume defined by  $r \leq 2$  m and  $0 \leq z \leq 3$  m if  $\rho_v = 20rz$  (mC/m<sup>3</sup>).

4.4 If the line charge density is given by  $\rho_l = 24y^2$  (mC/m), find the total charge distributed on the  $y$  axis from  $y = -5$  to  $y = 5$ .

4.5 Find the total charge on a circular disk defined by  $r \leq a$  and  $z = 0$  if:

- a.  $\rho_s = \rho_{s0} \cos \phi$  (C/m<sup>2</sup>)
- b.  $\rho_s = \rho_{s0} \sin^2 \phi$  (C/m<sup>2</sup>)
- c.  $\rho_s = \rho_{s0} e^{-r}$  (C/m<sup>2</sup>)
- d.  $\rho_s = \rho_{s0} e^{-r} \sin^2 \phi$  (C/m<sup>2</sup>)

where  $\rho_{s0}$  is a constant.

4.8 An electron beam shaped like a circular cylinder of radius  $r_0$  carries a charge density given by

$$\rho_v = \left( \frac{-\rho_0}{1+r^2} \right) \text{ (C/m}^3\text{)}$$

where  $\rho_0$  is a positive constant and the beam's axis is coincident with the  $z$  axis.

- a. Determine the total charge contained in length  $L$  of the beam.
- b. If the electrons are moving in the  $+z$  direction with uniform speed  $u$ , determine the magnitude and direction of the current crossing the  $z$ -plane.

4.9 A circular beam of charge of radius  $a$  consists of electrons moving with a constant speed  $u$  along the  $+z$  direction. The beam's axis is coincident with the  $z$  axis and the electron charge density is given by

$$\rho_v = -cr^2 \quad (\text{C/m}^3)$$

where  $c$  is a constant and  $r$  is the radial distance from the axis of the beam.

- \*a. Determine the charge density per unit length.
- b. Determine the current crossing the  $z$ -plane.

- 4.12 Three point charges, each with  $q = 3 \text{ nC}$ , are located at the corners of a triangle in the  $x$ - $y$  plane, with one corner at the origin, another at  $(2 \text{ cm}, 0, 0)$ , and the third at  $(0, 2 \text{ cm}, 0)$ . Find the force acting on the charge located at the origin.
- 4.16 A line of charge with uniform density  $\rho_l$  extends between  $z = -L/2$  and  $z = L/2$  along the  $z$  axis. Apply Coulomb's law to obtain an expression for the electric field at any point  $P(r, \phi, 0)$  on the  $x$ - $y$  plane. Show that your result reduces to the expression given by Eq. (4.33) as the length  $L$  is extended to infinity.
- \*4.17 Repeat Example 4-5 for the circular disk of charge of radius  $a$ , but in the present case, assume the surface charge density to vary with  $r$  as

$$\rho_s = \rho_{s0} r^2 \quad (\text{C/m}^2)$$


where  $\rho_{s0}$  is a constant.

- \*4.19 Three infinite lines of charge, all parallel to the  $z$  axis, are located at the three corners of the kite-shaped arrangement shown in Fig. P4.19. If the two right triangles are symmetrical and of equal corresponding sides, show that the electric field is zero at the origin.

Figure P4.19 Kite-shaped arrangement of line charges for Problem 4.19.

$$\therefore \cos \theta = 1/2$$

$$\begin{aligned} d^2 &= x^2 + y^2 \\ \vec{S}_O \cdot \vec{S}_I &= 0 \\ [-x, -y] \cdot [-x, H-y] &= 0 \\ &= x^2 - y(H-y) = x^2 + y^2 - Hy = 0 \\ \therefore d^2 &= x^2 + y^2 = Hy \end{aligned}$$



See next page for Eq. (4.33) and Example 4-5 for problems 4.16 and 4.17

(4.33)


$$\mathbf{E} = \frac{\mathbf{D}}{\epsilon_0} = \hat{\mathbf{r}} \frac{D_r}{\epsilon_0} = \hat{\mathbf{r}} \frac{\rho_\ell}{2\pi\epsilon_0 r}.$$

**(infinite line charge)**

### Example 4-5: Electric Field of a Circular Disk of Charge

Find the electric field at point  $P$  with Cartesian coordinates  $(0, 0, h)$  due to a circular disk of radius  $a$  and uniform charge density  $\rho_s$  residing in the  $x$ - $y$  plane (Fig. 4-7). Also, evaluate  $\mathbf{E}$  due to an infinite sheet of charge density  $\rho_s$  by letting  $a \rightarrow \infty$ .

Figure 4-7 Circular disk of charge with surface charge density  $\rho_s$ . The electric field at  $P = (0, 0, h)$  points along the  $z$  direction (Example 4-5).



**Solution:** Building on the expression obtained in [Example 4-4](#) for the on-axis electric field due to a circular ring of charge, we can determine the field due to the circular disk by treating the disk as a set of concentric rings. A ring of radius  $r$  and width  $dr$  has an area  $ds = 2\pi r dr$  and contains charge  $dq = \rho_s ds = 2\pi\rho_s r dr$ . Upon using this expression in [Eq. \(4.23\)](#) and also replacing  $b$  with  $r$ , we obtain the following expression for the field due to the ring:

$$d\mathbf{E} = \hat{\mathbf{z}} \frac{h}{4\pi\epsilon_0(r^2 + h^2)^{3/2}} (2\pi\rho_s r dr).$$

The total field at  $P$  is obtained by integrating the expression over the limits  $r = 0$  to  $r = a$ :

(4.24)

$$\mathbf{E} = \hat{\mathbf{z}} \frac{\rho_s h}{2\epsilon_0} \int_0^a \frac{r dr}{(r^2 + h^2)^{3/2}} = \pm \hat{\mathbf{z}} \frac{\rho_s}{2\epsilon_0} \left[ 1 - \frac{|h|}{\sqrt{a^2 + h^2}} \right],$$

with the plus sign for  $h > 0$  ( $P$  above the disk) and the minus sign when  $h < 0$  ( $P$  below the disk).


For an infinite sheet of charge with  $a = \infty$ ,

(4.25)

$$\mathbf{E} = \pm \hat{\mathbf{z}} \frac{\rho_s}{2\epsilon_0} .$$

**(infinite sheet of charge)**

We note that for an infinite sheet of charge  $\mathbf{E}$  is the same at all points above the  $x$ - $y$  plane, and a similar statement applies for points below the  $x$ - $y$  plane.



- \*3.50 A vector field  $\mathbf{D} = \hat{\mathbf{r}}r^3$  exists in the region between two concentric cylindrical surfaces defined by  $r = 1$  and  $r = 2$  with both cylinders extending between  $z = 0$  and  $z = 5$ .

Verify the divergence theorem by evaluating the following:  $SA = 2\pi r \cdot H$

$$\begin{aligned} \text{a. } \oint_S \mathbf{D} \cdot d\mathbf{s} &= 2\pi(r=2)(5) \cdot 2^3 - 2\pi(r=1)(5) \cdot 2^1 = 150\pi \\ \text{b. } \int_V \nabla \cdot \mathbf{D} dv &= \nabla \cdot \mathbf{D} = \frac{1}{r} \frac{d(rD_r)}{dr} = \frac{1}{r} 4r^3 = 4r^2 \cdot \int_0^5 \int_0^{2\pi} \int_1^2 4r^2 dr d\phi dz = [2^4 - 1^4] 10\pi = 150\pi \end{aligned}$$

- 4.22 Given the electric flux density

$$\mathbf{D} = \hat{\mathbf{x}}2(x+y) + \hat{\mathbf{y}}(3x-2y) \quad (\text{C/m}^2),$$

determine

$$\rho_v = \frac{\partial}{\partial x} [2(x+y)] + \frac{\partial}{\partial y} (3x-2y) = 2 - 2 = 0$$

- a.  $\rho_v$  by applying Eq. (4.26)  $\square$ ,

- b. the total charge  $Q$  enclosed in a cube 2 m on a side, located in the first octant with three of its sides coincident with the  $x$ ,  $y$ , and  $z$  axes and one of its corners at the origin, and

- c. the total charge  $Q$  in the cube, obtained by applying Eq. (4.29)  $\square$ .

Eq. (4.26):

$$\begin{aligned} \text{face: } x=2 &\rightarrow \iint_A \mathbf{D} \cdot \hat{\mathbf{n}} dA = \int_0^2 \int_0^2 D_x dz dy = \int_0^2 \int_0^2 (4+2y) dz dy = \int_0^2 [8+4y] dy = 16 + 2(2^2) = 24 \\ \text{face: } z=2 &\rightarrow 0 \end{aligned}$$

(4.26)

$$\text{face: } x=0 \rightarrow \iint_A \mathbf{D} \cdot \hat{\mathbf{n}} dA = -\int_0^2 \int_0^2 D_x dz dy = -\int_0^2 \int_0^2 2y dz dy = -2y^2 \Big|_0^2 = -8$$

(differential form of Gauss's law)

$$\begin{aligned} \text{face: } y=2 &\rightarrow \iint_A \mathbf{D} \cdot \hat{\mathbf{n}} dA = \int_0^2 \int_0^2 D_y dx dz = \int_0^2 \int_0^2 (3x-4) dx dz = \int_0^2 [3x^2 - 4x]_0^2 dz = [3(2^2) - 8] \Big|_0^2 = -4 \\ \text{face: } z=0 &\rightarrow 0 \end{aligned}$$

Eq. (4.29):

$$\text{face: } y=0 \rightarrow \iint_A \mathbf{D} \cdot \hat{\mathbf{n}} dA \rightarrow -\int_0^2 \int_0^2 D_y dz dx = -\int_0^2 6x dx = -3x^2 \Big|_0^2 = -12$$

(4.29)

$$\oint_S \mathbf{D} \cdot d\mathbf{s} = Q.$$

(integral form of Gauss's law)

$$\therefore \oint_S \mathbf{D} \cdot d\mathbf{s} = Q = 24 - 8 - 12 - 4 = 0$$

- \*4.25 The electric flux density inside a dielectric sphere of radius  $a$  centered at the origin is given by

$$\mathbf{D} = \hat{\mathbf{R}}\rho_0 R \quad (\text{C/m}^2)$$

See Page 12

where  $\rho_0$  is a constant. Find the total charge inside the sphere.

- 4.26 In a certain region of space, the charge density is given in cylindrical coordinates by the function

$$\rho_v = 5re^{-r} \quad (\text{C/m}^3).$$

Apply Gauss's law to find  $\mathbf{D}$ .

- \*4.27 An infinitely long cylindrical shell extending between  $r = 1 \text{ m}$  and  $r = 3 \text{ m}$  contains a uniform charge density  $\rho_{v0}$ . Apply Gauss's law to find  $\mathbf{D}$  in all regions.


- 4.29 A spherical shell with outer radius  $b$  surrounds a charge-free cavity of radius  $a < b$  (Fig. P4.29). If the shell contains a charge density given by

$$\rho_v = -\frac{\rho_{v0}}{R^2}, \quad a \leq R \leq b,$$

See Pg. 13

where  $\rho_{v0}$  is a positive constant, determine  $\mathbf{D}$  in all regions.

Figure P4.29 Problem 4.29.



**Solutions to starred problems:**

4.1  $Q = 2.62 \text{ (nC)}$

4.17  $\mathbf{E} = \hat{\mathbf{z}} (\rho_{s0} h / 2\epsilon_0) [\sqrt{a^2 + h^2} + h^2 / \sqrt{a^2 + h^2} - 2h]$

4.19  $\mathbf{E} = 0$

$$3.50 \blacksquare \quad (\text{a}) \oint \mathbf{D} \cdot d\mathbf{s} = 150\pi$$


$$(\text{b}) \iiint \nabla \cdot \mathbf{D} dv = 150\pi$$

$$4.25 \blacksquare \quad Q = 4\pi\rho_0 a^3 \text{ (C)}$$

$$4.27 \blacksquare \quad \mathbf{D} = \hat{\mathbf{r}} \frac{\rho_{v0}(r^2 - 1)}{2r}, \quad 1 \leq r \leq 3 \text{ m}$$

$$\mathbf{D} = \hat{\mathbf{r}} D_r = \hat{\mathbf{r}} \frac{4\rho_{v0}}{r}, \quad r \geq 3 \text{ m}$$

4.1:  $\rho = xy^2 e^{-2z} \text{ (mC/m}^3\text{)}$



$$Q = \int_0^2 \int_0^2 \int_0^2 \rho \, dx \, dy \, dz; I_x = \int_0^2 \rho \, dx = \int_0^2 xy^2 e^{-2z} \, dx = \left[ \frac{x^2}{2} y^2 e^{-2z} \right]_0^2 = 2y^2 e^{-2z}$$

$$I_y = \int_0^2 2y^2 e^{-2z} \, dy = \left[ \frac{2}{3} y^3 e^{-2z} \right]_0^2 = \frac{8}{3} e^{-2z}; Q = I_z = \int_0^2 \frac{8}{3} e^{-2z} \, dz = \left[ -\frac{8}{6} e^{-2z} \right]_0^2$$

$$Q = \frac{8}{3} (1 - e^{-4}) \approx 2.618 \text{ mC}$$

4.2.  $\rho_r = 20r^2 z \text{ (nC/m}^3\text{)}$ ;  $\int_0^3 \int_0^{2\pi} \int_0^2 20r^2 z \, dr \, d\phi \, dz$ ;  $I_r = \left[ \frac{20}{3} r^3 z \right]_0^2 = \frac{160}{3} z$ ;

$$I_\phi = \frac{320\pi}{3} \cdot z; Q = I_z = \left[ \frac{320}{6} \cdot z^2 \right]_0^3 = \frac{480\pi}{1000} \text{ C} = \frac{12}{25} \pi \text{ C}$$

4.4.  $\rho_z = 24y^2 \text{ (nC/m)}$ ;  $Q = \int_{-5}^5 24y^2 \, dy = \left[ 8y^3 \right]_{-5}^5 = \left[ 8(5^3 - (-5)^3) \right] = 20$

4.5 a.  $Q = \rho_{s0} \int_0^{2\pi} \int_0^a r \cos \phi \, dr \, d\phi$ ;  $I_r = \left[ \frac{r^2}{2} \cos \phi \right]_0^a$ ;  $Q = \rho_{s0} \int_0^{2\pi} \frac{a^2}{2} \cos \phi \, d\phi$

$$Q = \rho_{s0} \left[ -\frac{a^2}{2} \sin \phi \right]_0^{2\pi} = 0$$

b.  $Q = \int_0^{2\pi} \int_0^a \sin^2 \phi \, r \, dr \, d\phi \left( \frac{C}{m} \right)$ ;  $I_r = \left[ \frac{r^2}{2} \sin^2 \phi \right]_0^a = \frac{a^2}{2} \sin^2 \phi$ ;  $I_\phi = \frac{a^2}{2} \rho_{s0} \int_0^{2\pi} \sin^2 \phi \, d\phi$

$$\int_0^{2\pi} \sin^2 \phi \, d\phi = \int_0^{2\pi} \frac{1 - \cos(2\phi)}{2} \, d\phi = \int_0^{2\pi} \frac{1}{2} \, d\phi - \int_0^{2\pi} \frac{\cos(2\phi)}{2} \, d\phi = \pi - \left[ \frac{\sin(2\phi)}{4} \right]_0^{2\pi}$$

$$Q = \frac{\pi a^2 \rho_{s0}}{2} \text{ C}$$

c.  $Q = \rho_{s0} \int_0^{2\pi} \int_0^a r e^{-r} \, dr \, d\phi$ ;  $u = r; du = dr$ ;  $\int_0^a r e^{-r} \, dr = -r e^{-r} + \int e^{-r} \, dr = \left[ -e^{-r} (r+1) \right]_0^a = -e^{-a} (a+1) + 1$

$$Q = \rho_{s0} (1 - e^{-a} (a+1)) \int_0^{2\pi} d\phi; Q = 2\pi \rho_{s0} (1 - e^{-a} (a+1)) \text{ C}$$

d.  $Q = \rho_{s0} \int_0^{2\pi} \int_0^a r e^{-r} \sin^2 \phi \, dr \, d\phi = \rho_{s0} \pi (1 - e^{-a} (a+1)) \text{ C}$

$$4.8 \text{ a. } \rho_r = \left( \frac{-\rho_0}{1+r^2} \right) \text{ C/m}^3; Q = 2\pi \cdot L \int_0^{r_0} \frac{-\rho_0 \cdot r}{1+r^2} dr \text{ C/m}^3$$

$$-2\pi\rho_0 L \int_0^{r_0} \frac{r}{r^2+1} dr; u=r^2+1; du=2r dr; dr=\frac{du}{2r}; -\pi\rho_0 L \int_0^{r_0} \frac{1}{u} du = -\pi\rho_0 L \left[ \ln|r^2+1| \right]_0^{r_0}$$

$$Q = -\pi\rho_0 L (\ln|r_0^2+1|) \text{ C}$$

$$\text{b. } \mathbf{J} = \rho_0 u \hat{z}; \mathbf{I} = \iint_S \mathbf{J} \cdot d\mathbf{S}; d\mathbf{S} = \hat{z} r dr d\phi; \mathbf{I} = \int_0^{2\pi} \int_0^{r_0} \frac{-\rho_0 u}{1+r^2} r dr d\phi$$

$$\mathbf{I} = -\pi\rho_0 u \ln(1+r_0^2) \hat{z}; |\mathbf{I}| = \rho_0 u \ln(1+r_0^2); \mathbf{D} = -\hat{z}$$

$$4.9: \rho_r = -cr^2 (\text{C/m}^3); dQ = \rho_r dV; \therefore dV = (2\pi r dr dz); \therefore \lambda = \frac{dQ}{dz} = \int_0^a -2\pi cr^3 dr dz$$

$$\lambda = -2\pi c \int_0^a r^3 dr = -\frac{c}{2} \pi a^4 \text{ C/m}$$


$$\hat{z}$$

$$\therefore A = \pi r^2; dA = 2\pi r dr$$

$$\therefore dV = 2\pi r dr dz$$

$$\text{b. } \mathbf{I} = 2u = -\frac{c}{2} \pi a^4 u \hat{z}$$

$$4.12: q = 3nC; @ (0,0,0); \mathbf{E} = \frac{Q}{4\pi\epsilon_0 R^2} \hat{R}; \mathbf{E}_{\text{origin}} = \frac{q}{4\pi\epsilon_0 r^2} (\hat{x} + \hat{y}); \mathbf{F}_{\text{origin}} = \mathbf{E}_{\text{origin}} \cdot q;$$




$$\mathbf{F} = \frac{q^2}{4\pi\epsilon_0 r^2} (\hat{x} + \hat{y}) = -2.023 \times 10^{-4} (\hat{x} + \hat{y}) \text{ N}; |\mathbf{F}| = 2.86 \times 10^{-4} \text{ N}$$

$$4.16: \text{Uniform } \rho_L; \mathbf{R} = r\hat{r} - z\hat{z}; |\mathbf{R}| = \sqrt{r^2 + z^2}; dE = \frac{1}{4\pi\epsilon_0} \frac{dQ R}{|\mathbf{R}|^3} = \frac{\rho_L}{4\pi\epsilon_0} \left( \frac{r\hat{r} - z\hat{z}}{(r^2 + z^2)^{3/2}} \right) dz$$


$$\mathbf{E} = \hat{r} \frac{\rho_L r}{4\pi\epsilon_0} \int_{-L/2}^{L/2} \frac{dz}{(r^2 + z^2)^{3/2}} = \frac{L}{r^2 \sqrt{r^2 + (L/2)^2}} \hat{r} \frac{\rho_L r}{4\pi\epsilon_0}$$

$$\mathbf{E} = \frac{\rho_L L}{4\pi\epsilon_0 \sqrt{r^2 + (L/2)^2}} \hat{r}; \lim_{L \rightarrow \infty} \frac{L}{\sqrt{r^2 + (L/2)^2}} = \lim_{L \rightarrow \infty} \frac{L}{\frac{L}{2} \sqrt{1 + 4r^2/L^2}} = 2$$

$$\therefore \mathbf{E} = \frac{\rho_L}{2\pi\epsilon_0} \hat{r} \text{ as } L \rightarrow \infty$$




$$4.17. P_s = P_{s0} r^2 \text{ (C/m}^2\text{)}; dq = P_s dA = (P_{s0} r^2) (2\pi r dr) = 2\pi P_{s0} r^3 dr; dE_z = \frac{1}{4\pi\epsilon_0} \left( \frac{h dq}{(r^2 + h^2)^{3/2}} \right);$$



$$E_z = \frac{P_{s0} h}{2\epsilon_0} \int_0^a \frac{r^3}{(r^2 + h^2)^{3/2}} dr; U = r^2 + h^2; dU = 2r dr; r dr = \frac{dU}{2}; r^2 = U - h^2; \text{ thus:}$$

$$\int \frac{r^2(r dr)}{(r^2 + h^2)^{3/2}} = \frac{1}{2} \int \frac{U - h^2}{U^{3/2}} dU = \frac{1}{2} \left( \int \frac{1}{\sqrt{U}} dU - h^2 \int \frac{1}{U^{3/2}} dU \right) = \sqrt{U} + \frac{h^2}{\sqrt{U}} = \sqrt{r^2 + h^2} + \frac{h^2}{\sqrt{r^2 + h^2}}$$

for  $h > 0$ :  $E(0,0,h) = \frac{P_{s0} h}{2\epsilon_0} \left[ \sqrt{a^2 + h^2} + \frac{h^2}{\sqrt{a^2 + h^2}} - 2h \right]$

$$4.25. D = \hat{r} P_0 R \text{ (C/m}^2\text{)}; SA = 4\pi R^2; Q = \oint_S D \cdot d\mathbf{s} = (P_0 \cdot a) (4\pi a^2) = P_0 a^3 4\pi$$


$$4.26. P_v = 5re^{-r} \text{ (C/m}^3\text{)}; \iiint_V P_v dV = Q; \int_0^L \int_0^{2\pi} \int_0^R 5r e^{-r} dr d\phi dz = Q = 10\pi L \int_0^R r^2 e^{-r} dr$$

$$\therefore D(R) (2\pi RL) = 10\pi L \int_0^R r^2 e^{-r} dr; D(R) = \frac{5}{R} \int_0^R r^2 e^{-r} dr; \int u dv = uv - \int v du; u = r^2, du = 2r dr; dv = e^{-r}, v = -e^{-r}$$

$$= -r^2 e^{-r} + 2 \int r e^{-r} dr; u = r, du = dr; \int r e^{-r} dr = -r e^{-r} - \int -e^{-r} dr = -r e^{-r} - e^{-r}$$

$$= -e^{-r} (r^2 + 2r + 2)$$

$$4.27. 1m \leq R \leq 3m; P_{v0}; D(R) (2\pi RL) = Q_{enc}$$

$$R < 1 \text{ No } Q_{enc} \rightarrow D = 0$$


$$1 \leq R \leq 3: Q_{enc} = P_{v0} \pi (R^2 - 1) L; D(R) = \frac{P_{v0} (R^2 - 1)}{2R} \hat{r}$$

$$R \geq 3: Q_{enc} = 8\pi P_{v0} L; D(R) = \frac{4P_{v0}}{R} \hat{r}$$

$$D(R) = \frac{5}{R} \left[ -e^{-r} (r^2 + 2r + 2) \right]_0^R$$

$$= \frac{5}{R} \left[ 2 - e^{-R} (R^2 + 2R + 2) \right] \hat{r}$$

4.29:  $D_R(R)(4\pi R^2) = Q_{enc.}$ ;  $R < a$ :  $Q_{enc} = 0 = D$ ;  $a \leq R \leq b$ :  $\rightarrow$



$$Q_{enc} = \int_a^R \left(-\frac{\rho_0}{s^2}\right) 4\pi s^2 ds = -4\pi \rho_0 (R-a) \rightarrow D(R) = \frac{\rho_0 (a-R)}{R^2} \hat{r}$$

$$R \geq b: Q_{enc} = -4\pi \rho_0 (b-a) \rightarrow D(R) = \frac{\rho_0 (a-b)}{R^2} \hat{r}$$