

Normal Adult Physiological Values Relevant to Emergency Cardiac & Respiratory Conditions

This document summarizes normal adult physiological values used as reference standards when assessing tachycardia, bradycardia, hypertensive emergencies, atrial fibrillation, and suspected myocardial infarction. Values apply to resting, non-pregnant adults. Reference numbers are indicated in brackets.

Parameter	Normal Value	Reference
Heart Rate	60–100 beats/min	[1,2]
Respiratory Rate	12–20 breaths/min	[2,3]
Blood Pressure	<120/80 mmHg	[1,3]
Core Body Temperature	36.5–37.5 °C	[4]
Oxygen Saturation (SpO ₂)	≥95% (room air)	[1,5]
Cardiac Rhythm	Regular sinus rhythm	[1,2]
PR Interval	120–200 ms	[1]
QRS Duration	<120 ms	[1]
QTc Interval	<440 ms (men), <460 ms (women)	[1,2]
Serum Troponin I/T	Below 99th percentile (lab specific)	[1,2]
Systolic BP (lower limit)	≥90 mmHg	[3]
Mean Arterial Pressure (MAP)	≥65 mmHg	[3]

References

1. American Heart Association (AHA). ACLS Guidelines and ECG Interpretation Standards.
2. Harrison's Principles of Internal Medicine, 21st Edition.
3. Davidson's Principles and Practice of Medicine, 24th Edition.
4. World Health Organization (WHO). Normal Body Temperature Ranges.
5. UpToDate®. Normal Vital Signs in Adults.