



Master's thesis
Astronomy

Your Title Here

Anni Järvenpää

June 19, 2017

Tutor: Associate Professor Peter Johansson
Dr. Till Sawala

Censors: prof. Smith
doc. Smythe

UNIVERSITY OF HELSINKI
DEPARTMENT OF PHYSICS

PL 64 (Gustaf Hållströmin katu 2a)
00014 University of Helsinki

Contents

1	Introduction	1
1.1	TL;DR version of prerequisite information	1
1.2	History of Local Group Research	1
1.3	Aim of This Thesis	2
2	Theoretical Background	3
2.1	Local Group	3
2.1.1	Structure	3
2.1.2	Evolution	3
2.2	Expanding universe	4
2.2.1	Discovery	4
2.2.2	Λ CDM Cosmology	4
2.2.3	Hubble flow	4
3	Mathematical and statistical methods	5
3.1	Statistical Background	5
3.1.1	Hypothesis testing and p-values	6
3.1.2	Distribution functions	7
3.2	Regression Analysis	10
3.3	Error analysis	10
3.4	Comparing two samples drawn from unknown distributions	10

3.4.1	Kolmogorov-Smirnov test	11
3.5	Cluster Analysis	13
4	general simulation thingies	14
4.1	N-body simulations	14
4.1.1	Hierarchical Tree Algorithm	14
4.1.2	Numerical Integrators	14
4.1.3	Halo Finding with Subfind	14
4.2	Description of actual simulations used	14
5	Findings from DMO Halo Catalogue Analysis	15
5.1	Selection of Local Group analogues	15
5.2	Local Anisotropy of the Hubble Flow	15
5.2.1	Hubble Flow Fitting	15
5.3	Statistical Estimate of the Local Group Mass	15
6	Conclusions	16
	Bibliography	17

1. Introduction

1.1 TL;DR version of prerequisite information

1. galaxies form
 - Why?
 - When?
 - How?
 - Where?
2. galaxies form in groups
3. our local group is one of these
4. something about large scale distribution of galaxies

1.2 History of Local Group Research

LG objects visible with naked eye -> realization they are something outside our galaxy -> realization they are something very much like our galaxy

First determining distance was difficult, now mass is more interesting question

1.3 Aim of This Thesis

Whatever the main results end up being, presented in somewhat coherent manner and hopefully sugar-coated enough to sound Important and Exciting.

2. Theoretical Background

Think whether LG or LCDM first

2.1 Local Group

Definition of galaxy group, our local group is one of these.

Mass estimate (Li, Yang masses for the LG and MW)

Maybe something about scale of things in our universe, what are galaxy groups made of, what do you get if you go one distance scale up, what's different in galaxy clusters

2.1.1 Structure

Galaxies that are part of LG, distribution of smaller ones around bigger ones

Current mass estimates (at least timing argument, hubble flow and maybe satellites)

2.1.2 Evolution

How have we ended up in a situation described earlier? What will happen in future?

2.2 Expanding universe

2.2.1 Discovery

Make maths, add cosmological constant, make observations, remove cosmological constant

Enough cosmology here or in other sections to make other parts of thesis to make sense and to suffice as master's thesis = basic textbook cosmology and galaxy formation theory

2.2.2 Λ CDM Cosmology

2.2.3 Hubble flow

What is, where seen, what means, how to measure, hotness/coldness

Plot: observations with fitted hubble flow

3. Mathematical and statistical methods

TODO: tämä ei
sovi tänne
yksinään/
muutoksitta

Precision of the used equipment limits accuracy of all data gathered from physical experiments, simulations or observations. Therefore the results are affected by the measurement process and the results have to be presented as estimates with some error, magnitude of which is affected by both number of data points and accuracy of the measurement equipment. [Bohm and Zech, 2010]

Estimating errors for measured quantities offers a way to test hypotheses and compare different experiments[Bohm and Zech, 2010]. This is done using different statistical methods, a few of which are covered here. Methods used in this work are shortly introduced in the following sections together with basic statistical concepts that are necessary to understand the methods.

3.1 Statistical Background

täällä tarvittavat esitiedot ja önnönnöö, listaa mm. mitä aiot kertoa kunhan tiedät itsekkään

3.1.1 Hypothesis testing and p-values

A common situation in scientific research is that one has to compare a sample to either a model or another sample in order to derive a conclusion from the dataset. In statistics, this is known as hypothesis testing. For example, this can mean testing hypotheses like "these two variables are not correlated" or "this sample is from a population with a mean of 1.0". [J. V. Wall, 2003] Next paragraphs shortly introduce the basic concept of hypothesis testing and methods that can be used to test the hypothesis "these two samples are drawn from the same distribution".

Typically the process of hypothesis testing begins with forming of null hypothesis H_0 that is formatted such that the aim for the next steps is to either reject it or deduce that it cannot be rejected with a chosen significance level. Negation of the null hypothesis is often called research hypothesis or alternative hypothesis and denoted as H_1 . For example, this can lead to H_0 "this dataset is sampled from a normal distribution" and H_1 "this dataset is not sampled from a normal distribution". Choosing the hypothesis in this manner is done because often the research hypothesis is difficult to define otherwise. [Bohm and Zech, 2010; J. V. Wall, 2003]

After setting the hypothesis one must choose an appropriate test statistic. Ideally this is chosen such that the difference between cases H_0 and H_1 is as large as possible. Then one must choose the significance level α which corresponds to the probability of rejecting H_0 in the case where H_0 actually is true. This fixes the critical region i.e. the values of test statistic that lead to the rejection of the H_0 . [Bohm and Zech, 2010; J. V. Wall, 2003]

It is crucial not to look at the test results before choosing α in order to avoid intentional or unintentional fiddling with the data or changing the criterion of acceptance or rejection to give desired results. Only after these steps should the test statistic be calculated. If the test statistic falls within the critical region, H_0 should be rejected and otherwise stated that H_0 cannot be rejected at this significance level.

[Bohm and Zech, 2010; J. V. Wall, 2003]

This kind of probability based decision making is always prone to error. It is easy to see that α corresponds to the chance of H_0 being rejected when it is true. This is known as error of the first kind. However, this is not the only kind of error possible. It might also occur that H_0 is false but it does not get rejected, which is known as error of the second kind. [Bohm and Zech, 2010]

Despite statistical tests having a binary outcome " H_0 rejected" or " H_0 not rejected", a continuous output is often desired. This is what p-values are used for. The name p-value hints towards probability, but despite it's name p-value is not equal to the probability that the null hypothesis is true. These p-values are functions of test statistic and the p-value for a certain value t_{obs} of test statistic gives the probability that under the condition that H_0 is true, the value of a test statistics for a randomly drawn sample is at least as extreme as t_{obs} . Therefore if p-value is smaller than α , H_0 is to be rejected. [Bohm and Zech, 2010]

3.1.2 Distribution functions

[insert lyhyt aloituskappale here]

As the name suggests, PDF is a function the value of which at some point x represents the likelihood that the value of the random variable would equal x . This is often denoted $f(x)$. Naturally for continuous functions the probability of drawing any single value from the distribution is zero, so these values should be interpreted as depicting relative likelihood of different values. For example if $f(a) = 0.3$ and $f(b) = 0.6$ we can say that drawing value a is twice as likely as drawing value b . [Heino et al., 2012]

Another way to use the PDF is to integrate it over semi closed interval from negative infinity to some value a to obtain CDF, often denoted with $F(x)$:

$$F(x) = \int_{-\infty}^x f(x') dx'. \quad (3.1)$$




Figure 3.1: Cumulative distribution function (top panel) for three random samples (PDFs shown on bottom panel) drawn from different distributions, two of which are normal and one is uniform.

This gives the probability of a random value drawn from the distribution having value that is smaller than x . Relation between PDF and CDF is illustrated in figure 3.1, where PDFs and CDFs are shown for three different distributions. It is easy to see the integral relation between PDF and CDF and how wider distributions have wider CDFs.[Heino et al., 2012]

Both PDF and CDF both apply to whole population or the set of all possible outcomes of a measurement. In reality the sample is almost always smaller than this. Therefore one cannot measure the actual CDF. Nevertheless, it is possible to calculate a similar measure of how big a fraction of measurements falls under a given




Figure 3.2: Empirical distribution function for four random samples ($N=35$) drawn from same distributions as in figure 3.1. Note that both blue and cyan data are drawn from the same distribution.

value. This empirical counterpart of the CDF is known as empirical distribution function, often denoted $\hat{F}(x)$, and for a dataset X_1, X_2, \dots, X_n containing n samples it is defined to be

$$\hat{F}(x) = \frac{1}{n} \sum_{i=1}^n I[X_i \leq x] \quad (3.2)$$

where I is the indicator function, value of which is 1 if the condition in brackets is true, otherwise 0. [Feigelson and Babu, 2012]

Due to EDF being a result of random sampling, it may deviate from the underlying CDF considerably as can be seen by comparing CDFs in figure 3.1 and corresponding EDFs in figure 3.2. Latter figure also has EDFs corresponding to two random samples drawn from the distribution of the green curve in the first figure to illustrate the differences that can arise from random sampling. This randomness also makes determining whether two samples are drawn from same distribution difficult.

3.2 Regression Analysis

line fitting and other trivial things

3.3 Error analysis

3.4 Comparing two samples drawn from unknown distributions

TODO: oispa
parempi otsikko.
mieti, onko tämä
muutenkaan hyvä
nyt kun on siirretty
yksi otsikkotasoa
ylöspäin

A common question in multiple fields of science is whether two or more samples are drawn from the same distribution. This can occur for example when comparing effectiveness of two procedures, determining if instrument has changed over time or whether observed data is compatible with simulations. There are multiple two-sample tests that can address this kind of questions, e.g. χ^2 , Kolmogorov-Smirnov, Cramér-von Mises and Anderson-Darling tests. [Bohm and Zech, 2010; Feigelson and Babu, 2012]

In addition to comparing two samples, these tests can be used as one-sample tests to determine whether it is expected that the sample is from a particular distribution. However, some restrictions apply when using the one-sample variants. [Feigelson and Babu, 2012]

Out of these, the χ^2 test uses categorical data, for example "number of data points between values 1.5 and 1.6" or "number of galaxies that are active", and compares numbers of samples in different categories whereas the others compare empirical distribution functions (EDF) of the datasets. Feigelson and Babu [2012]

3.4.1 Kolmogorov-Smirnov test

For astronomers one of the most well-known of these tests is the Kolmogorov-Smirnov test, also known as the KS test. It is computationally inexpensive to calculate, easy to understand and does not require binning of data. It is also non-parametric test i.e. the data does not have to be drawn from a particular distribution. In astrophysical context this is often important because astrophysical models often do not fix a specific statistical distribution for observables and it is common to carry out calculations with logarithms of observables, after which the originally possibly normally distributed residuals will no longer follow normal distribution. Feigelson and Babu [2012]

The test can be used as either one-sample or two-sample test, both of which are very similar. For two-sample variate the test statistic for KS test is calculated based on empirical distribution functions \hat{F}_1 and \hat{F}_2 derived from two samples and the test statistic

$$D = \sup_x |\hat{F}_1(x) - \hat{F}_2(x)| \quad (3.3)$$

uses the maximum vertical distance of the e.d.f's. This test statistic is then used to determine the p-value and thus decide whether the null hypothesis can be rejected. For one-sample variate the procedure is similar, but EDF \hat{F}_2 is substituted with the CDF that corresponds to the null hypothesis. [Bohm and Zech, 2010; Feigelson and Babu, 2012]

As an example, let's consider two pairs of samples from figure 3.2: green and blue (two samples drawn from different normal distributions) and blue and cyan (two samples drawn from same normal distribution). We can formulate the test and null hypotheses for both pairs as H_0 ="the two samples are drawn from the same distribution" and H_1 ="the two samples are not drawn from the same distribution" and choose a significance level of for example $\alpha = 0.05$ or $\alpha = 0.01$.




Figure 3.3: KS test parameter values (magenta vertical lines) shown graphically for three samples from figure 3.2.




Figure 3.4: KS test run on another pair of samples drawn from blue and green distributions in figure 3.1.

The test statistic is then calculated and for these samples we get $D = 0.51$ for the green-blue pair and $D = 0.20$ for the blue-cyan pair. Test statistics are illustrated in figure 3.3 where the test statistics D are shown as vertical magenta lines. These values of D correspond to p-values 9.9×10^{-5} and 0.44 respectively, which means that the null hypothesis "green and blue samples are drawn from the same distribution" is rejected at both 0.05 and 0.01 significance levels but the null hypothesis "blue and cyan samples are drawn from the same distribution" cannot be rejected.

In this case the KS test resulted in result that matches the actual distributions from which the samples are drawn from. Using a different sample might have resulted in different conclusion, for example one shown in figure 3.4 results in $D = 0.17$ that corresponds to p-value of 0.64 i.e. null hypothesis could not have been rejected using α specified earlier.

3.5 Cluster Analysis

DBSCAN

4. general simulation thingies

Data used here from EAGLE which uses modified GADGET-2 which is a tree-code that uses leapfrog, other integrators also briefly introduced?

4.1 N-body simulations

4.1.1 Hierarchical Tree Algorithm

4.1.2 Numerical Integrators

4.1.3 Halo Finding with Subfind

4.2 Description of actual simulations used

Volume, number of particles, compare to other simulations, where better and where maybe worse

Resimulation of interesting regions

Simulation has same parameters as EAGLE 800 Mpc volume used schaye 2015 paper DM-only parts: Volker-Springer Gadget and Gadget 2 papers 1999 and 2005 or something, gravity part is more interesting than SPH Zooms can use multiple meshes, only one is used here gravitational softening

5. Findings from DMO Halo Catalogue Analysis

5.1 Selection of Local Group analogues

criteria, how many found, what are like (some plots maybe? distributions of masses, separations, velocities or correlations between two of those?). This might be part of previous chapter too (relevant to resimulation)?

5.2 Local Anisotropy of the Hubble Flow

Hopefully there's something at least mildly interesting to report when I get to look at the new data

5.2.1 Hubble Flow Fitting

5.3 Statistical Estimate of the Local Group Mass

Analysis similar to Fattahi et al 2016 paper

6. Conclusions

Bibliography

G. Bohm and G. Zech. *Introduction to statistics and data analysis for physicists*. DESY, 2010. ISBN 9783935702416. URL http://www-library.desy.de/preparch/books/vstatmp_engl.pdf.

Eric D. Feigelson and G. Jogesh Babu. *Modern Statistical Methods for Astronomy: With R Applications*. Cambridge University Press, 2012. doi: 10.1017/CBO9781139015653.

R. Heino, K. Ruostenoja, and J. Räisänen. *Havaintojen tilastollinen käsittely*. Department of Physics (University of Helsinki), 2012.

C. R. Jenkins J. V. Wall. *Practical Statistics for Astronomers*. Cambridge Observing Handbooks for Research Astronomers. Cambridge University Press, illustrated edition edition, 2003. ISBN 9780521454162,0521454166.