



Master's thesis  
Astronomy

**Your Title Here**

Anni Järvenpää

May 26, 2017

Tutor: Associate Professor Peter Johansson  
Dr. Till Sawala

Censors: prof. Smith  
doc. Smythe

UNIVERSITY OF HELSINKI  
DEPARTMENT OF PHYSICS

PL 64 (Gustaf Hällströmin katu 2a)  
00014 University of Helsinki





# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	TL;DR version of prerequisite information . . . . .	1
1.2	History of Local Group Research . . . . .	1
1.3	Aim of This Thesis . . . . .	2
<b>2</b>	<b>Theoretical Background</b>	<b>3</b>
2.1	Local Group . . . . .	3
2.1.1	Structure . . . . .	3
2.1.2	Evolution . . . . .	3
2.2	Expanding universe . . . . .	4
2.2.1	Discovery . . . . .	4
2.2.2	$\Lambda$ CDM Cosmology . . . . .	4
2.2.3	Hubble flow . . . . .	4
<b>3</b>	<b>Mathematical and statistical methods</b>	<b>5</b>
3.1	Regression Analysis . . . . .	5
3.2	Error analysis . . . . .	6
3.3	Goodness of fit testing . . . . .	6
3.3.1	Comparing two samples drawn from unknown distributions . .	7
3.4	Cluster Analysis . . . . .	8

<b>4</b>	<b>general simulation thingies</b>	<b>9</b>
4.1	N-body simulations . . . . .	9
4.1.1	Hierarchical Tree Algorithm . . . . .	9
4.1.2	Numerical Integrators . . . . .	9
4.1.3	Halo Finding with Subfind . . . . .	9
4.2	Description of actual simulations used . . . . .	9
<b>5</b>	<b>Findings from DMO Halo Catalogue Analysis</b>	<b>10</b>
5.1	Selection of Local Group analogues . . . . .	10
5.2	Local Anisotropy of the Hubble Flow . . . . .	10
5.2.1	Hubble Flow Fitting . . . . .	10
5.3	Statistical Estimate of the Local Group Mass . . . . .	10
<b>6</b>	<b>Conclusions</b>	<b>11</b>
	<b>Bibliography</b>	<b>12</b>

# 1. Introduction

## 1.1 TL;DR version of prerequisite information

1. galaxies form
  - Why?
  - When?
  - How?
  - Where?
2. galaxies form in groups
3. our local group is one of these
4. something about large scale distribution of galaxies

## 1.2 History of Local Group Research

LG objects visible with naked eye -> realization they are something outside our galaxy -> realization they are something very much like our galaxy

First determining distance was difficult, now mass is more interesting question

## 1.3 Aim of This Thesis

Whatever the main results end up being, presented in somewhat coherent manner and hopefully sugar-coated enough to sound Important and Exciting.

## 2. Theoretical Background

Think whether LG or LCDM first

### 2.1 Local Group

Definition of galaxy group, our local group is one of these.

Mass estimate (Li, Yang masses for the LG and MW)

Maybe something about scale of things in our universe, what are galaxy groups made of, what do you get if you go one distance scale up, what's different in galaxy clusters

#### 2.1.1 Structure

Galaxies that are part of LG, distribution of smaller ones around bigger ones

Current mass estimates (at least timing argument, hubble flow and maybe satellites)

#### 2.1.2 Evolution

How have we ended up in a situation described earlier? What will happen in future?



## 2.2 Expanding universe

### 2.2.1 Discovery

Make maths, add cosmological constant, make observations, remove cosmological constant

Enough cosmology here or in other sections to make other parts of thesis to make sense and to suffice as master's thesis = basic textbook cosmology and galaxy formation theory

### 2.2.2 $\Lambda$ CDM Cosmology

### 2.2.3 Hubble flow

What is, where seen, what means, how to measure, hotness/coldness

Plot: observations with fitted hubble flow

## 3. Mathematical and statistical methods

Precision of the used equipment limits accuracy of all data gathered from physical experiments, simulations or observations. Therefore the results are affected by the measurement process and the results have to be presented as estimates with some error, magnitude of which is affected by both number of data points and accuracy of the measurement equipment. [Bohm and Zech, 2010]

Estimating errors for measured quantities offers a way to test hypotheses and compare different experiments [Bohm and Zech, 2010]. This is done using different statistical methods, a few of which are covered here. Methods used in this work are shortly introduced in the following sections.

### 3.1 Regression Analysis

line fitting and other trivial things

## 3.2 Error analysis

## 3.3 Goodness of fit testing

A common situation in scientific research is that one has to compare a sample to either a model or another sample in order to derive a conclusion from the dataset. In statistics, this is known as hypothesis testing. For example, this can mean testing hypotheses like "these two variables are not correlated" or "this sample is from a population with a mean of 1.0". [J. V. Wall, 2003] Next paragraphs shortly introduce the basic concept of hypothesis testing and methods that can be used to test the hypothesis "these two samples are drawn from the same distribution".

Typically the process of hypothesis testing begins with forming of null hypothesis  $H_0$  that is formatted such that the aim for the next steps is to either reject it or deduce that it cannot be rejected with a chosen significance level. Negation of the null hypothesis is often called research hypothesis or alternative hypothesis and denoted as  $H_1$ . For example, this can lead to  $H_0$  "this dataset is sampled from a normal distribution" and  $H_1$  "this dataset is not sampled from a normal distribution". Choosing the hypotheses in this manner is done because often the research hypothesis is difficult to define otherwise. [Bohm and Zech, 2010; J. V. Wall, 2003]

After setting the hypotheses one must choose an appropriate test statistic. Ideally this is chosen such that the difference between cases  $H_0$  and  $H_1$  is as large as possible. Then one must choose the significance level  $\alpha$  which corresponds to the probability of rejecting  $H_0$  in the case where  $H_0$  actually is true. This fixes the critical region i.e. the values of test statistic that lead to the rejection of the  $H_0$ . [Bohm and Zech, 2010; J. V. Wall, 2003]

It is crucial not to look at the test results before choosing  $\alpha$  in order to avoid intentional or unintentional fiddling with the data or changing the criterion of acceptance or rejection to give desired results. Only after these steps should the test

statistic be calculated. If the test statistic falls within the critical region,  $H_0$  should be rejected and otherwise stated that  $H_0$  cannot be rejected at this significance level. [Bohm and Zech, 2010; J. V. Wall, 2003]

This kind of probability based decision making is always prone to error. It is easy to see that  $\alpha$  corresponds to the chance of  $H_0$  being rejected when it is true. This is known as error of the first kind. However, this is not the only kind of error possible. It might also occur that  $H_0$  is false but it does not get rejected, which is known as error of the second kind. [Bohm and Zech, 2010]

Despite statistical tests having a binary outcome " $H_0$  rejected" or " $H_0$  not rejected", a continuous output is often desired. This is what p-values are used for. The name p-value hints towards probability, but despite it's name p-value is not equal to the probability that the null hypothesis is true. These p-values are functions of test statistic and the p-value for a certain value  $t_{obs}$  of test statistic gives the probability that under the condition that  $H_0$  is true, the value of a test statistics for a randomly drawn sample is at least as extreme as  $t_{obs}$ . Therefore if p-value is smaller than  $\alpha$ ,  $H_0$  is to be rejected. [Bohm and Zech, 2010]

### 3.3.1 Comparing two samples drawn from unknown distributions

A common question in multiple fields of science is whether two or more samples are drawn from the same distribution. This can occur for example when comparing effectiveness of two procedures, determining if instrument has changed over time or whether observed data is compatible with simulations. There are multiple two-sample tests that can address this kind of questions, e.g.  $\chi^2$ , Kolmogorov-Smirnov, Cramér-von Mises and Anderson-Darling tests. [Bohm and Zech, 2010; Feigelson and Babu, 2012]

All of these compare empirical distribution functions of the samples except

the  $\chi^2$ , which uses binned or categorical data and compares numbers of samples in different categories. [kerro tässä tai muualla edf:stä. ehkä kuva edf:stä? esim parille normaalijakaumalle, tophatille ja poissonille tms?]. In addition to comparing two samples, these tests can be used as one-sample tests to determine whether it is expected that the sample is from a particular distribution. Some precautions [Feigelson and Babu, 2012]

For astronomers the most well-known of these tests is the Kolmogorov-Smirnov test, which is also known as the KS test. Test statistic is calculated based on empirical distribution functions  $\hat{F}_1$  and  $\hat{F}_2$  derived from two samples and the test statistic

$$D = \sup_x |\hat{F}_1(x) - \hat{F}_2(x)| \quad (3.1)$$

uses the maximum vertical distance of the e.d.f's as shown in [ehdottomasti kuva tähän]. [Bohm and Zech, 2010; Feigelson and Babu, 2012]

## 3.4 Cluster Analysis

DBSCAN

## 4. general simulation thingies

Data used here from EAGLE which uses modified GADGET-2 which is a tree-code that uses leapfrog, other integrators also briefly introduced?

### 4.1 N-body simulations

#### 4.1.1 Hierarchical Tree Algorithm

#### 4.1.2 Numerical Integrators

#### 4.1.3 Halo Finding with Subfind

### 4.2 Description of actual simulations used

Volume, number of particles, compare to other simulations, where better and where maybe worse

Resimulation of interesting regions

Simulation has same parameters as EAGLE 800 Mpc volume used schaye 2015 paper DM-only parts: Volker-Springer Gadget and Gadget 2 papers 1999 and 2005 or something, gravity part is more interesting than SPH Zooms can use multiple meshes, only one is used here gravitational softening

## **5. Findings from DMO Halo Catalogue Analysis**

### **5.1 Selection of Local Group analogues**

criteria, how many found, what are like (some plots maybe? distributions of masses, separations, velocities or correlations between two of those?). This might be part of previous chapter too (relevant to resimulation)?

### **5.2 Local Anisotropy of the Hubble Flow**

Hopefully there's something at least mildly interesting to report when I get to look at the new data

#### **5.2.1 Hubble Flow Fitting**

### **5.3 Statistical Estimate of the Local Group Mass**

Analysis similar to Fattahi et al 2016 paper

## 6. Conclusions



# Bibliography

G. Bohm and G. Zech. *Introduction to statistics and data analysis for physicists*. DESY, 2010. ISBN 9783935702416. URL [http://www-library.desy.de/preparch/books/vstatmp\\_engl.pdf](http://www-library.desy.de/preparch/books/vstatmp_engl.pdf).

Eric D. Feigelson and G. Jogesh Babu. *Modern Statistical Methods for Astronomy: With R Applications*. Cambridge University Press, 2012. doi: 10.1017/CBO9781139015653.

C. R. Jenkins J. V. Wall. *Practical Statistics for Astronomers*. Cambridge Observing Handbooks for Research Astronomers. Cambridge University Press, illustrated edition edition, 2003. ISBN 9780521454162,0521454166.