



Master's thesis
Astronomy

Your Title Here

Anni Järvenpää

March 30, 2017

Tutor: Associate Professor Peter Johansson
Dr. Till Sawala

Censors: prof. Smith
doc. Smythe

UNIVERSITY OF HELSINKI
DEPARTMENT OF PHYSICS

PL 64 (Gustaf Hällströmin katu 2a)
00014 University of Helsinki

Contents

1	Introduction	1
1.1	TL;DR version of prerequisite information	1
1.2	History of Local Group Research	1
1.3	Aim of This Thesis	2
2	Local Group	3
2.1	Galaxy Groups	3
2.2	Structure	3
2.3	Evolution	3
3	Expanding universe	4
3.1	Discovery	4
3.2	Hubble flow	4
4	Simulation stuff	5
5	Findings from DMO Halo Catalogue Analysis	6
5.1	Selection of Local Group analogues	6
5.2	Local Anisotropy of the Hubble Flow	6
5.3	Statistical Estimate of the Local Group Mass	6
6	Conclusions	15

1. Introduction

1.1 TL;DR version of prerequisite information

1. galaxies form
 - Why?
 - When?
 - How?
 - Where?
2. galaxies form in groups
3. our local group is one of these
4. something about large scale distribution of galaxies

1.2 History of Local Group Research

LG objects visible with naked eye -> realization they are something outside our galaxy -> realization they are something very much like our galaxy

First determining distance was difficult, now mass is more interesting question

1.3 Aim of This Thesis

Whatever the main results end up being, presented in somewhat coherent manner and hopefully sugar-coated enough to sound Important and Exciting.

2. Theoretical Background

Think whether LG or LCDM first

2.1 Local Group

Definition of galaxy group, our local group is one of these.

Mass estimate (Li, Yang masses for the LG and MW)

Maybe something about scale of things in our universe, what are galaxy groups made of, what do you get if you go one distance scale up, what's different in galaxy clusters

2.1.1 Structure

Galaxies that are part of LG, distribution of smaller ones around bigger ones

Current mass estimates (at least timing argument, hubble flow and maybe satellites)

2.1.2 Evolution

How have we ended up in a situation described earlier? What will happen in future?

2.2 Expanding universe

2.2.1 Discovery

Make maths, add cosmological constant, make observations, remove cosmological constant

Enough cosmology here or in other sections to make other parts of thesis to make sense and to suffice as master's thesis. How much is enough for the latter?

2.2.2 Λ CDM Cosmology

2.2.3 Hubble flow

What is, where seen, what means, how to measure, hotness/coldness

Plot: observations with fitted hubble flow

2.3 Mathematical and statistical methods

2.3.1 Regression Analysis

line fitting and other trivial things

2.3.2 Statistical testing

ks-test, maybe others if used, background info about distributions etc in general

2.4 Cluster Analysis

DBSCAN

3. general simulation thingies

Data used here from EAGLE which uses modified GADGET-2 which is a tree-code that uses leapfrog, other integrators also briefly introduced?

3.1 N-body simulations

3.1.1 Hierarchical Tree Algorithm

3.1.2 Numerical Integrators

3.1.3 Halo Finding with Subfind

3.2 Description of actual simulations used

Volume, number of particles, compare to other simulations, where better and where maybe worse

Resimulation of interesting regions

Simulation has same parameters as EAGLE 800 Mpc volume used schaye 2015 paper DM-only parts: Volker-Springer Gadget and Gadget 2 papers 1999 and 2005 or something, gravity part is more interesting than SPH Zooms can use multiple meshes, only one is used here gravitational softening

4. Findings from DMO Halo Catalogue Analysis

4.1 Selection of Local Group analogues

criteria, how many found, what are like (some plots maybe? distributions of masses, separations, velocities or correlations between two of those?). This might be part of previous chapter too (relevant to resimulation)?

4.2 Local Anisotropy of the Hubble Flow

Hopefully there's something at least mildly interesting to report when I get to look at the new data

4.3 Statistical Estimate of the Local Group Mass

Analysis similar to Fattahi et al 2016 paper




Figure 4.1: Radial velocities of haloes as a function of distance. Best fit to Hubble flow shown with solid line. Nearby points ignored when fitting shown in gray.




Figure 4.2: Hubble flow with colours depicting angular distance from line connecting Milky Way and Andromeda counterparts in simulation.




Figure 4.3: Velocity dispersion of Hubble flow.




Figure 4.4: Haloes with distances between 2 and 5 Mpc as seen from Milky Way counterpart in simulation. Colours depict deviations from best linear Hubble flow fit ignoring haloes up to 2 Mpc away, blue end meaning haloes coming closer faster than expected and redder colours moving away.




Figure 4.5: Dark matter haloes with distances from 2 to 5 Mpc grouped to clusters using DB-SCAN clustering algorithm. Parameters used for this plot were $ms=5$ and $eps=2$.




Figure 4.6: Median number of clusters found with constant eps on different minsamples.




Figure 4.7: Mean number of clusters found with constant ms on different eps.




Figure 4.8: Standard deviation of velocities within cluster as a function of distance.

5. Conclusions