

# A Toy Model in Three Dimensions Indicating Inverse-Square-Law Emergence from Quantum Entanglement

The HoloCosmo Project  
HC-003-DOC

April 14, 2025

## Abstract

We present a simplified extension of a one-dimensional toy model to three dimensions, demonstrating that an effective force with an inverse-square radial dependence may, in principle, emerge from the decay of mutual information between spatially separated quantum subsystems. This work serves as a sanity check to complement previous 1D investigations, highlighting key challenges and potential avenues toward a more realistic emergent-gravity framework.

## 1 Introduction

Recent theoretical work has explored the idea that spacetime and gravitational dynamics may emerge from the entanglement properties of an underlying quantum state. In one spatial dimension, a toy model has shown that a power-law decay of mutual information between two subsystems can be associated with an effective force obeying an inverse-square law. This note extends that idea into three dimensions, providing a higher-dimensional generalization and exploring the conditions under which similar inverse-square-law behavior emerges.

## 2 Three-Dimensional Setup

Consider a three-dimensional cubic lattice composed of qubits. Each lattice site is labeled by a coordinate vector

$$\mathbf{x} = (x_1, x_2, x_3),$$

with  $x_i \in \{1, 2, \dots, L\}$  for a lattice of side length  $L$ . We assume either open or periodic boundary conditions.

Define two disjoint subsystems,  $A$  and  $B$ , each consisting of a contiguous block (or cluster) of qubits. Let  $\mathbf{r}_A$  and  $\mathbf{r}_B$  denote the centers of mass of the subsystems, and define the separation

$$r = \|\mathbf{r}_A - \mathbf{r}_B\|.$$

The quantum correlations between  $A$  and  $B$  are quantified by the mutual information,

$$I(A : B) = S(A) + S(B) - S(A \cup B),$$

where  $S(X)$  is the von Neumann entropy of subsystem  $X$ .

### 3 Power-Law Decay of Mutual Information

We postulate that the mutual information decays with the distance  $r$  according to a power law:

$$I(A : B) \sim \frac{c}{r^\alpha},$$

where  $c > 0$  is a constant and  $\alpha > 0$  characterizes the decay. In many three-dimensional systems—especially those at criticality or in gapless phases—power-law decays of correlation functions (and thus entanglement measures) are commonplace.

### 4 Entanglement Potential and Effective Force

We define an *entanglement potential* as

$$V_{\text{ent}}(r) = -kI(A : B),$$

with  $k > 0$  setting the overall coupling strength. Following classical mechanics, we define an effective force as the negative radial derivative of the potential:

$$F(r) = -\frac{dV_{\text{ent}}}{dr} = k\frac{d}{dr}I(A : B).$$

Substituting our power-law ansatz,

$$I(A : B) \sim \frac{c}{r^\alpha},$$

we compute

$$F(r) = k\frac{d}{dr}\left(\frac{c}{r^\alpha}\right) = -k\alpha c\frac{1}{r^{\alpha+1}}.$$

### 5 Recovering the Inverse-Square Law

To mimic the Newtonian gravitational force, which scales as  $1/r^2$ , we require that

$$\frac{1}{r^{\alpha+1}} \propto \frac{1}{r^2}.$$

This condition implies:

$$\alpha + 1 = 2 \implies \alpha = 1.$$

Thus, if the mutual information decays as  $I(A : B) \sim 1/r$ , the effective force becomes

$$F(r) \sim \frac{1}{r^2},$$

which is the familiar inverse-square law observed in Newtonian gravity.

## 6 Discussion and Outlook

This three-dimensional extension represents a straightforward, yet important, step toward more realistic models of emergent gravity:

- **Exponent Tuning:** In a true 3D quantum many-body system, engineering a precise  $1/r$  decay may require fine-tuning or may naturally occur only under special critical conditions.
- **Static versus Dynamic Models:** Our analysis is entirely static. For a complete gravitational theory, one must consider time-dependent entanglement correlations and their impact on emergent spacetime dynamics.
- **Towards Continuum Theories:** Ultimately, it is desirable to derive a continuum description linking entanglement dynamics with an effective field theory that includes gravitational degrees of freedom.

While this model does not provide a complete theory of gravity, it offers a proof-of-principle that quantum entanglement can encode an effective inverse-square interaction—thereby complementing earlier 1D studies and guiding future research in emergent gravity scenarios.

## 7 Conclusion

We have extended a toy model based on the power-law decay of mutual information from one to three dimensions. By positing that  $I(A : B) \sim 1/r$  for well-chosen subsystems in a 3D lattice, the resulting entanglement potential yields an effective force that scales as  $1/r^2$ . Although highly idealized, this extension reinforces the possibility that gravity-like interactions may emerge from the underlying quantum entanglement structure, and it serves as a stepping stone for more comprehensive explorations connecting quantum information theory with the geometry of spacetime.

## Acknowledgments

The author thanks the broader community exploring emergent gravity and quantum informational approaches for inspiring this work.