- 1. Which one of the following generates new genetic combinations leading to variation?
 - Sexual reproduction
 - (2) Nucellar polyembryony
 - (3) Vegetative reproduction
 - (4) Parthenogenesis
- 2. Match Column-I with Column-II and select the correct option using the codes given below:

Column—I

Column—II

- a. Pistils fused together
- (i) Gametogenesis
- b. Formation of
- (ii) Pistillate
- gametes c. Hyphae of higher
- (iii) Syncarpous
- Ascomycetes d. Unisexual female flower
- (iv) Dikaryotic

Codes:

- а b c d (1)(i) (ii) (iv) (iii) (iii) (i) (iv) (ii) (3) (iv) (iii) (i) (ii) (4) (ii) (i) (iv) (iii)
- 3. In majority of angiosperms
 - (1) reduction division occurs the megaspore mother cells
 - (2) a small central cell is present in the embryo sac
 - (3) egg has a filiform apparatus
 - (4) there are numerous antipodal cells
- 4. Pollination in water hyacinth and water lily is brought about by the agency of
 - (1) birds
 - (2) bats
 - √3) water
 - (4) insects or wind
- 5. The ovule of an angiosperm is technically equivalent to
 - (1) megaspore mother cell
 - 42) megaspore
 - (3) megasporangium
 - (4) megasporophyll

- (1) translocation رك) crossing-over

replication on

(3) Vinca rosea

(4) Vicia faba

(2) E. coli

(1) Drosophila melanogaster

- (3) inversion
- (4) duplication
- 8. The equivalent of a structural gene is

6. Taylor conducted the experiments to prove

7. The mechanism that causes a gene to move

from one linkage group to another is called

semiconservative mode of chromosome

- (1) operon
- (2) recon
- (3) muton
- (4) cistron
- 9. A true breeding plant is
 - (1) near homozygous and produces offspring of its own kind
 - (2) always homozygous recessive in its genetic constitution
 - (3) one that is able to breed on its own
- produced due to cross-pollination among unrelated plants
- 10. Which of the following rRNAs acts as structural RNA as well as ribozyme in bacteria?
 - (1) 23 S rRNA
 - (2) 5.8 S rRNA
 - (3) 5 S rRNA
 - (4) 18 S rRNA
- 11. Stirred-tank bioreactors have been designed
 - (1) availability of oxygen throughout the process
 - (2) ensuring anaerobic conditions in the culture vessel
 - (3) purification of product
 - (4) addition of preservatives to the product

JMD/E3

orove some

nove

:d

ring

its

ong

28

red

ict

- 12. A foreign DNA and plasmid cut by the same restriction endonuclease can be joined to form a recombinant plasmid using
 - (1) polymerase III
 - (12) ligase
 - (3) Eco RI
 - (4) Taq polymerase
- **13.** Which of the following is **not** a component of downstream processing?
 - (1) Preservation
 - (2) Expression
 - (3) Separation
 - (4) Purification
- **14.** Which of the following restriction enzymes produces blunt ends?
 - 11) Xho I
 - (2) Hind III
 - (3) Sal I
 - Eco RV

15. Which kind of therapy was given in 1990 to a four-year-old girl with adenosine deaminase (ADA) deficiency?

- (1) Immunotherapy
- (2) Radiation therapy
- (3) Gene therapy
 - (4) Chemotherapy
- 16. How many hot spots of biodiversity in the world have been identified till date by Norman Myers?
 - (1) 34
 - (2) 43

.(4) 25

JMD/E3

- **17.** The primary producers of the deep-sea hydrothermal vent ecosystem are
 - blue-green algae
 - (2) coral reefs
 - (3) green algae
 - (4) chemosynthetic bacteria
- 18. Which of the following is **correct** for r-selected species?
- (1) Small number of progeny with small size
 - (2) Small number of progeny with large size
 - (3) Large number of progeny with small size
 - (4) Large number of progeny with large size
- 19. If '+' sign is assigned to beneficial interaction, '-' sign to detrimental and '0' sign to neutral interaction, then the population interaction represented by '+' '-' refers to
 - (1) commensalism
 - parasitism
 - (3) mutualism
 - (4) amensalism
- 20. Which of the following is correctly matched?
 - (1) Parthenium hysterophorus—Threat to biodiversity
 - Stratification—Population
 - (3) Aerenchyma—Opuntia
 - (4) Age pyramid—Biome
- 21. Red List contains data or information on
 - 11 threatened species
 - (2) marine vertebrates only
 - (3) all economically important plants
 - (4) plants whose products are in international trade

[P.T.O.

3

- **22.** Which one of the following is **wrong** for fungi?
 - (1) They are heterotrophic.
 - (2) They are both unicellular and multicellular.
 - (3) They are eukaryotic.
 - All fungi possess a purely cellulosic cell wall.
- 23. Methanogens belong to
 - (1) Dinoflagellates
 - (2) Slime moulds
 - (3) Eubacteria
 - (A) Archaebacteria
- 24. Select the wrong statement.
 - (1) Diatoms are chief producers in the oceans.
 - (2) Diatoms are microscopic and float passively in water.
 - (3) The walls of diatoms are easily destructible.
 - (4) 'Diatomaceous earth' is formed by the cell walls of diatoms.
- **25.** The label of a herbarium sheet **does not** carry information on
 - (1) local names
 - (2) height of the plant
 - (3) date of collection
 - name of collector
- **26.** Conifers are adapted to tolerate extreme environmental conditions because of
 - thick cuticle
 - (2) presence of vessels
 - (3) broad hardy leaves
 - (4) superficial stomata
- 27. Which one of the following statements is wrong?
 - (1) Agar-agar is obtained from *Gelidium* and *Gracilaria*.
 - Laminaria and Sargassum are used as food.
 - (3) Algae increase the level of dissolved oxygen in the immediate environment.
 - (4) Algin is obtained from red algae, and carrageenan from brown algae.

- 28. The term 'polyadelphous' is related to '
 - (1) corolla
 - (2) calyx
 - (3) gynoecium
 - (4) androecium
- 29. How many plants among Indigofera, Sesbania, Salvia, Allium, Aloe, mustard, groundnut, radish, gram and turnip have stamens with different lengths in their flowers?
 - (1) Five
 - (2) Six
 - (3) Three
 - 14 Four
- 30. Radial symmetry is found in the flowers of
 - (1) Pisum
 - (2) Cassia
 - , 13) Brassica
 - (4) Trifolium
- 31. Free-central placentation is found in
 - JY Brassica
 - (2) Citrus
 - (3) Dianthus
 - JAT Argemone
- 32. Cortex is the region found between
 - (1) endodermis and pith
 - (2) endodermis and vascular bundle
 - (3) epidermis and stele
 - (4) pericycle and endodermis
- **33.** The balloon-shaped structures called tyloses
 - are extensions of xylem parenchyma cells into vessels
 - (2) are linked to the ascent of sap through xylem vessels
 - (3) originate in the lumen of vessels
 - (4) characterize the sapwood

- 34. A non-proteinaceous enzyme is
 - (1) ligase
 - (2) deoxyribonuclease
 - (3) lysozyme
 - (4) ribozyme

zofera,

istard,

have

their

ers of

35. Select the mismatch.

- (1) Protists-Eukaryotes
- (2) Methanogens-Prokaryotes
- (3) Gas vacuoles-Green bacteria
- Large central vacuoles—Animal cells
- 36. Select the wrong statement.
 - Cyanobacteria lack flagellated cells.
 - (2) Mycoplasma is a wall-less microorganism.
 - (3) Bacterial cell wall is made up of peptidoglycan.
 - (4) Pili and fimbriae are mainly involved in motility of bacterial cells.
- **37.** A cell organelle containing hydrolytic enzymes is
 - ribosome
 - 12) mesosome
 - (3) lysosome
 - (4) microsome
- 38. During cell growth, DNA synthesis takes place in
 - (1) G₂ phase
 - (2) M phase
 - (3) S phase
 - (4) G₁ phase
- 39. Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins?
 - (1) Pyruvic acid
 - (2) Acetyl CoA
 - (3) Glucose-6-phosphate
 - (4) Fructose 1,6-bisphosphate

40. A few drops of sap were collected by cutting across a plant stem by a suitable method. The sap was tested chemically. Which one of the following test results indicates that it is phloem sap?

(1) Low refractive index

- (2) Absence of sugar
- (3) Acidic

(4) Alkaline

- 41. You are given a tissue with its potential for differentiation in an artificial culture. Which of the following pairs of hormones would you add to the medium to secure shoots as well as roots?
 - (1) Auxin and abscisic acid
 - (2) Gibberellin and abscisic acid
 - (3) IAA and gibberellin
 - Auxin and cytokinin
- 42. Phytochrome is a
 - (1) lipoprotein
 - (2) chromoprotein
 - (3) flavoprotein
 - (4) glycoprotein
- 43. Which is essential for the growth of root tip?

Ul Ca

(2) Mn

(3) Zn

- (4) Fe
- 44. The process which makes major difference between C₃ and C₄ plants is
 - (1) photorespiration
 - (2) respiration
 - (3) glycolysis
 - (4) Calvin cycle
- **45.** Which one of the following statements is **not** correct?
 - In potato, banana and ginger, the plantlets arise from the internodes present in the modified stem.
 - (2) Water hyacinth, growing in the standing water, drains oxygen from water that leads to the death of fishes.
 - (3) Offspring produced by the asexual reproduction are called clone.
 - (4) Microscopic, motile asexual reproductive structures are called zoospores.

JMD/E3

5

- **46.** The part of nephron involved in active reabsorption of sodium is
 - (1) Bowman's capsule
 - (2) descending limb of Henle's loop
 - (3) distal convoluted tubule
 - (4) proximal convoluted tubule
- **47.** Which of the following is hormone-releasing IUD?
 - (1) Lippes loop
 - (2) Cu7
 - 48) LNG-20
 - (4) Multiload 375
- **48.** Which of the following is **incorrect** regarding vasectomy?
 - (1) Vasa deferentia is cut and tied
 - (2) Irreversible sterility
 - (3) No sperm occurs in seminal fluid
 - A No sperm occurs in epididymis
- **49.** Embryo with more than 16 blastomeres formed due to *in vitro* fertilization is transferred into
 - (1) fimbriae
 - (2) cervix
 - (3) uterus
 - (4) fallopian tube
- **50.** Which of the following depicts the **correct** pathway of transport of sperms?
 - (1) Rete testis \rightarrow Vas deferens \rightarrow Efferent ductules \rightarrow Epididymis
 - (2) Efferent ductules → Rete testis → Vas deferens → Epididymis
 - Rete testis → Efferent ductules → Epididymis → Vas deferens
 - (4) Rete testis → Epididymis → Efferent ductules → Vas deferens
- 51. Match Column—I with Column—II and select the correct option using the codes given below:

Column—I

Column—II


- a. Mons pubis
- (i) Embryo formation
- b. Antrum
- (ii) Sperm
- c. Trophectoderm
- (iii) Female external genitalia
- d. Nebenkern
- (iv) Graafian follicle

Codes:

	а	b	С	d
(1)	(iii)	(i)	(iv)	(ii)
(2)	(i)	(iv)	(iii)	(ii)
(3)	(iii)	(iv)	(ii)	(i)
142	(iii)	(iv)	(i)	(ii)

JMD/E3

- **52.** Several hormones like hCG, hPL, estrogen, progesterone are produced by
 - (1) fallopian tube
 - (2) pituitary
 - (3) ovary
 - (4) placenta
- **53.** If a colour-blind man marries a woman who is homozygous for normal colour vision, the probability of their son being colour-blind is
 - (1) 0.75
- (2) 1
- (3) 0
- (4)-0.5
- 54. Genetic drift operates in
 - (1) non-reproductive population
 - (2) slow reproductive population
 - (3) small isolated population
 - (4) large isolated population
- **55.** In Hardy-Weinberg equation, the frequency of heterozygous individual is represented by
 - (1) pq
- (2) q^2
- (3) p^2
- JAY 2pq
- **56.** The chronological order of human evolution from early to the recent is
 - (1) Ramapithecus → Homo habilis Australopithecus → Homo erectus
 - (2) Australopithecus → Homo habilis Ramapithecus → Homo erectus
 - (3) Australopithecus → Ramapithecus Homo habilis → Homo erectus
 - (A) Ramapithecus → Australopithecus → Homo habilis → Homo erectus
- 57. Which of the following is the correct sequence of events in the origin of life?
 - I. Formation of protobionts
 - II. Synthesis of organic monomers
 - III. Synthesis of organic polymers
 - IV. Formation of DNA-based genetic systems
 - (1) II, III, I, IV
 - (2) II, III, IV, I
 - (3) I, II, III, IV
 - (4) I, III, II, IV



6

estrogen,.

man who ision, the

·blind is

- 58. A molecule that can act as a genetic material must fulfill the traits given below, except
 - it should be unstable structurally and chemically
 - (2) it should provide the scope for slow changes that are required for evolution
 - (3) it should be able to express itself in the form of 'Mendelian characters'
 - (4) it should be able to generate its replica
- **59.** DNA-dependent RNA polymerase catalyzes transcription on one strand of the DNA which is called the
 - (1) alpha strand
 - (2) antistrand
 - (13) template strand
 - (4) coding strand
- 60. Interspecific hybridization is the mating of

requency nted by

volution

bilis

:bilis

?cus

?cus

life?

correct

systems

- superior males and females of different breeds
 - (2) more closely related individuals within same breed for 4-6 generations
 - animals within same breed without having common ancestors
 - (4) two different related species

61. Which of the following is correct regarding

AIDS causative agent HIV?

- (1) HIV is unenveloped retrovirus.
- (2) HIV does not escape but attacks the acquired immune response.
- HIV is enveloped virus containing one molecule of single-stranded RNA and one molecule of reverse transcriptase.
 - (4) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptase.
- 62. Among the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids?
 - Mrigala ﴿لل
 - (2) Mackerel
 - (3) Mystus
 - Mangur Mangur

63. Match Column-II with Column-II and select the correct option using the codes given below:

5,,,	C11 DC.				
Column—I					Column—II
a.	Citric acid			(i)	Trichoderma
b.	Cyclosporin A			(ii)	Clostridium
c.	Statins			(iii)	Aspergillus
d.	Butyric acid (iv)			Monascus	
Co	des :				
	а	b	С	d	
(1)	(i)	(iv)	(ii)	(iii)	
(2)	(iii)	(iv)	(i)	(ii)	
(3)	(iii)	(i)	(ii)	(iv)	

(iii) 64. Biochemical Oxygen Demand (BOD) may not be a good index for pollution for water bodies receiving effluents from

(iv)

(ii)

(1) petroleum industry

(i)

- (2) sugar industry
- (3) domestic sewage
- (4) dairy industry
- 65. The principle of competitive exclusion was stated by
 - (1) MacArthur
 - (2) Verhulst and Pearl
 - 431 C. Darwin
 - (4) G. F. Gause
- 66. Which of the following National Parks is home to the famous musk deer or hangul?
 - (1) Eaglenest Wildlife Sanctuary, Arunachal Pradesh
 - (2) Dachigam National Park, Jammu & Kashmir
 - (3) Keibul Lamjao National Park, Manipur
 - (4) Bandhavgarh National Park, Madhya Pradesh
- 67. A lake which is rich in organic waste may result in
 - (1) increased population of fish due to lots of nutrients
 - (2) mortality of fish due to lack of oxygen
 - population (3) increased of aquatic organisms due to minerals
 - drying of the lake due to algal bloom
- 68. The highest DDT concentration in aquatic food chain shall occur in

(1) crab

- De eel
- (3) phytoplankton
- (4) seagull

JMD/E3

- 69. Which of the following sets of diseases is caused by bacteria?
 - (1) Tetanus and mumps
 - (2) Herpes and influenza
 - (3) Cholera and tetanus
 - (4) Typhoid and smallpox
- 70. Match Column—I with Column—II for housefly classification and select the correct option using the codes given below:

Column—I Column-II Diptera a. Family Arthropoda b. Order (ii) (iii) Muscidae c. Class (iv) Insecta d. Phylum Codes: (1) (iv) (iii) (ii) (i) (iii) (2) (iv) (ii) (i)

(iv)

(iv)

(i) (4) (iii) (ii) 71. Choose the correct statement.

(i)

137 (iii)

- (1) All reptiles have a three-chambered
 - (2) All Pisces have gills covered by an operculum.


(ii)

- (3) All mammals are viviparous.
- (4) All cyclostomes do not possess jaws and paired fins.
- 72. Study the four statements (A-D) given below and select the two correct ones out of them:
 - A. Definition of biological species was given by Ernst Mayr.
 - B. Photoperiod does not affect reproduction in plants.
 - C. Binomial nomenclature system was given by R. H. Whittaker.
 - D. In unicellular organisms, reproduction is synonymous with growth.

The two correct statements are

- (1) A and D
- (2) A and B
- (3) B and C
- (4) C and D
- 73. In male cockroaches, sperms are stored in which part of the reproductive system?
 - (1) Testes
 - (2) Vas deferens
 - (3) Seminal vesicles
 - (4) Mushroom glands
- 74. Smooth muscles are
 - (H) involuntary, cylindrical, striated
 - (2) voluntary, spindle-shaped, uninucleate
 - (3) involuntary, fusiform, non-striated
 - (4) voluntary, multinucleate, cylindrical

- 75. Oxidative phosphorylation is
 - (1) addition of phosphate group to ATP
 - (2) formation of ATP by energy released from electrons removed during substrate oxidation
 - transfer of (3) formation of ATP by phosphate group from a substrate to ADP
 - (A) oxidation of phosphate group in ATP
- 76. Which of the following is the least likely to be involved in stabilizing the three-dimensional folding of most proteins?
 - (1) Hydrophobic interaction
 - (2) Ester bonds
 - (3) Hydrogen bonds
 - (4) Electrostatic interaction
- 77. Which of the following describes the given graph correctly?



- (1) Endothermic reaction with energy A in absence of enzyme and B in presence of
- Exothermic reaction with energy A in absence of enzyme and B in presence of enzyme
 - (3) Endothermic reaction with energy A in presence of enzyme and B in absence of enzyme
 - (4) Exothermic reaction with energy A in presence of enzyme and B in absence of
- 78. When cell has stalled DNA replication fork, which checkpoint should be predominantly activated?
 - (1) M
 - (2) Both G₂/M and M
 - (3) G_1/S
 - (4) G_2/M

) ATP ased from substrate

nsfer of substrate

in ATP kely to be aensional

he given

rgy A in

gy A in sence of

sence of

sence of

rgy A in

gy A in sence of

ion fork. ninantly 79. Match the stages of meiosis in Column—I to their characteristic features in Column-II and select the correct option using the codes given below:

Column—I

Column—II

- a. Pachytene
- (i) Pairing of homologous chromosomes
- Metaphase I
- Terminalization of chiasmata
- Diakinesis
- (iii) Crossing-over takes place
- d. Zygotene
- (iv) Chromosomes align at equatorial plate

Codes:

	а	b	c	d
(1)	(ii)	(iv)	(iii)	(i)
(2)	(iv)	(iii)	(ii)	(i)
48ل	(iii)	(iv)	(ii) -	(i)
(4)	(i) `	(iv)	(ii)	(iii)


- 80. Which hormones do stimulate the pancreatic and production of juice bicarbonate?
 - (1) Cholecystokinin and secretin
 - (2) Insulin and glucagon
 - (3) Angiotensin and epinephrine
 - (4) Gastrin and insulin
- 81. The partial pressure of oxygen in the alveoli of the lungs is
 - (1) less than that in the blood
 - (2) less than that of carbon dioxide
 - (3) equal to that in the blood
 - more than that in the blood
- 82. Choose the correct statement.
 - Photoreceptors in the human eye are depolarized during darkness and become hyperpolarized in response to the light stimulus.
 - do not produce graded (2) Receptors potentials.
 - (3) Nociceptors respond to changes in pressure.
 - (4) Meissner's corpuscles thermoreceptors.
- 83. Graves' disease is caused due to
 - Hyposecretion of adrenal gland
 - hypersecretion of adrenal gland
 - (3) hyposecretion of thyroid gland
 - (4) hypersecretion of thyroid gland

- 84. Name the ion responsible for unmasking of active sites for myosin for cross-bridge activity during muscle contraction.
 - (1) Sodium
- (2) Potassium
- (3) Calcium
- (4) Magnesium
- 85. Name the blood cells, whose reduction in number can cause clotting disorder, leading to excessive loss of blood from the body.
 - (1) Neutrophils
 - (2) Thrombocytes
 - (3) Erythrocytes
 - (4) Leucocytes
- 86. Name a peptide hormone which acts mainly on hepatocytes, adipocytes and enhances cellular glucose uptake and utilization.
 - (1) Secretin
- (2) Gastrin
- (3) Insulin
- (A) Glucagon
- 87. Osteoporosis, an age-related disease of skeletal system, may occur due to
 - (1) decreased level of estrogen
 - (2) accumulation of uric acid leading to inflammation of joints
 - immune disorder affecting muscular junction leading to fatigue
 - (4) high concentration of Ca++ and Na+
- 88. Serum differs from blood in
 - (1) lacking clotting factors
 - (2) lacking antibodies
 - (3) lacking globulins
 - (4) Tacking albumins
- 89. Lungs do not collapse between breaths and some air always remains in the lungs which can never be expelled because
 - (1) there is a positive intrapleural pressure
 - (2) pressure in the lungs is higher than the atmospheric pressure
 - (3) there is a negative pressure in the lungs
 - there is a negative intrapleural pressure pulling at the lung walls
- 90. The posterior pituitary gland is not a 'true' endocrine gland because
 - it is under the regulation of hypothalamus
 - (2) it secretes enzymes
 - (3) it is provided with a duct
 - (4) it only stores and releases hormones


JMD/E3

9

91. A parallel-plate capacitor of area A, plate separation d and capacitance C is filled with four dielectric materials having dielectric constants k_1 , k_2 , k_3 and k_4 as shown in the figure below. If a single dielectric material is to be used to have the same capacitance C in this capacitor, then its dielectric constant k is given by



- (1) $\frac{2}{k} = \frac{3}{k_1 + k_2 + k_3} + \frac{1}{k_4}$
- (2) $\frac{1}{k} = \frac{1}{k_1} + \frac{1}{k_2} + \frac{1}{k_3} + \frac{3}{2k_4}$
- (3) $k = k_1 + k_2 + k_3 + 3k_4$
- (4) $k = \frac{2}{3}(k_1 + k_2 + k_3) + 2k_4$
- **92.** The potential difference $(V_A V_B)$ between the points A and B in the given figure is




- (1) + 6 V
- (2) +9 V
- (3) -3 V
- (4) +3 V
- 93. A filament bulb (500 W, 100 V) is to be used in a 230 V main supply. When a resistance R is connected in series, it works perfectly and the bulb consumes 500 W. The value of R is
 - (1) 26 Ω
- (2) 13 Ω
- (3) 230 Ω
- (4) 46 Ω
- 94. A long wire carrying a steady current is bent into a circular loop of one turn. The magnetic field at the centre of the loop is B. It is then bent into a circular coil of n turns. The magnetic field at the centre of this coil of n turns will be
 - 2nB
- \cdot (2) $2n^2B$
- (3) nB
- (4) n^2B
- JMD/E3

10

- 95. A bar magnet is hung by a thin cotton thread in a uniform horizontal magnetic field and is in equilibrium state. The energy required to rotate it by 60° is W. Now the torque required to keep the magnet in this new position is
 - (1) $\frac{\sqrt{3}W}{2}$
- (2) $\frac{2W}{\sqrt{3}}$
- $(3) \ \frac{W}{\sqrt{3}}$
- (4) √3W
- **96.** An electron is moving in a circular path under the influence of a transverse magnetic field of 3.57×10^{-2} T. If the value of e/m is 1.76×10^{11} C/kg, the frequency of revolution of the electron is
 - (1) 62·8 MHz
- (2) 6·28 MHz
- (3) 1 GHz
- (4) 100 MHz
- **97.** Which of the following combinations should be selected for better tuning of an *L-C-R* circuit used for communication?
 - (1) $R = 15 \Omega$, L = 3.5 H, $C = 30 \mu\text{F}$
 - (2) $R = 25 \Omega$, L = 1.5 H, $C = 45 \mu F$
 - (3) $R = 20 \Omega$, L = 1.5 H, $C = 35 \mu F$
 - (4) $R = 25 \Omega$, L = 2.5 H, $C = 45 \mu F$
- **98.** A uniform magnetic field is restricted within a region of radius r. The magnetic field changes with time at a rate $\frac{d\vec{B}}{dt}$. Loop 1 of radius R > r encloses the region r and loop 2

radius R > r encloses the region r and loop 2 of radius R is outside the region of magnetic field as shown in the figure below. Then the e.m.f. generated is





- (1) $-\frac{d\vec{B}}{dt}\pi R^2$ in loop 1 and zero in loop 2
- (2) $-\frac{d\vec{B}}{dt}\pi r^2$ in loop 1 and zero in loop 2
- (3) zero in loop 1 and zero in loop 2
- (4) $-\frac{d\vec{B}}{dt}\pi r^2$ in loop 1 and
 - $-\frac{d\vec{B}}{dt}\pi r^2$ in loop 2

313 65

i thread eld and required torque is new

circular nsverse e value iency of

should 1 *L-C-R*

within ic field

op 1 of

loop 2 agnetic nen the

э**ор** 2

юр 2

3

loop 2

99. The potential differences across the resistance, capacitance and inductance are 80 V, 40 V and 100 V respectively in an *L-C-R* circuit. The power factor of this circuit is

- (1) 0.8
- (2) 1.0
- (3) 0.4
- (4) 0.5

100. A $100~\Omega$ resistance and a capacitor of $100~\Omega$ reactance are connected in series across a 220 V source. When the capacitor is 50% charged, the peak value of the displacement current is

- (1) 4·4 A
- (2) $11\sqrt{2}$ A
- (3) 2·2 A
- (4) 11 A

101. Two identical glass ($\mu_g = 3/2$) equiconvex lenses of focal length f each are kept in contact. The space between the two lenses is filled with water ($\mu_w = 4/3$). The focal length of the combination is

- (1) 4f/3
- (2) 3f/4
- (3) f/3
- (4) f

102. An air bubble in a glass slab with refractive index 1.5 (near normal incidence) is 5 cm deep when viewed from one surface and 3 cm deep when viewed from the opposite face. The thickness (in cm) of the slab is

- (1) 12
- (2) 16
- (3) 8
- (4) 10

103. The interference pattern is obtained with two coherent light sources of intensity ratio n. In the interference pattern, the ratio

$$\frac{I_{\max} - I_{\min}}{I_{\max} + I_{\min}}$$

will be

- (1) $\frac{\sqrt{n}}{(n+1)^2}$
- $(2) \ \frac{2\sqrt{n}}{\left(n+1\right)^2}$
- $(3) \ \frac{\sqrt{n}}{n+1}$
- $(4) \ \frac{2\sqrt{n}}{n+1}$

the the lare they lie between 50 cm and 400 cm from his eyes. In order to increase the maximum distance of distinct vision to infinity, the type and power of the correcting lens, the person has to use, will be

- (1) concave, -0.2 diopter
- (2) convex, +0·15 diopter
- (3) convex, +2.25 diopter
- (4) concave, -0.25 diopter

105. A linear aperture whose width is 0.02 cm is placed immediately in front of a lens of focal length 60 cm. The aperture is illuminated normally by a parallel beam of wavelength 5×10^{-5} cm. The distance of the first dark band of the diffraction pattern from the centre of the screen is

(1) 0·20 cm

27.000

- (2) 0·15 cm
- (3) 0·10 cm
- (4) 0·25 cm

106. Electrons of mass m with de-Broglie wavelength λ fall on the target in an X-ray tube. The cutoff wavelength (λ_0) of the emitted X-ray is

(1)
$$\lambda_0 = \frac{2m^2c^2\lambda^3}{h^2}$$

(2) $\lambda_0 = \lambda$

(3)
$$\lambda_0 = \frac{2mc\lambda^2}{h}$$

$$(4) \quad \lambda_0 = \frac{2h}{mc}$$


107. Photons with energy 5 eV are incident on a cathode C in a photoelectric cell. The maximum energy of emitted photoelectrons is 2 eV. When photons of energy 6 eV are incident on C, no photoelectrons will reach the anode A, if the stopping potential of A relative to C is

- (1) -1 V
- (2) -3 V
- (3) +3 V
- (4) +4 V


11

JMD/E3

- 108. If an electron in a hydrogen atom jumps | 113. Planck's constant (h), speed of light in from the 3rd orbit to the 2nd orbit, it emits a photon of wavelength λ . When it jumps from the 4th orbit to the 3rd orbit, the corresponding wavelength of the photon
- $(2) \ \frac{20}{13} \lambda$
- (3) $\frac{16}{25}\lambda$
- $(4) \frac{9}{16}\lambda$
- 109. The half-life of a radioactive substance is 30 minutes. The time (in minutes) taken between 40% decay and 85% decay of the same radioactive substance is
 - (1) 45
- (2) 60
- (3) 15
- (4) 30
- 110. For CE transistor amplifier, the audio signal voltage across the collector resistance of $\bar{2}\,k\Omega$ is 4 V. If the current amplification factor of the transistor is 100 and the base resistance is $1\,k\Omega$, then the input signal voltage is
 - (1) 30 mV
- (2) 15 mV
- (3) 10 mV
- (4) 20 mV
- 111. The given circuit has two ideal diodes connected as shown in the figure below. The current flowing through the resistance R_1



- (1) 1·43 A
- (2) 3·13 A
- (3) 2·5 A
- (4) 10·0 A
- 112. What is the output Y in the following circuit, when all the three inputs A, B, C are first 0 and then 1?



- (3) 0, 1
- (4) 0, 0

- vacuum (c) and Newton's gravitational constant (G) are three fundamental constants. Which of the following combinations of these has the dimension of length?

- 114. Two cars P and Q start from a point at the same time in a straight line and their positions are represented by $x_P(t) = at + bt^2$ and $x_Q(t) = ft - t^2$. At what time do the cars and $x_Q(t) = ft$.


 have the same velocity?

 (1) $\frac{a+f}{2(1+b)}$ (2) $\frac{f-a}{2(1+b)}$ (3) $\frac{a-f}{1+b}$ (4) $\frac{a+f}{2(b-1)}$

- 115. In the given figure, $a = 15 \text{ m/s}^2$ represents the total acceleration of a particle moving in the clockwise direction in a circle of radius R = 2.5 m at a given instant of time. The speed of the particle is



- (1) 5·7 m/s
- (2) 6·2 m/s
- (3) 4.5 m/s
- (4) 5·0 m/s
- 116. A rigid ball of mass m strikes a rigid wall at 60° and gets reflected without loss of speed as shown in the figure below. The value of impulse imparted by the wall on the ball will be



- (4) 2mV

ght in ational mental llowing sion of


at the their $at + bt^2$ ie cars


esents ving in radius e. The


wall at speed ilue of ie ball


- 117. A bullet of mass 10 g moving horizontally | 121. A solid sphere of mass m and radius R is with a velocity of 400 m s⁻¹ strikes a wooden block of mass 2 kg which is suspended by a light inextensible string of length 5 m. As a result, the centre of gravity of the block is found to rise a vertical distance of 10 cm. The speed of the bullet after it emerges out horizontally from the block will be
 - (1) 120 m s^{-1}
 - (2) 160 m s^{-1}
 - (3) 100 m s^{-1}
 - (4) 80 m s^{-1}
- 118. Two identical balls A and B having velocities of 0.5 m/s and -0.3 m/s respectively collide elastically in one dimension. The velocities of B and A after the collision respectively will be
 - (1) -0.3 m/s and 0.5 m/s
 - (2) 0.3 m/s and 0.5 m/s
 - (3) -0.5 m/s and 0.3 m/s
 - (4) 0.5 m/s and -0.3 m/s
- 119. A particle moves from a point $(-2\hat{i} + 5\hat{j})$ to $(4\hat{j}+3\hat{k})$ when a force of $(4\hat{i}+3\hat{j})$ N is applied. How much work has been done by the force?
 - (1) 5 J
 - (2) 2 J
 - (3) 8 J
 - (4) 11 J
- 120. Two rotating bodies A and B of masses mand 2m with moments of inertia I_A and $I_B(I_B > I_A)$ have equal kinetic energy of rotation. If L_A and L_B be their angular momenta respectively, then
 - (1) $L_B > L_A$
 - (2) $L_A > L_B$
 - $(3) L_A = \frac{L_B}{2}$
 - $(4) L_A = 2L_B$

- rotating about its diameter. A solid cylinder of the same mass and same radius is also rotating about its geometrical axis with an angular speed twice that of the sphere. The ratio of their kinetic energies of rotation $(E_{\text{sphere}} / E_{\text{cylinder}})$ will be
 - (1) 1:4
 - (2) 3:1
 - (3) 2:3
 - (4) 1:5
- **122.** A light rod of length l has two masses m_1 and m_2 attached to its two ends. The moment of inertia of the system about an axis perpendicular to the rod and passing through the centre of mass is
 - $(1) (m_1 + m_2)l^2$
- (3) $\frac{m_1 m_2}{m_1 + m_2} l^2$
- 123. Starting from the centre of the earth having radius R, the variation of g (acceleration due to gravity) is shown by









- 124. A satellite of mass m is orbiting the earth (of radius R) at a height h from its surface. The total energy of the satellite in terms of g_0 , the value of acceleration due to gravity at the earth's surface, is
 - $2mg_0R^2$
 - $_{2mg_0R^2}$
 - mg_0R^2 2(R+h)
 - mg_0R^2

- from $(4 \text{ cm} \times 2 \text{ cm})$ to $(5 \text{ cm} \times 4 \text{ cm})$. If the work done is 3×10^{-4} J, the value of the surface tension of the liquid is
 - (1) 0·2 N m⁻¹
- (2) 8·0 N m⁻¹
- (3) 0.250 N m^{-1} (4) 0.125 N m^{-1}
- 126. Three liquids of densities ρ_1 , ρ_2 and ρ_3 (with $\rho_1 > \rho_2 > \rho_3$), having the same value of surface tension T, rise to the same height in three identical capillaries. The angles of contact θ_1 , θ_2 and θ_3 obey
 - $(1) \ \frac{\pi}{2} < \theta_1 < \theta_2 < \theta_3 < \pi$
 - (2) $\pi > \theta_1 > \theta_2 > \theta_3 > \frac{\pi}{2}$
 - (3) $\frac{\pi}{2} > \theta_1 > \theta_2 > \theta_3 \ge 0$
 - (4) $0 \le \theta_1 < \theta_2 < \theta_3 < \frac{\pi}{2}$
 - 127. Two identical bodies are made of a material for which the heat capacity increases with temperature. One of these is at 100 °C, while the other one is at 0 °C. If the two bodies are brought into contact, then, assuming no heat loss, the final common temperature is
 - (1) less than 50 °C but greater than 0 °C
 - (2) 0 °C
 - (3) 50 °C
 - (4) more than 50 °C
 - 128. A body cools from a temperature 3T to 2Tin 10 minutes. The room temperature is T. Assume that Newton's law of cooling is applicable. The temperature of the body at the end of next 10 minutes will be
- (3) $\frac{7}{4}T$
- 129. One mole of an ideal monatomic undergoes a process described by the equation PV^3 = constant. The heat capacity of the gas during this process is
- (3) $\frac{3}{2}R$
- (4) $\frac{5}{2}R$

- 125. A rectangular film of liquid is extended 130. The temperature inside a refrigerator is t_2 °C amount of heat delivered to the room for each joule of electrical energy consumed ideally will be
 - $t_2 + 273$
- (2) $\frac{t_1 + t_2}{t_1 + 273}$ (4) $\frac{t_1 + 273}{t_1 t_2}$

- 131. A given sample of an ideal gas occupies a volume V at a pressure P and absolute temperature T. The mass of each molecule of the gas is m. Which of the following gives the density of the gas?
 - (1) P/(kTV)
- (2) mkT
- (3) P/(kT)
- (4) Pm/(kT)
- 132. A body of mass m is attached to the lower end of a spring whose upper end is fixed. The spring has negligible mass When the mass m is slightly pulled down and released, it oscillates with a time period of $3 ext{ s}$. When the mass m is increase by 1 kg, the time period of oscillation becomes 5 s. The value of m in kg is
- (3)
- 133. The second overtone of an open orga pipe has the same frequency as the fir overtone of a closed pipe L metre long. T length of the open pipe will be
 - (1) $\frac{L}{\hat{c}}$
- (2) 4L
- (3) L
- (4) 2L
- 134. Three sound waves of equal amplitude have frequencies (n-1), n, (n+1). Then superimpose to give beats. The number beats produced per second will be
 - (1) 3
- (2) 2
- (3) 1
- (4) 4
- 135. An electric dipole is placed at an angle of with an electric field intensity 2×10^5 N/0 experiences a torque equal to 4 N m. charge on the dipole, if the dipole lengt 2 cm, is
 - (1) 5 mC
 - (2) 7 µC
 - (3) 8 mC
 - (4) 2 mC

ator is t2°C t₁°C. The 2 room for consumed

s occupies id absolute h molecule owing gives

ed to the upper end gible mass. ulled down time period increased oscillations kg is

pen organ as the first e long. The

amplitudes 1+1). They number of be

angle of 30° 10⁵ N/C. It 4 Nm. The ple length is 136. Hot concentrated sulphuric acid is a 142. Jahn-Teller effect is not observed in high moderately strong oxidizing agent. Which of the following reactions does not show oxidizing behaviour?

(1)
$$C + 2H_2SO_4 \rightarrow CO_2 + 2SO_2 + 2H_2O$$

$$J2$$
 $CaF_2 + H_2SO_4 \rightarrow CaSO_4 + 2HF$

(3)
$$Cu + 2H_2SO_4 \rightarrow CuSO_4 + SO_2 + 2H_2O$$

(4)
$$3S + 2H_2SO_4 \rightarrow 3SO_2 + 2H_2O$$

137. Which of the following pairs of d-orbitals will have electron density along the axes?

$$d_{z^2}, d_{x^2-y^2}$$

- (2) d_{xy} , $d_{x^2-y^2}$
- (3) d_{z^2} , d_{xz}
- (4) d_{xz} , d_{yz}
- 138. The correct geometry and hybridization for XeF₄ are
 - (1) planar triangle, sp^3d^3
 - (2) square planar, sp^3d^2
 - (3) octahedral, sp^3d^2

trigonal bipyramidal, sp³d

- 139. Among the following, which one is a wrong statement?
 - (1) SeF₄ and CH₄ have same shape.
 - (2) I₃⁺ has bent geometry.

PH₅ and BiCl₅ do not exist.

- (4) $p\pi d\pi$ bonds are present in SO₂.
- 140. The correct increasing order of trans-effect of the following species is

(1)
$$Br^- > CN^- > NH_3 > C_6H_5^-$$

(2)
$$CN^- > Br^- > C_6H_5^- > NH_3$$

(3)
$$NH_3 > CN^- > Br^- > C_6H_5^-$$

$$CN^- > C_6H_5^- > Br^- > NH_3$$

- 141. Which one of the following statements related to lanthanons is incorrect?
 - (1) All the lanthanons are much more reactive than aluminium.
 - (2) Ce (+4) solutions are widely used as oxidizing agent in volumetric analysis.
 - (3) Europium shows +2 oxidation state.
 - (4) The basicity decreases as the ionic radius decreases from Pr to Lu.

spin complexes of

(1)
$$d^4$$
 (2) d^9

(3) d^7

143. Which of the following can be used as the halide component for Friedel-Crafts reaction?

(Chloroethene

- (2) Isopropyl chloride
- (3) Chlorobenzene
- (4) Bromobenzene

144. In which of the following molecules, all atoms are coplanar?

$$\begin{array}{cccc} \text{(1)} & \begin{array}{cccc} \text{CH}_3 & \text{(2)} \\ \text{CH}_3 & \text{C} & \text{(2)} \end{array} \end{array}$$



145. Which one of the following structures represents nylon 6,6 polymer?

(1)
$$\begin{pmatrix} H_2 & H_3 & COOH \\ & & & & & & & & & & & & & & & & \\ & & & & & & & & & & & & & \\ & & & & & & & & & & & & & \\ & & & & & & & & & & & & & \\ & & & & & & & & & & & & \\ & & & & & & & & & & & & \\ & & & & & & & & & & & & \\ & & & & & & & & & & & & \\ & & & & & & & & & & & & \\ & & & & & & & & & & & & \\ & & & & & & & & & & & \\ & & & & & & & & & & & \\ & & & & & & & & & & & \\ & & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & & \\ & & & & & & & & & \\ & & & & & & & & & \\ & & & & & & & & & \\ & & & & & & & & & \\ & & & & & & & & & \\ & & & & & & & & \\ & & & & & & & & \\ & & & & & & & & \\ & & & & & & & & \\ & & & & & & & & \\ & & & & & & & & \\ & & & & & & & & \\ & & & & & & \\ & & & & & & & \\ & & & & & \\ & & & & & & \\ & & & & & \\ & & & & & & \\ & & & & & \\ & &$$

(2)
$$\left(\begin{array}{c} 0 & H_2 & H_2 \\ C & C & N + CH_2 \\ H_2 & 0 \end{array}\right)_n$$

(3)
$$\begin{pmatrix} H_2 & H_2 & H_2 \\ C & C & C \\ NH_2 & CH_3 \end{pmatrix}_{6}$$

$$(4) \quad \begin{pmatrix} H_2 & H_2 & H_2 \\ C & C & C \\ & NH_2 & NH_2 \end{pmatrix}_{66}$$

JMD**/E3**

15

146. In pyrrole

the electron density is maximum on

- (1) 2 and 4
- (2) 2 and 5
- (3) 2 and 3
- (4) 3 and 4
- 147. Which of the following compounds shall **not** produce propene by reaction with HBr followed by elimination or direct only elimination reaction?

- (2) H₃C—C—CH₂Br
- $\begin{array}{ccc} \text{(3)} & \text{H}_2\text{C} & \text{CH}_2 \\ & \text{H}_2 & \end{array}$
- (4) H₃C—C—CH₂OH
- 148. Which one of the following nitro-compounds does not react with nitrous acid?

$$(2) \quad H_3C \bigvee_{O}^{CH_3}_{H \setminus_{NO_2}}$$

(4)
$$H_3C$$
 CH
 C
 NO


149. The central dogma of molecular genetics states that the genetic information flows from

- (2) DNA \rightarrow RNA \rightarrow Carbohydrates
- (3) Amino acids \rightarrow Proteins \rightarrow DNA
- (4) DNA → Carbohydrates → Proteins
- **150.** The **correct** corresponding order of names of four aldoses with configuration given below

respectively, is

- (1) L-erythrose, L-threose, D-erythrose, D-threose
- D-erythrose, D-threose, L-erythrose, L-threose
- (3) L-erythrose, L-threose, L-erythrose, D-threose
- (4) D-threose, D-erythrose, L-threose, L-erythrose
- 151. In the given reaction

the product P is



r genetics tion flows

s

NA

teins

of names of iven below

CHO
H—OH
O—H

ĊH₂OH

-erythrose,

-erythrose,

-erythrose,

L-threose,

152. A given nitrogen-containing aromatic compound A reacts with Sn/HCl, followed by HNO₂ to give an unstable compound B. B, on treatment with phenol, forms a beautiful coloured compound C with the molecular formula C₁₂H₁₀N₂O. The structure of compound A is

153. Consider the reaction

 $\mathsf{CH_3CH_2CH_2Br} + \mathsf{NaCN} \to \mathsf{CH_3CH_2CH_2CN} + \mathsf{NaBr}$

This reaction will be the fastest in

- (1) N, N'-dimethylformamide (DMF)
- (2) water
- (3) ethanol
- (4) methanol

154. The **correct** structure of the product A formed in the reaction

$$\frac{\text{H}_2 \text{ (gas, 1 atmosphere)}}{\text{Pd/carbon, ethanol}} A$$

is



JMD**/ЕЗ**

(4)

17

aromatic 155. Which among the given molecules can exhibit tautomerism?

- (1) Both I and II
- (2) Both II and III
- (3) III only
- (4) Both I and III

156. The **correct** order of strengths of the carboxylic acids

is

- (1) III > II > I
- (2) II > I > III
- (3) I > II > III
- (4) II > III > I

157. The compound that will react most readily with gaseous bromine has the formula

- (1) C_4H_{10}
- (2) C₂H₄
- (3) C_3H_6
- (4) C_2H_2

- shows the presence of intramolecular hydrogen bond?
 - (1) Cellulose
 - (2) Concentrated acetic acid
 - (3) H_2O_2
 - (4) HCN
- 159. The molar conductivity of a $0.5 \text{ mol}/\text{dm}^3$ solution of AgNO3 with electrolytic of $5.76 \times 10^{-3} \text{ S cm}^{-1}$ conductivity 298 K is
 - (1) $0.086 \text{ S cm}^2/\text{mol}$
 - (2) 28.8 S cm²/mol
 - (3) $2.88 \text{ S cm}^2/\text{mol}$
 - (4) 11.52 S cm²/mol
- 160. The decomposition of phosphine (PH₃) on tungsten at low pressure is a first-order reaction. It is because the
 - (1) rate is independent of the surface coverage
 - (2) rate of decomposition is very slow
 - (3) rate is proportional to the surface coverage
 - (4) rate is inversely proportional to the surface coverage
- 161. The coagulation values in millimoles per litre of the electrolytes used for the coagulation of As₂S₃ are given below:
 - I. (NaCl) = 52,
- II. $(BaCl_2) = 0.69$,
- III. (MgSO₄) = 0.22

The correct order of their coagulating power is

- (1) III > II > I
- (2) III > I > II
- (3) I > II > III
- (4) II > I > III
- 162. During the electrolysis of molten sodium chloride, the time required to produce 0.10 mol of chlorine gas using a current of 3 amperes is
 - (1) 220 minutes
 - (2) 330 minutes
 - (3) 55 minutes
 - (4) 110 minutes

- 158. Which one of the following compounds | 163. How many electrons can fit in the orbital for which n = 3 and l = 1?
 - (1) 10
- (2) 14
- (3) 2
- (4) 6
- 164. For a sample of perfect gas when its pressure is changed isothermally from p_i to p_f , the entropy change is given by
 - (1) $\Delta S = nRT \ln \left(\frac{p_f}{p_i} \right)$
 - (2) $\Delta S = RT \ln$
 - (3) $\Delta S = nR \ln \left| \frac{p_f}{r} \right|$
 - (4) $\Delta S = nR \ln \left(\frac{p_i}{p_f} \right)$
- 165. The van't Hoff factor (i) for a dilute aqueous solution of the strong electrolyte barium hydroxide is
 - (1) 2
- (2) 3
- (3) 0
- (4) 1
- 166. The percentage of pyridine (C_5H_5N) that forms pyridinium ion ($C_5H_5N^+H$) in a 0.10~M (K_{b}) aqueous pyridine solution $C_5H_5N = 1.7 \times 10^{-9}$) is
 - (1) 0.77%
 - (2) 1.6%
 - (3) 0.0060%
 - (4) 0.013%
- 167. In calcium fluoride, having the fluorite structure, the coordination numbers for calcium ion (Ca²⁺) and fluoride ion (F⁻) are
 - (1) 8 and 4
 - (2) 4 and 8
 - (3) 4 and 2
 - (4) 6 and 6
- **168.** If the E_{cell}° for a given reaction has a negativ value, which of the following gives th correct relationships for the values of ΔC and K_{eq} ?
 - (1) $\Delta G^{\circ} < 0$; $K_{\text{eq}} > 1$
 - (2) $\Delta G^{\circ} < 0$; $K_{eq} < 1$
 - (3) $\Delta G^{\circ} > 0$; $K_{eq} < 1$
 - (4) $\Delta G^{\circ} > 0$; $K_{eq} > 1$

e orbital for

its pressure

to p_f , the

169. Which one of the following is incorrect for 175. Zinc can be coated on iron to produce ideal solution?

- (1) $\Delta P = P_{\text{obs}} P_{\text{calculated by Raoult's law}} = 0$
- (2) $\Delta G_{\text{mix}} = 0$
- (3) $\Delta H_{\text{mix}} = 0$
- (4) $\Delta U_{\text{mix}} = 0$

170. The solubility of AgCl (s) with solubility product 1.6×10^{-10} in 0.1 M NaCl solution

- (1) $1.6 \times 10^{-11} M$
- (2) zero
- (3) 1.26×10^{-5} M
- (4) $1.6 \times 10^{-9} M$

ate aqueous 171. Suppose the elements X and Y combine to 177. The hybridizations of atomic orbitals of form two compounds XY_2 and X_3Y_2 . When 0.1 mole of XY_2 weighs 10 g and 0.05 mole of X_3Y_2 weighs 9 g, the atomic weights of X and Y are

- C5H5N) that (1) 20, 30
- (2) 30, 20
- (3) 40, 30
- (4) 60, 40

in a 0.10 M (K_b) for

the fluorite

numbers for on (F) are

as a negative

g gives the

alues of ΔG

172. The number of electrons delivered at the cathode during electrolysis by a current of 1 ampere in 60 seconds is (charge on electron = 1.60×10^{-19} C)

- (1) 3.75×10^{20}
- (2) 7.48×10^{23}
- (3) 6×10^{23}
- (4) 6×10^{20}

73. Boric acid is an acid because its molecule

- (1) accepts OH from water releasing proton
- with proton from water (2) combines molecule
- (3) contains replaceable H⁺ion
- (4) gives up a proton

74. AlF₃ is soluble in HF only in presence of KF. It is due to the formation of

- (1) AlH₃
- (2) $K[AIF_3H]$
- (3) $K_3[A1F_3H_3]$
- (4) $K_3[A1F_6]$

MD/E3

19

galvanized iron but the reverse is not possible. It is because

- (1) zinc has lower negative electrode potential than iron
- (2) zinc has higher negative electrode potential than iron
- (3) zinc is lighter than iron
- (4) zinc has lower melting point than iron

176. The suspension of slaked lime in water is known as

- (1) milk of lime
- (2) aqueous solution of slaked lime
- (3) limewater
- (4) quicklime

nitrogen in NO₂⁺, NO₃⁻ and NH₄⁺ respectively

- (1) sp, sp^2 and sp^3
- (2) sp^2 , sp and sp^3
- (3) sp, sp^3 and sp^2
- (4) sp^2 , sp^3 and sp

178. Which of the following fluoro-compounds is 35.54 most likely to behave as a Lewis base?

- (1) CF₄ (3) BF₃
- (2) SiF₄
- (4) PF₃
- 82.446

179. Which of the following pairs of ions isoelectronic and isostructural? 17423

- (1) SO_3^{2-} , NO_3^{-}
- (2) $C10_3^-$, $S0_3^{2-}$
- 2) 40

(3) CO_3^{2-} , NO_3^{-}

- (4) CIO_3^- , CO_3^{2-}

16+22

14+49 61 @1012+50 180. In context with beryllium, which one of the following statements is incorrect? 12+46

- (1) Its salts rarely hydrolyze.
- 358
- (2) Its hydride is electron-deficient and polymeric. 35+
- (3) It is rendered passive by nitric acid.
- JAY It forms Be 2C.

Bezel

AglaSem Admission