

Lecture 1

Tuesday 28th October, 2014

Contents

1	General Information	2
1.1	Office Hours	2
2	Functions	3
2.1	Notation	3
2.2	Definitions	3
2.2.1	Domain, Range and Variables	3
2.2.2	Image of a function	3
2.2.3	Existence domain	3
2.2.4	Graph	3
2.2.5	Even function	3
2.2.6	Odd function	4
2.2.7	Periodical function	4
2.2.8	Shifting with respect to y -axis	4
2.2.9	Shifting with respect to x -axis	4
2.2.10	Monotonic function	4
2.2.11	One-to-one function	4

1 General Information

Dr. Yakov Yakubov

1.1 Office Hours

Monday

16:15 - 17:15

Room 233, MathBuilding

Tel: 03-6405357

2 Functions

2.1 Notation

$$\mathbb{N} = \text{Set of all natural numbers} \quad (1)$$

$$\mathbb{Z} = \text{Set of all integers} \quad (2)$$

$$\mathbb{N} = \left\{ \frac{m}{n} : m \in \mathbb{Z}, n \in \mathbb{N} \right\} \quad (3)$$

2.2 Definitions

2.2.1 Domain, Range and Variables

Let D and E be two sets of real numbers. A function f from D into E is a well defined law which, to each $x \in D$ corresponds to a unique number $y \in E$. The set D is called the domain of f and the set E is called the range of f .

Denote $f : D \rightarrow E$ or $y = f(x)$.

The variable x is called independent variable and the variable y is called dependent variable.

The variable x is also called the origin of y and y is also called the image of x .

2.2.2 Image of a function

Given $f : D \rightarrow E$. Then the image of f is a set of all $y \in E$ s.t. $\exists x \in D, y = f(x) : I(f) = \{y \in E : \exists x \in D, y = f(x)\}$

2.2.3 Existence domain

The biggest possible domain of a function f is called the existence domain of f .

2.2.4 Graph

A set of point $\{(x, f(x)) : x \in D\}$ in the plane \mathbb{R}^2 is called a graph of a function $y = f(x)$.

2.2.5 Even function

If $f(-x) = f(x); (x, -x \in D)$ then, f is called an even function.

Each even function is symmetric about the y -axis.

2.2.6 Odd function

If $f(-x) = -f(x)$; $(x, -x \in D)$ then, f is called an odd function.
Each odd function is symmetric about the origin.

2.2.7 Periodical function

A function $y = f(x)$ which is defined on D is called periodical if $\exists T \neq 0$ which is called a period of f s.t. $\forall x \in D \Rightarrow x + T \in D$ and $f(x + T) = f(x)$.
The smallest such $T > 0$ (if it exists) is called the minimal period.

2.2.8 Shifting with respect to y -axis

$f(x+a)$ is the graph of $f(x)$, shifted by a , in the direction of the x -axis, opposite to the sign of a .

2.2.9 Shifting with respect to x -axis

$f(x) + a$ is the graph of $f(x)$, shifted by a , in the direction of the y -axis, according to the sign of a .

2.2.10 Monotonic function

A function $y = f(x)$ is called monotonic increasing (strongly increasing) in D if $\forall x_1, x_2 \in D, x_1 < x_2 \Rightarrow f(x_1) \leq f(x_2)$ ($f(x_1) < f(x_2)$).

A function $y = f(x)$ is called monotonic decreasing (strongly decreasing) in D if $\forall x_1, x_2 \in D, x_1 > x_2 \Rightarrow f(x_1) \geq f(x_2)$ ($f(x_1) > f(x_2)$).

2.2.11 One-to-one function

A function $f : D(f) \rightarrow E$ is called one-to-one if $\forall y \in I(f) \Rightarrow \exists! x \in D(f)$ s.t. $y = f(x)$.

Equivalently, $\forall x_1, x_2 \in D(f)$, if $f(x_1) = f(x_2) \Rightarrow x_1 = x_2$.