

# Lecture 2

Thursday 30<sup>th</sup> October, 2014

## Contents

<b>1</b>	<b>Functions</b>	<b>2</b>
1.1	Continuity . . . . .	2
1.2	Elementary Operations between Functions . . . . .	2
1.2.1	$h = f \pm g$ . . . . .	2
1.2.2	$h = kf$ . . . . .	2
1.2.3	$h = fg$ . . . . .	2
1.2.4	$h = \frac{f}{g}$ . . . . .	2
1.3	Composite Functions . . . . .	2
1.4	Elementary Functions . . . . .	3
1.4.1	Polynomial . . . . .	3
1.4.2	Power Function . . . . .	3
1.4.3	Exponential Function . . . . .	3
1.4.4	Logarithmic Function . . . . .	3
1.4.5	Trigonometric Functions . . . . .	3
1.4.6	Inverse Trigonometric Functions . . . . .	4
1.4.7	Hyperbolic Functions . . . . .	4
1.4.8	Absolute Value . . . . .	5
1.4.9	Floor Function . . . . .	5

# 1 Functions

## 1.1 Continuity

If  $f : D(f) \rightarrow I(f)$  is one-to-one and onto. Then, we can define  $g : I(f) \rightarrow D(f)$ , which is one-to-one and onto, by  $g(y) = x$ , where  $y = f(x)$ . Therefore,  $g(f(x)) = x$ .  $g$  is called the inverse function of  $f$ .

The inverse function is denoted as  $g = f^{-1}$  (Note:  $f^{-1} \neq \frac{1}{f}$ )

$$D(f) = I(f^{-1}) \quad (1)$$

$$I(f) = D(f^{-1}) \quad (2)$$

The graphs of a  $f$  and  $f^{-1}$  are symmetric about  $y = x$ .

## 1.2 Elementary Operations between Functions

### 1.2.1 $h = f \pm g$

$$h(x) = f(x) \pm g(x) \quad (3)$$

$$D(h) = D(f) \cap D(g) \quad (4)$$

### 1.2.2 $h = kf$

$$h(x) = kf(x) \quad (5)$$

$$D(h) = D(f) \quad (6)$$

### 1.2.3 $h = fg$

$$h(x) = f(x)g(x) \quad (7)$$

$$D(h) = D(f) \cap D(g) \quad (8)$$

### 1.2.4 $h = \frac{f}{g}$

$$h(x) = \frac{f(x)}{g(x)} \quad (9)$$

$$D(h) = \{x \in D(f) \cap D(g) : g(x) \neq 0\} \quad (10)$$

## 1.3 Composite Functions

Let  $f : D(f) \rightarrow E$  and  $g : D(g) \rightarrow F$  be two functions. A composition of  $f$  with  $g$  is a function  $h : D(h) \rightarrow F$  where  $h(x) = g(f(x))$ . It is denoted as  $g \circ f$

$$D(h) = \{x \in D(f) : f(x) \in D(g)\} \quad (11)$$

## 1.4 Elementary Functions

### 1.4.1 Polynomial

$$y = f(x) = a_0 + a_1x + \cdots + a_nx^n; a_0, \dots, a_n \in \mathbb{R} \quad (12)$$

$$D(f) = \mathbb{R} \quad (13)$$

1. If  $n = 0, y = f(x) = a_0$  represents a constant function.
2. If  $n = 1, y = f(x) = a_0 + a_1x$  represents a straight line in the  $X - Y$  plane.
3. If  $n = 2, y = f(x) = a_0 + a_1x + a_2x^2$  represents a parabola in the  $X - Y$  plane.

### 1.4.2 Power Function

$$y = f(x) = x^a; a \in \mathbb{R} \quad (14)$$

$$D(f) \text{ depends on } a \quad (15)$$

### 1.4.3 Exponential Function

$$y = f(x) = a^x; a > 0, a \neq 1 \quad (16)$$

$$D(f) = \mathbb{R} \quad (17)$$

$$I(f) = (0, \infty) \quad (18)$$

### 1.4.4 Logarithmic Function

$$y = f(x) = \log_a x \quad (19)$$

$$D(f) = (0, \infty) \quad (20)$$

$$I(f) = \mathbb{R} \quad (21)$$

### 1.4.5 Trigonometric Functions

$$y = f(x) = \sin x \quad (22)$$

$$y = f(x) = \cos x \quad (23)$$

$$y = f(x) = \tan x = \frac{\sin x}{\cos x} \quad (24)$$

$$y = f(x) = \cot x = \frac{1}{\tan x} = \frac{\cos x}{\sin x} \quad (25)$$

$$y = f(x) = \csc x = \frac{1}{\sin x} \quad (26)$$

$$y = f(x) = \sec x = \frac{1}{\cos x} \quad (27)$$

#### 1.4.6 Inverse Trigonometric Functions

$$y = f^{-1}(x) = \sin^{-1} x = \arcsin x \quad (28)$$

$$D(\arcsin x) = [-1, 1] \quad (29)$$

$$I(\arcsin x) = \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \quad (30)$$

$$y = f^{-1}(x) = \cos^{-1} x = \arccos x \quad (31)$$

$$D(\arccos x) = [-1, 1] \quad (32)$$

$$I(\arccos x) = [0, \pi] \quad (33)$$

$$y = f^{-1}(x) = \tan^{-1} x = \arctan x \quad (34)$$

$$D(\arctan x) = \mathbb{R} \quad (35)$$

$$I(\arctan x) = \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \quad (36)$$

#### 1.4.7 Hyperbolic Functions

$$\sinh x \doteq \frac{e^x - e^{-x}}{2} \quad (37)$$

$$D(\sinh x) = \mathbb{R} \quad (38)$$

$$I(\sinh x) = \mathbb{R} \quad (39)$$

$$\cosh x \doteq \frac{e^x + e^{-x}}{2} \quad (40)$$

$$D(\cosh x) = \mathbb{R} \quad (41)$$

$$I(\cosh x) = [1, \infty) \quad (42)$$

$$\tanh x \doteq \frac{\sinh x}{\cosh x} = \frac{e^x - e^{-x}}{e^x + e^{-x}} \quad (43)$$

$$D(\tanh x) = \mathbb{R} \quad (44)$$

$$I(\tanh x) = (-1, 1) \quad (45)$$

#### 1.4.7.1 Identities of Hyperbolic Functions

$$\sinh(2x) = 2 \sinh x \cosh x \quad (46)$$

$$\cosh^2 x + \sinh^2 x = \cosh(2x) \quad (47)$$

$$\cosh^2 x - \sinh^2 x = 1 \quad (48)$$

$$\frac{\cosh(2x) - 1}{2} = \sinh^2 x \quad (49)$$

$$\frac{\cosh(2x) + 1}{2} = \cosh^2 x \quad (50)$$

#### 1.4.8 Absolute Value

$$y = f(x) = \begin{cases} x; & x > 0 \\ 0; & x = 0 \\ -x; & x < 0 \end{cases} \quad (51)$$

#### 1.4.9 Floor Function

$$y = f(x) = \lfloor x \rfloor = \text{the largest integer less than or equal to } x \quad (52)$$