

Lecture 21

Aakash Jog


Thursday 8th January, 2015

Contents


1	Forced Oscillations	2
2	Accelerating Systems	3

1 Forced Oscillations

Example 1. Find



Solution.



$$m\ddot{x} + \frac{k}{m}x = \frac{F_0}{m} \cos \omega t$$

$$\therefore \ddot{x} + \frac{k}{m}x = \frac{F_0}{m} \cos \omega t$$

Therefore, solving

$$x = A \cos \omega_0 t + B \sin \omega_0 t + \frac{F_0}{k - m\omega^2} \cos \omega t$$

$$\therefore \dot{x} = \omega_0(-A \sin \omega_0 t + B \cos \omega_0 t) - \frac{F_0}{k - m\omega^2} \omega \sin \omega t$$

Substituting initial conditions,

$$x = \frac{\frac{F_0}{m}}{\frac{k}{m} - \omega^2} (-\cos \omega_0 t + \cos \omega t)$$

Let $\frac{F_0}{m} = f_0$

$$\therefore x = -\frac{2f_0}{\omega_0^2 - \omega^2} \sin\left(\frac{\omega t - \omega_0 t}{2}\right) \sin\left(\frac{\omega t + \omega_0 t}{2}\right)$$

$\omega - \omega_0 = \Delta\omega$ and $\omega + \omega_0 \approx 2\omega_0$

$$\therefore x \approx \frac{2f_0}{\Delta\omega \cdot 2\omega_0} \sin\left(\frac{\Delta\omega}{2}t\right) \cdot \sin(\omega_0 t)$$