

Physics 2 : Recitations

Aakash Jog

2014-15

Contents

1	Instructor Information	3
I	Electrostatics	4
1	Gravitation and Electromagnetism	4
2	Coulomb's Law	4
	Recitation 1 – Exercise 1.	4
	Recitation 1 – Solution 1.	4
	Recitation 1 – Exercise 2.	5
	Recitation 1 – Solution 2.	5
	Recitation 1 – Exercise 3.	6
	Recitation 1 – Solution 3.	6
	Recitation 1 – Exercise 4.	6
	Recitation 1 – Solution 4.	6
	Recitation 1 – Exercise 5.	6
	Recitation 1 – Solution 5.	7



This work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>.

3	Gauss' Law	9
	Recitation 2 – Exercise 2.	9
	Recitation 2 – Solution 2.	10
	Recitation 2 – Exercise 3.	11
	Recitation 2 – Solution 3.	11
	Recitation 2 – Exercise 4.	12
	Recitation 2 – Solution 4.	12
4	Electric Potential	12
	Recitation 3 – Exercise 2.	12
	Recitation 3 – Solution 2.	13
	Recitation 3 – Exercise 3.	14
	Recitation 3 – Solution 3.	14
	Recitation 3 – Exercise 4.	15
	Recitation 3 – Solution 4.	15
	Recitation 4 – Exercise 1.	17
	Recitation 4 – Solution 1.	17
	Recitation 4 – Exercise 3.	18
	Recitation 4 – Solution 3.	18
	Recitation 4 – Exercise 4.	19
	Recitation 4 – Solution 4.	19
	Recitation 4 – Exercise 5.	19
	Recitation 4 – Solution 5.	20

1 Instructor Information

Dr. Richard Spitzberg

Office: Ma Aabadot 119

E-mail: rms9999@gmail.com

Part I

Electrostatics

1 Gravitation and Electromagnetism

Gravitation

$$F_G = G \frac{m_1 m_2}{r^2}$$

$$G = 6.7 \times 10^{-11} \text{N m}^2 \text{kg}^{-2}$$

Electromagnetism

$$F_E = k \frac{q_1 q_2}{r^2}$$

$$8.99 \times 10^9 \text{N m}^2 \text{C}^{-2}$$


2 Coulomb's Law

Recitation 1 – Exercise 1.

Four identical charges q are placed in the corners of a square of length a . A fifth charge Q is free to move along the straight line perpendicular to the square plane and passing through its centre. When the charge Q is in the same plane as the other charges, all the forces in the system cancel out.

1. Calculate Q for a given q and a .
2. Find the force $\overrightarrow{F(z)}$ acting on the charge Q when it is at height z above the square.

Recitation 1 – Solution 1.



Consider q on the top right corner of the square. The total force acting on it is 0. Therefore

$$0 = \frac{kq^2}{a^2} \cos \frac{\pi}{4} + \frac{kq^2}{a^2} \cos \frac{\pi}{4} + \frac{kq^2}{2a^2} + \frac{2kQq}{a^2}$$

$$\therefore Q = -\frac{1 + 2\sqrt{2}}{4}q$$

If Q is at a height z from the plane, the distance between each q and Q is

$$r = \sqrt{z^2 + \left(\frac{a}{\sqrt{2}}\right)^2}$$

Therefore the force of each q on Q is $\frac{kQq}{r^2}$.

Due to symmetry, the components of the forces in the z direction will add up, and all other components will cancel out.

Let the angle between the z direction and the line joining q and Q be φ .

Therefore, the net force is

$$F = 4 \frac{kQq}{r^2} \cos \varphi$$

$$= 4 \frac{kQq}{r^2} \frac{z}{r}$$

$$= 4 \frac{kQq}{z^2 \left(1 + \frac{a^2}{2z^2}\right)^{3/2}}$$

Recitation 1 – Exercise 2.

1. A wire of length 3 metre is charged with 2 C m^{-1} . What is the wire's total charge?

Recitation 1 – Solution 2.

$$\lambda = \frac{Q}{L}$$

$$\therefore Q = L\lambda$$

$$= 6\text{C}$$

Recitation 1 – Exercise 3.

A wire of length L has the following charge distribution: $\lambda = \lambda_0 \cos \frac{\pi x}{L}$, where x is the distance from the wire's edge. What is the wire's total charge?

Recitation 1 – Solution 3.

$$\begin{aligned}\lambda &= \frac{dq}{dx} \\ \therefore \frac{dq}{dx} &= \lambda_0 \cos \frac{\pi x}{L} \\ \therefore q &= \int_0^L \lambda_0 \cos \frac{\pi x}{L} dx \\ &= 0\end{aligned}$$

Recitation 1 – Exercise 4.

A hollow sphere of radius R is uniformly charged with a charge Q . Calculate the charge distribution on the surface of the sphere.

Recitation 1 – Solution 4.


$$\begin{aligned}\sigma &= \frac{Q}{A} \\ &= \frac{Q}{4\pi R^2}\end{aligned}$$

Recitation 1 – Exercise 5.

A straight thin wire is uniformly charged with distribution λ . A charge q is positioned at distance y_1 beneath the wire and r away from it.

1. Find the force acting on the charge q .
2. Show that when the charge is positioned in front of the centre of the wire the \hat{y} component of the force is cancelled.
3. Calculate the force an infinite straight wire will exert on the charge q .

Recitation 1 – Solution 5.



Consider an elemental charge dQ of length dy , at distance y as shown. Let the angle between the line joining dQ and q and the y direction be θ .

$$F_y = F \cos \theta$$

$$F_z = F \sin \theta$$

Let

$$a = L + y_1$$

$$\therefore R = \sqrt{r^2 + (a - y)^2}$$

Therefore,

$$\cos \theta = \frac{a - y}{R}$$

$$\sin \theta = \frac{r}{R}$$

Therefore,

$$\begin{aligned} F_y &= kq \int_0^L \frac{\lambda dy}{R^2} \frac{(a - y)}{R} \\ &= kq\lambda \int_0^L \frac{dy(a - y)}{((a - y)^2 + r^2)^{3/2}} \\ &= kq\lambda \left(\frac{1}{\sqrt{y_1^2 + r^2}} - \frac{1}{\sqrt{a^2 + r^2}} \right) \end{aligned}$$

$$\begin{aligned}
F_z &= kq \int_0^L \frac{\lambda \, dy}{R^2} \frac{r}{R} \\
&= kq\lambda \int_0^L \frac{r \, dy}{((a-y)^2 + r^2)^{3/2}} \\
&= \frac{kq\lambda}{r} \left(\frac{a}{\sqrt{r^2 + a^2}} - \frac{y_1}{\sqrt{r^2 + y_1^2}} \right)
\end{aligned}$$

When the charge is positioned above the centre of the wire,

$$\begin{aligned}
y_1 &= -\frac{L}{2} \\
\therefore a &= \frac{L}{2}
\end{aligned}$$

Therefore,

$$\begin{aligned}
F_y &= kq\lambda \left(\frac{1}{\sqrt{y_1^2 + r^2}} - \frac{1}{\sqrt{a^2 + r^2}} \right) \\
&= kq\lambda \left(\frac{1}{\sqrt{-\frac{L^2}{4} + r^2}} - \frac{1}{\sqrt{\frac{L^2}{4} + r^2}} \right) \\
&= 0
\end{aligned}$$

$$\begin{aligned}
F_z &= \frac{kq\lambda}{r} \left(\frac{a}{\sqrt{r^2 + a^2}} - \frac{y_1}{\sqrt{r^2 + y_1^2}} \right) \\
&= \frac{kq\lambda}{r} \left(\frac{\frac{L}{2}}{\sqrt{r^2 + \frac{L^2}{2}}} - \frac{-\frac{L}{2}}{\sqrt{r^2 + -\frac{L^2}{2}}} \right) \\
&= \frac{kq\lambda}{r} \left(\frac{L}{\sqrt{r^2 + \frac{L^2}{2}}} \right) \\
&= \frac{kq\lambda}{r} \left(\frac{1}{\sqrt{\frac{1}{4} + \left(\frac{r}{L}\right)^2}} \right)
\end{aligned}$$

If the line is infinite, $L \rightarrow \infty$. Therefore

$$\begin{aligned}
F_z &= \frac{kq\lambda}{r} \left(\frac{1}{\sqrt{\frac{1}{4} + \left(\frac{r}{L}\right)^2}} \right) \\
&= \frac{2kq\lambda}{r}
\end{aligned}$$

3 Gauss' Law

Recitation 2 – Exercise 2.

A ball of radius a is charged with distribution $\rho = \rho_0 \frac{r}{a}$. Find the electric field everywhere.

Recitation 2 – Solution 2.

Consider a spherical Gaussian surface of radius r .

If $r \leq a$, the charge in the interior of the Gaussian surface is

$$\begin{aligned} q(r) &= \int_0^r \frac{\rho_0 r}{a} \cdot 4\pi r^2 \, dr \\ &= \frac{\rho_0}{a} \pi r^4 \end{aligned}$$

Therefore, by Gauss' Law,

$$\begin{aligned} E \cdot 4\pi r^2 &= \frac{q(r)}{\varepsilon_0} \\ \therefore E &= \frac{\rho_0 \pi r^4}{4\pi a r^2} \\ &= \frac{\rho_0 r^2}{4a\varepsilon_0} \end{aligned}$$

If $r \geq a$, the entire ball of charge is in the interior of the Gaussian surface.

Therefore,

$$\begin{aligned} Q &= q(a) \\ &= \frac{\rho_0}{a} \cdot \pi a^4 \\ &= \rho_0 \pi a^3 \end{aligned}$$

Therefore, by Gauss' Law,

$$\begin{aligned} E \cdot 4\pi r^2 &= \frac{Q}{\varepsilon_0} \\ \therefore E &= \frac{Q}{4\pi r^2 \varepsilon_0} \\ &= \frac{\rho_0 a^3}{4r^2 \varepsilon_0} \end{aligned}$$

Therefore,

$$E = \begin{cases} \frac{\rho_0 r^2}{4a\varepsilon_0} & ; \quad r \leq a \\ \frac{\rho_0 a^3}{4r^2 \varepsilon_0} & ; \quad r \geq a \end{cases}$$

Recitation 2 – Exercise 3.

An infinitely long cylinder of radius a is charged with distribution $\rho = \rho_0 \frac{r}{a}$. Find the electric field everywhere.

Recitation 2 – Solution 3.

Consider a infinite cylindrical Gaussian surface with radius r . If $r \leq a$, the charge in the interior of the Gaussian surface is

$$\begin{aligned} q(r) &= \int_0^r \frac{\rho_0 r}{a} \pi r^2 \, dr \\ &= \frac{2\pi\rho_0 L r^3}{3a} \end{aligned}$$

Therefore, by Gauss' Law,

$$\begin{aligned} E \cdot 2\pi r L &= \frac{2\pi\rho_0 L r^3}{3a\varepsilon_0} \\ \therefore E &= \frac{\rho_0 r^2}{3a\varepsilon_0} \end{aligned}$$

If $r \geq a$, the entire cylinder of charge is in the interior of the Gaussian surface. Therefore,

$$\begin{aligned} Q &= q(a) \\ &= \frac{2\pi\rho_0 L a^3}{3a} \\ &= \frac{2\pi\rho_0 L a^2}{3} \end{aligned}$$

Therefore, by Gauss' Law,

$$\begin{aligned} E \cdot 2\pi r L &= \frac{2\pi\rho_0 L a^2}{3\varepsilon_0} \\ \therefore E &= \frac{\rho_0 a^2}{3\varepsilon_0 r} \end{aligned}$$

Therefore,


$$E = \begin{cases} \frac{\rho_0 r^2}{3a\varepsilon_0} & ; \quad r \leq a \\ \frac{\rho_0 a^2}{3\varepsilon_0 r} & ; \quad r \geq a \end{cases}$$

Recitation 2 – Exercise 4.

Find the electric field due to a thin infinite plane of uniform charge distribution σ .

Recitation 2 – Solution 4.

Consider a cylindrical Gaussian surface, with ends of area A , as shown.



The charge in the interior of the surface is

$$dq = A\sigma$$


Therefore, by Gauss' Law,

$$\begin{aligned} E_1 \cdot A_1 + E_2 \cdot A_2 &= \frac{A\sigma}{\varepsilon_0} \\ \therefore 2EA &= \frac{A\sigma}{\varepsilon_0} \\ \therefore E &= \frac{\sigma}{2\varepsilon_0} \end{aligned}$$

4 Electric Potential

Recitation 3 – Exercise 2.


A system of four charges is constructed as shown.



- a) Calculate the work needed to build this system.
- b) What is the potential in the centre of the system?
- c) Calculate the potential in each of the corners (calculate as if there is no charge in the corner you are calculating for).

Recitation 3 – Solution 2.

- a) Let the positions of the charges be A, B, C, D.



The work done to bring the first charge from infinity to A is

$$W_A = 0$$

The work done to bring the first charge from infinity to B is

$$W_B = \frac{1}{4\pi\epsilon_0} \frac{q^2}{a\sqrt{2}}$$

Similarly for the other two charges.

Therefore,

$$\begin{aligned} W &= 0 + \frac{q^2}{4\sqrt{2}\pi\epsilon_0} + \frac{-2q^2}{4\pi\epsilon_0 a} + \left(\frac{-2q^2}{4\pi\epsilon_0 a} + \frac{q^2}{4\sqrt{2}\pi\epsilon_0} \right) \\ &= \frac{q^2}{2\sqrt{2}\pi\epsilon_0} - \frac{q^2}{\pi\epsilon_0 a} \end{aligned}$$

b)

$$\begin{aligned} V_{\text{centre}} &= V_q + V_q + V_{-q} + V_{-q} \\ &= \frac{q}{4\pi\epsilon_0 \left(\frac{a}{\sqrt{2}}\right)} + \frac{q}{4\pi\epsilon_0 \left(\frac{a}{\sqrt{2}}\right)} - \frac{q}{4\pi\epsilon_0 \left(\frac{a}{\sqrt{2}}\right)} - \frac{q}{4\pi\epsilon_0 \left(\frac{a}{\sqrt{2}}\right)} \\ &= 0 \end{aligned}$$

c)

$$\begin{aligned} V_A &= \frac{1}{4\pi\epsilon_0} \frac{-q}{a} + \frac{1}{4\pi\epsilon_0} \frac{-q}{a} + \frac{1}{4\pi\epsilon_0} \frac{q}{\sqrt{2}a} \\ V_B &= \frac{1}{4\pi\epsilon_0} \frac{-q}{a} + \frac{1}{4\pi\epsilon_0} \frac{-q}{a} + \frac{1}{4\pi\epsilon_0} \frac{q}{\sqrt{2}a} \\ V_C &= \frac{1}{4\pi\epsilon_0} \frac{q}{a} + \frac{1}{4\pi\epsilon_0} \frac{q}{a} + \frac{1}{4\pi\epsilon_0} \frac{-q}{\sqrt{2}a} \\ V_D &= \frac{1}{4\pi\epsilon_0} \frac{q}{a} + \frac{1}{4\pi\epsilon_0} \frac{q}{a} + \frac{1}{4\pi\epsilon_0} \frac{-q}{\sqrt{2}a} \end{aligned}$$

Recitation 3 – Exercise 3.

A ring of radius R is charged with total charge Q .

- a) Calculate the electric field in the centre of the ring.
- b) Calculate the potential in the centre of the ring by integrating the contributions of the infinitesimal charge elements of the ring.

Recitation 3 – Solution 3.

- a) Due to the symmetry of the ring, the field due to every elemental charge dq will be cancelled out by the field due to a elemental charge diametrically opposite to dq .

Therefore,

$$\vec{E} = 0$$

b)

$$\mathrm{d}V = \frac{\mathrm{d}q}{4\pi\epsilon_0 R}$$
$$\therefore V = \frac{Q}{4\pi\epsilon_0 R}$$

Recitation 3 – Exercise 4.

Calculate the potential resulting from a ball charged with constant volume distribution ρ . Use the expression

$$\varphi(r_2) - \varphi(r_1) = - \int_{r_1}^{r_2} E(r) \, \mathrm{d}r$$

Repeat the calculation twice:

a) Set $\varphi(r = R) = 0$

b) Set $\varphi(r = \infty) = 0$

Recitation 3 – Solution 4.

a) Let $\varphi(r = \infty) = 0$.

If $r > R$,

$$\begin{aligned} E &= -\frac{d\varphi}{dr} \\ \therefore -\frac{d\varphi}{dr} &= \frac{Q}{4\pi\epsilon_0 r^2} \\ \therefore \int d\varphi &= \int_{\infty}^r \frac{Q}{4\pi\epsilon_0 r^2} dr \\ \therefore \varphi(r) - \varphi(\infty) &= \frac{Q}{4\pi\epsilon_0 r} - 0 \\ \therefore \varphi(r) &= \frac{Q}{4\pi\epsilon_0 r} \end{aligned}$$

If $r < R$,

$$\begin{aligned} E &= \frac{q(r)}{4\pi\epsilon_0 r^2} \\ &= \frac{\rho \cdot \frac{4}{3}\pi r^3}{4\pi\epsilon_0 r^2} \\ &= \frac{\rho r}{3\epsilon_0} \end{aligned}$$

Therefore

$$\begin{aligned} E &= -\frac{d\varphi}{dr} \\ \therefore \int d\varphi &= \int_r^R \frac{\rho}{3\epsilon_0} r dr \\ \therefore \varphi(R) - \varphi(r) &= \frac{\rho}{6\epsilon_0} (r^2 - R^2) \\ \therefore \varphi(r) &= \frac{Q}{4\pi\epsilon_0 R} + \frac{\rho(R^2 - r^2)}{6\epsilon_0} \\ &= \frac{Q}{8\pi\epsilon_0 R} \left(3 - \frac{r^2}{R^2} \right) \end{aligned}$$

b) Let $\varphi(r = R) = 0$.

$$\begin{aligned}\therefore \varphi(R) - \varphi(r) &= \frac{\rho}{6\epsilon_0} (r^2 - R^2) \\ \therefore \varphi(r) &= \frac{\rho}{6\epsilon_0} (R^2 - r^2) \\ &= \frac{Q}{8\pi R\epsilon_0} \left(1 - \frac{r^2}{R^2}\right)\end{aligned}$$

Recitation 4 – Exercise 1.

A point charge Q is surrounded by a spherical grounded shell of radius R_1 .

1. What is the charge accumulated on the shell? Where did it come from?
2. The entire system is covered with another spherical shell of radius R_2 and charged with q . What will be the charge accumulated on the grounded shell?

Recitation 4 – Solution 1.

1. The charge accumulated on the shell comes from the ground.

Let the charge on the shell be Q_1 .

As the shell is grounded, the net potential on it must be zero.

$$\begin{aligned}\varphi &= \varphi \text{ due to } Q + \varphi \text{ due to } Q' \\ \therefore 0 &= \frac{Q}{4\pi\epsilon_0 R_1} + \frac{Q_1}{4\pi\epsilon_0 R_1} \\ \therefore Q_1 &= -Q\end{aligned}$$

2. Let the charge on the grounded shell be Q_2 .

$$\begin{aligned}\varphi &= \varphi \text{ due to } Q + \varphi \text{ due to } Q_2 + \varphi \text{ due to } q \\ \therefore 0 &= \frac{Q}{4\pi\epsilon_0 R_1} + \frac{Q_2}{4\pi\epsilon_0 R_1} + \frac{q}{4\pi\epsilon_0 R_2} \\ \therefore Q_2 &= Q - \frac{qR_1}{R_2}\end{aligned}$$

Recitation 4 – Exercise 3.

A point charge Q is set in the center (same distance from all corners) of a perfect tetrahedron. The bottom face of the tetrahedron is uniformly charged with charge density σ .

Recitation 4 – Solution 3.

1. Consider a spherical Gaussian surface passing through the vertices of the tetrahedron.

Therefore by Gauss' Law, the total flux passing through the sphere, due to Q is $\oint \vec{E} \cdot d\vec{A}$.

Hence, by symmetry, the flux through every surface of the tetrahedron, due to Q is

$$\begin{aligned}\Phi &= \frac{1}{4} \oint \vec{E} \cdot d\vec{A} \\ &= \frac{1}{4} \frac{Q}{\epsilon_0}\end{aligned}$$

The flux on the bottom face due to the bottom face itself is zero. Therefore the total flux through the bottom face is the flux due to Q only.

2. As the charge on Q and the bottom face of the tetrahedron are similar in charge, the force between them is repulsive in nature. Hence, the force on the bottom face is directed downwards.
3. Let the area of the bottom face

$$\begin{aligned}F &= (\sigma A)E \\ &= \sigma(EA)\end{aligned}$$

As Q is exactly above the centre of the bottom face, due to symmetry, $\oint \vec{E} \cdot d\vec{A} = EA$

$$\begin{aligned}\therefore F &= \sigma \left(\oint \vec{E} \cdot d\vec{A} \right) \\ &= \sigma \left(\frac{Q}{4\epsilon_0} \right)\end{aligned}$$

Recitation 4 – Exercise 4.

The following electric field is given:

$$\vec{E} = \alpha y^2 \hat{i} + \alpha(2xy + z^2) \hat{j} + 2\alpha yz \hat{k}$$

Calculate the potential at $\vec{r} = (x, y, z)$, set the potential at the origin to be zero.

Recitation 4 – Solution 4.

Let $\varphi(0, 0, 0) = 0$.

Therefore,

$$\begin{aligned} E_x &= -\frac{d\varphi}{dx} \\ &= \alpha y^2 \\ \therefore \varphi &= -\alpha xy^2 + f_1(y, z) \\ E_y &= -\frac{d\varphi}{dy} \\ &= \alpha(2xy + z^2) \\ \therefore \varphi &= -\alpha xy^2 - \alpha z^2 y + f_1(x, z) \\ E_z &= -\frac{d\varphi}{dz} \\ &= 2\alpha yz \\ \therefore \varphi &= -\alpha yz^2 + f_2(x, y) \end{aligned}$$

Comparing the three expressions of φ ,

$$\varphi = -\alpha xy^2 - \alpha yz^2 + c$$

As $\varphi(0, 0, 0) = 0$, $c = 0$.

Therefore,

$$\varphi = -\alpha xy^2 - \alpha yz^2$$

Recitation 4 – Exercise 5.

A thin rod of length L is charged with a uniform charge density λ is laid along the x -axis.

1. Calculate the electric potential along the x -axis (where $x > L$)
2. Calculate the electric field along the x -axis (where $x > L$)

3. A second identical thin rod is placed along the x -axis at distance L from the edge of the first rod. Calculate the force due to left rod, acting on the right one.

Recitation 4 – Solution 5.

1. Consider an elemental charge dq with length dx at a distance x from the origin.

Therefore, the potential at a distance d from the origin is

$$\begin{aligned} d\varphi &= \frac{dq}{4\pi\epsilon_0(L+d-x)} \\ \therefore \int d\varphi &= \int_0^L \frac{\lambda dx}{4\pi\epsilon_0(L+d-x)} \\ \therefore \varphi &= \frac{\lambda}{4\pi\epsilon_0} \ln\left(\frac{L+d}{d}\right) \end{aligned}$$

2.

$$\begin{aligned} dE &= \frac{d\varphi}{dr} \\ \therefore dE &= \frac{dq}{4\pi\epsilon_0(L+d-x)^2} \\ \therefore E &= \frac{\lambda}{4\pi\epsilon_0} \int_{L+d}^d \frac{du}{u^2} \\ &= \frac{\lambda}{4\pi\epsilon_0} \left(\frac{L}{(L+d)(d)} \right) \end{aligned}$$

3. Consider an elemental charge dq with length dx , on the second rod, at a distance x from the end of the first rod.

$$\begin{aligned}
dF &= \frac{\lambda}{4\pi\epsilon_0} \left(\frac{L}{(L+x)(x)} \right) dq \\
&= \frac{\lambda}{4\pi\epsilon_0} \left(\frac{L}{(L+x)(x)} \right) \lambda dx \\
\therefore F &= \int_L^{2L} \frac{\lambda^2}{4\pi\epsilon_0} \left(\frac{L}{(L+x)(x)} \right) dx \\
&= \frac{\lambda^2}{4\pi\epsilon_0} \ln \frac{x}{L+x} \Big|_L^{2L} \\
&= \frac{\lambda^2}{4\pi\epsilon_0} \left(\ln \frac{2}{3} - \ln \frac{1}{2} \right) \\
&= \frac{\lambda^2}{4\pi\epsilon_0} \left(\ln \frac{4}{3} \right)
\end{aligned}$$