

## Chapter 2

### 1. Compute mean

Arrange data in ascending order

35, 35, 37, 39, 40, 41, 45, 55, 55, 55, 55, 55, 60, 65

$$\bar{x} = \frac{\sum x}{n} = \frac{35+35+37+39+40+41+45+55+55+55+55+55+60+65}{14} = 48$$

$$Md = \frac{n-1}{2}^{\text{th}} \text{ item} = \frac{14+1}{2} = 7.5^{\text{th}} \text{ item}$$

$$\text{Median value} = \frac{45+55}{2} = 50$$

$M_O$  = Most repeated value = 55

### 2. The number of telephone calls

No. of calls (x)	0	1	2	3	4	5	6	
Frequency	15	22	28	35	42	34	24	$N = \sum f = 200$
c.f.	15	37	65	100	142	176	200	
$fx$	0	22	56	105	168	170	144	$\sum fx = 665$

$$\bar{x} = \frac{\sum fx}{N} = \frac{665}{200} = 3.325$$

$$Md = \frac{N}{2} = \frac{200}{2} = 100^{\text{th}} \text{ items} = 3$$

$M_o$  = No. of calls with maximum frequency = 4

### 3. The length in hours of 100 VCA cable

Length on inch	3.80 - 3.89	3.90 - 3.99	4.00 - 4.09	4.10 - 4.19	4.20 - 4.29	4.30 - 4.39	4.40 - 4.49	5.50 - 5.59
Frequency (f)	3	8	14	19	28	18	10	8
c.f.	3	11	25	44	72	90	100	108

For mode,

$$\text{Model class} = 4.20 - 4.29$$

It is inclusive class, hence adjusted model class = (4.195 - 4.295)

$$f_0 = 19, f_1 = 28, f_2 = 18$$

$$\begin{aligned}
 M_0 &= L + \frac{\Delta_1}{\Delta_1 + \Delta_2} \times h \\
 &= 4.195 + \frac{f_1 - f_0}{f_1 - f_0 + f_1 - f_2} \times h = 4.195 + \frac{28 - 19}{28 - 19 + 28 - 18} \times 0.1 \\
 &= 4.24
 \end{aligned}$$

$$\text{For Median } \frac{N}{2} = \frac{108}{2} = 54$$

$$\text{Median class} = (4.20 - 4.29)$$

$$\text{Adjusted median class} = (4.195 - 4.295)$$

$$M_d = L + \frac{\frac{N}{2} - c.f.}{f} \times h = 4.195 + \frac{45 - 44}{28} \times 0.1 = 4.23$$

**2 CHAPTER - 2 | A Complete Solutions of Statistics I for BSc. CSIT**

4. Compute the mean, median and mode

IQ	Frequency (f)	Mid-value (x)	$d' = \frac{x - 104.5}{10}$	$fd'$	c.f.
50 - 59	2	54.5	-5	-10	2
60 - 69	4	64.5	-4	-16	6
70 - 79	8	74.5	-3	-24	14
80 - 89	15	84.5	-2	-30	29
90 - 99	21	94.5	-1	-21	50
100 - 109	25	104.5	0	0	75
110 - 119	20	114.5	1	20	95
119 - 129	2	124.5	2	4	97
130 - 139	2	134.5	3	6	99
140 - 149	1	144.5	4	4	100
				$\sum fd' = -67$	

$$\bar{x} = A + \frac{\sum fd'}{N} \times h = 104.5 + \frac{(-67)}{100} \times 10 = 97.8$$

$$\frac{N}{2} = \frac{100}{2} = 50,$$

$$Md = L + \frac{\frac{N}{2} - c.f.}{f} \times h = 89.5 + \frac{50 - 25}{21} \times 10 = 99.5$$

Hence  $M_d$  class = (90 - 99)

Adjusted  $M_d$  class = 89.5 - 99.5

Model class = (100 - 109)

Adjusted model class = (99.5 - 109.5)

$$f_0 = 21, f_1 = 25, f_2 = 20, \Delta_1 = f_1 - f_0 = 4, \Delta_2 = f_1 - f_2 = 5$$

$$M_o = L + \frac{\Delta_1}{\Delta_1 + \Delta_2} \times h = 99.5 + \frac{4}{4 + 5} \times 10 = 103.94$$

5. Compute G.M.

$$G.M. = (38 \times 45 \times 50 \times 68 \times 82 \times 40)^{1/6} = 51.68$$

6. Compute G.M.

Weight	50 - 54	55 - 59	60 - 64	65 - 69	70 - 74
Mid-weight	52	57	62	67	72
No. of students (f)	30	25	32	20	13
$\log x$	1.716	1.755	1.792	1.826	1.857
$f \log x$	51.48	43.87	57.34	36.52	24.14

$$\sum f = 120; \sum f \log x = 213.35$$

$$G.M. = \text{Antilog} \left( \frac{\sum f \log x}{N} \right)$$

$$= \text{Antilog} \left( \frac{213.35}{120} \right) = \text{Antilog } 1.777 = 59.96 \text{ kg}$$

7. A boy travels

$$n = 5$$

$$\text{H.M.} = \frac{n}{\sum \frac{1}{x}} = \frac{5}{\frac{1}{40} + \frac{1}{38} + \frac{1}{35} + \frac{1}{36} + \frac{1}{39}} = \frac{5}{0.1333} = 37.509$$

8. The data relating to

Increase in wt	0-5	5-10	10-15	15-20	20-25	25-30	30-35	35-40
No. of animals (f)	3	7	18	35	20	8	5	4
c.f.	3	10	28	63	83	91	96	100

$$\frac{N}{4} = \frac{100}{4} = 25 \quad \therefore Q_1 \text{ class} = (10 - 15)$$

$$Q_1 = L + \frac{\frac{N}{4} - c.f.}{f} \times h = 20 + \frac{25 - 10}{18} \times 5 = 14.166$$

$$\frac{3N}{4} = 3 \times 25 = 75 \quad \therefore Q_3 \text{ class} = (20 - 25)$$

$$Q_3 = L + \frac{\frac{3N}{4} - c.f.}{f} \times h = 20 + \frac{75 - 63}{20} \times 5 = 23$$

$$\frac{9N}{10} = \frac{9 \times 100}{10} = 90 \quad \therefore D_9 \text{ class} = (25 - 30)$$

$$D_9 = L + \frac{\frac{9N}{10} - c.f.}{f} \times h = 25 + \frac{90 - 83}{8} \times 5 = 29.375$$

$$\frac{15N}{100} = \frac{15 \times 100}{100} = 15 \quad \therefore P_{15} \text{ class} = (10 - 15)$$

$$P_{15} = L + \frac{\frac{15N}{100} - c.f.}{f} \times h = 10 + \frac{15 - 10}{18} \times 5 = 11.388$$

$$\frac{90N}{100} = \frac{90 \times 100}{100} = 90 \quad \therefore P_{90} \text{ class} = (25 - 30)$$

$$P_{90} = L + \frac{\frac{90N}{100} - c.f.}{f} \times h = 25 + \frac{90 - 83}{8} \times 5 = 29.375$$

9. Random sample

Size of pen drive	No. of user (f)
4 - 5	10
5 - 6	35
6 - 7	70
7 - 8	35
8 - 9	12
9 - 10	6

**CHAPTER - 2 | A Complete Solutions of Statistics I for BSc. CSIT**  
Measure of central tendency in Mode.

Here modal class is (6 - 7)

$$L = 6, f_0 = 35, f_1 = 70, f_2 = 35, \Delta_{12} = f_1 - f_0 = 70 - 35 = 35 \\ \Delta_2 = f_1 - f_2 = 70 - 35 = 35$$

$$M_O = L + \frac{\Delta_1}{\Delta_1 + \Delta_2} \times h = 6 + \frac{35}{35 + 35} \times 1 = 6.5$$

✓ 10. A software

$$x_1 = 50, x_2 = 35, x_3 = 40, x_4 = 25, \bar{x}_1 = 30,000; \bar{x}_2 = 35,000, \bar{x}_3 = 40,000, \\ \bar{x}_4 = 50,000$$

$$\bar{x}_{1234} = \frac{n_1 \bar{x}_1 + n_2 \bar{x}_2 + n_3 \bar{x}_3 + n_4 \bar{x}_4}{n_1 + n_2 + n_3 + n_4} \\ = \frac{50 \times 30000 + 35 \times 35000 + 40 \times 40000 + 25 \times 50000}{50 + 35 + 40 + 25} = 37166.67$$

11. The percentage age

Age	0-4	5-9	10-14	15-19	20-24	25-29	30-34
Male pop <sup>n</sup> (f)	11.8	12.9	12.5	11.2	10.7	8.9	7.2
c.f.	11.8	24.7	37.2	48.4	59.1	68.0	75.2
	35-39	40-44	45-49	50-54	55-59	60 and above	
	6.2	4.7	4.0	2.9	2.3	4.7	
	81.4	86.1	90.1	93.0	95.3	100	

$$\frac{N}{4} = \frac{100}{4} = 25; \text{ Hence } Q_1 \text{ class} = (10 - 14)$$

Adjusted  $Q_1$  class = (9.5 - 14.5)

$$Q_1 = L + \frac{\frac{N}{4} - c.f.}{f} \times h = 9.5 + \frac{25 - 24.7}{12.5} \times 5 = 9.62$$

$$\frac{3N}{4} = 3 \times 25 = 75; \text{ Hence } Q_3 \text{ class} = (30 - 34)$$

Adjusted  $Q_3$  class = (29.5 - 34.5)

$$Q_3 = L + \frac{\frac{3N}{4} - c.f.}{f} \times h = 29.5 + \frac{75 - 68}{7.2} \times 5 = 34.36$$

$$\frac{8N}{10} = \frac{8 \times 100}{10} = 80$$

$D_{80}$  Class = (35 - 39)

Adj.  $D_{80}$  class = (34.5 - 39.5)

$$D_{80} = L + \frac{\frac{8N}{10} - c.f.}{f} \times h = 34.5 + \frac{80 - 75.2}{6.2} \times 5 = 38.37$$

$$\frac{70N}{100} = \frac{70 \times 100}{100} = 70$$

$P_{70}$  class (30 - 34) $\text{Adj. } P_{70} \text{ class} = (29.5 - 34.5)$ 

$$P_{70} = L + \frac{\frac{70 N}{100} - c.f.}{f} \times h = 29.5 + \frac{70 - 68}{7.2} \times 5 = 30.88$$

12. The percentage of

$$\begin{aligned} GM &= (x_1 \times x_2 \times x_3 \times x_4 \times x_5 \times x_6 \times x_7)^{1/7} \\ &= (49.19 \times 49.25 \times 50.34 \times 45.122 \times 49.87 \times 58.27 \times 83.58)^{1/7} \\ &= 48.717 \end{aligned}$$

13.

Marks	No. of students (f)	c.f.
10 - 20	15	15
20 - 40	20	35
40 - 70	30	65
70 - 90	20	85
90 - 100	15	100
	$N = \sum f = 100$	

Highest marks of weakest 30% students is  $P_{30}$ 

$$P_{30} = \frac{30 N}{100}^{\text{th}} \text{ item} = \frac{30 \times 100}{100} = 30$$

 $P_{30}$  class = (20 - 40)

$$P_{30} = L + \frac{\frac{30 N}{100} - c.f.}{f} \times h = 20 + \frac{30 - 15}{20} \times 20 = 35$$

Lowest marks of top 40% student is  $P_{60}$ 

$$P_{60} = \frac{60 N}{100} \text{ item} = \frac{60 \times 100}{100} = 60$$

 $P_{60}$  class = (40 - 70)

$$P_{60} = L + \frac{\frac{60 N}{100} - c.f.}{f} \times h = 40 + \frac{60 - 35}{30} \times 30 = 65$$

Lowest marks of top 20% students  $P_{80}$ 

$$P_{80} = \frac{80 N}{100} \text{ item} = \frac{80 \times 100}{100} = 80$$

 $P_{80}$  class = (70 - 90)

$$P_{80} = L + \frac{\frac{80 N}{100} - c.f.}{f} \times h = 70 + \frac{80 - 65}{20} \times 20 = 85$$

Limit of middle 50% student is  $P_{25}$  to  $P_{75}$ .

$$P_{25} = \frac{25 N}{100}^{\text{th}} \text{ item} = \frac{25 \times 100}{100} = 25$$

 $P_{25}$  class = (20 - 40)

$$P_{25} = L + \frac{\frac{25 N}{100} - c.f.}{f} \times h = 20 + \frac{25 - 15}{20} \times 20 = 30$$

$$P_{75} = L + \frac{\frac{75 N}{100} - c.f.}{f} \times h = 70 + \frac{75 - 65}{20} \times 20 = 80$$

$$\text{Range} = P_{75} - P_{25} = 80 - 30 = 50$$

14. Compute the quartile deviation

Size of screen	No. of laptop (f)	c.f.
9.5	1	1
10.0	8	9
10.5	20	29
11.0	30	59
11.5	50	109
12.0	95	204
12.5	110	314
13.0	150	464
13.5	200	664
14.0	250	914
14.5	280	1194
15.0	245	1439
15.5	80	1519
16.0	40	1559
16.5	35	1594
17.0	5	1599
$N = \sum f = 1599$		

$$Q_1 = \left( \frac{N+1}{4} \right)^{\text{th}} \text{ item} = \frac{1599+1}{4} = \frac{16000}{4} = 400 \quad \therefore Q_1 = 13,$$

$$Q_3 = \left( \frac{3(N+1)}{4} \right)^{\text{th}} \text{ item} = 3 \times 400 = 1200 \quad \therefore Q_3 = 15$$

$$\text{Quartile deviation} = \frac{Q_3 - Q_1}{2} = \frac{15 - 13}{2} = 1$$

✓ 15. The following frequency

Weight	4-5	5-6	6-7	7-8	8-9	9-10	10-11	11-12
Freq. (f)	20	24	35	48	32	24	8	2
c.f.	20	44	79	127	159	183	191	193

$$\frac{N}{4} = \frac{193}{4} = 48.25 \quad \therefore Q_1 \text{ class} = (6 - 7)$$

$$Q_1 = L + \frac{\frac{N}{4} - c.f.}{f} \times h = 6 + \frac{48.25 - 44}{35} \times 1 = 6.12$$

$$\frac{2N}{4} = 2 \times 48.25 = 96.5 \quad \therefore Q_2 \text{ class} = (7 - 8)$$

$$Q_2 = L + \frac{\frac{2N}{4} - c.f.}{f} \times h = 7 + \frac{96.5 - 79}{48} \times 1 = 7.36$$

$$\frac{3N}{4} = 3 \times 48.25 = 144.75 \quad \therefore Q_3 \text{ class} = (8 - 9)$$

$$Q_3 = L + \frac{\frac{3N}{4} - c.f.}{f} \times h = 8 + \frac{144.75 - 127}{32} \times 1 = 8.55$$

$$QD = (Q_3 - Q_1)/2 = (8.55 - 6.12)/2 = 1.215$$

16. The scores obtained

$$L = 65, S = 35$$

$$\text{Range} = L - S = 65 - 35 = 30$$

Score (x)	$x^2$
55	3025
35	1225
60	3600
55	3025
55	3025
65	4225
40	1600
45	2025
35	1225
42	1764
$\Sigma x = 487$	$\Sigma x^2 = 24739$

$$S.D. = \sqrt{\frac{\Sigma x^2}{n} - \left(\frac{\Sigma x}{n}\right)^2} = \sqrt{\frac{24739}{10} - \left(\frac{487}{10}\right)^2} = 10.109$$

17. Arrange data in ascending order

$$32, 34, 34, 35, 35, 35, 36$$

$$\text{Range} = L - S = 36 - 32 = 4$$

$$n = 7$$

$$Q_1 = \frac{n+1}{4}^{\text{th}} \text{ item} = \frac{7+1}{4} = 2^{\text{nd}} \text{ item} = 34$$

$$Q_3 = \frac{3(n+1)}{4}^{\text{th}} \text{ item} = 3 \times 2 = 6^{\text{th}} \text{ item} = 35$$

$$Q.D. = \frac{Q_3 - Q_1}{2} = \frac{35 - 34}{1} = 0.5$$

18. The time in minutes

x	10	12	11	15	18	20	24	23	26	16
$ x - \bar{x} $	7.5	5.5	6.5	2.5	0.5	2.5	6.5	5.5	8.5	1.5
$x^2$	100	144	121	225	324	400	576	529	676	256

$$\sum x = 175, \sum |x - \bar{x}| = 47, \sum x^2 = 3351$$

$$\bar{x} = \frac{\sum x}{n} = \frac{175}{10} = 17.5$$

$$\text{M.D. from } \bar{x} = \frac{\sum |x - \bar{x}|}{n} = \frac{47}{10} = 4.7$$

$$\text{S.D.} = \sqrt{\frac{\sum x^2}{n} - \left(\frac{\sum x}{n}\right)^2} = \sqrt{\frac{3351}{10} - (17.5)^2} = 5.37$$

19. To obtain information

Customer service time	$fd'^2$	No. of customers (f)	Mid-value (x)	$d' = \frac{x - 17.5}{5}$	$fd'$
0 - 5	18	2	2.5	-3	-6
5 - 10	32	3	7.5	-2	-16
10 - 15	26	26	12.5	-1	-26
15 - 20	0	30	17.5	0	0
20 - 25	28	28	22.5	1	28
25 - 30	24	6	27.5	2	12
	$\sum fd'^2 = 128$	$N = \sum f = 100$			$\sum fd' = -8$

$$\sigma = \sqrt{\frac{\sum fd'^2}{N} - \left(\frac{\sum fd'}{N}\right)^2} \times h = \sqrt{\frac{128}{100} - \left(\frac{-8}{100}\right)^2} \times 5 = 5.64$$

20. The frequency distribution

Time in sec	No. of customers (f)	Mid-value (x)	$d' = \frac{x - 17}{5}$	$fd'$	$fd'^2$
0 - 4	2	2	-3	-6	18
5 - 9	20	7	-2	-40	80
10 - 14	35	12	-1	-35	35
15 - 19	40	17	0	0	0
20 - 24	28	22	1	28	28
25 - 29	32	27	2	64	128
30 - 34	8	32	3	24	72
35 - 39	5	37	4	20	80
	170			55	441

$$N = \sum f = 170, \sum fd' = 55, \sum fd'^2 = 441$$

$$\text{S.D. } (\sigma) = \sqrt{\frac{\sum fd'^2}{N} - \left(\frac{\sum fd'}{N}\right)^2} \times h$$

$$= \sqrt{\frac{441}{170} - \left(\frac{55}{170}\right)^2} \times 5 = \sqrt{2.594 - 0.104} \times 5 = 7.88$$

21. The average marks secured by

$$\bar{x}_A = 78, \bar{x}_B = 80, \sigma_A^2 = 100, \sigma_B^2 = 81, x_A = 100, x_B = 150$$

$$\bar{x}_{AB} = \frac{n_A \bar{x}_A + n_B \bar{x}_B}{n_A + n_B} = \frac{100 \times 78 + 150 \times 80}{100 + 150} = 79.2$$

$$d_A = \bar{x}_A - \bar{x}_{AB} = 78 - 79.2 = -1.2$$

$$d_B = \bar{x}_B - \bar{x}_{AB} = 80 - 79.2 = 0.8$$

$$\begin{aligned}\sigma_{AB} &= \sqrt{\frac{n_A (\sigma_A^2 + d_B^2) + n_B (\sigma_B^2 + d_B^2)}{n_A + n_B}} \\ &= \sqrt{\frac{100(100 + 1.44) + 150(81 + 0.64)}{100 + 150}} = 9.46\end{aligned}$$

$$\sigma_{AB}^2 = 89.5$$

22. The number of runs

Group A ( $x_A$ )	Group B ( $x_B$ )	$x_A^2$	$x_B^2$
10	120	100	14400
25	15	625	225
85	30	7225	900
72	35	5184	1225
115	42	13225	1764
80	65	6400	4225
52	80	2704	6400
45	34	2025	1156
30	25	900	625
10	15	100	225
$\sum x_A = 524$	$\sum x_B = 461$	$\sum x_A^2 = 38488$	$\sum x_B^2 = 29929$

$$\bar{x}_A = \frac{\sum x_A}{n} = \frac{524}{10} = 52.4$$

$$\sigma_A = \sqrt{\frac{\sum x_A}{n} - \left(\frac{\sum x_A}{n}\right)^2} = \sqrt{\frac{38488}{10} - \left(\frac{524}{10}\right)^2} = 33.212$$

$$\bar{x}_B = \frac{\sum x_B}{n} = \frac{461}{10} = 46.1$$

$$\sigma_B = \sqrt{\frac{\sum x_B}{n} - \left(\frac{\sum x_B}{n}\right)^2} = \sqrt{\frac{29920}{10} - \left(\frac{461}{10}\right)^2} = 29.441$$

$$C.V._A = \frac{\sigma_A}{\bar{x}_A} \times 100\% = \frac{33.212}{52.4} \times 100\% = 63.38\%$$

$$C.V._B = \frac{\sigma_B}{\bar{x}_B} \times 100\% = \frac{29.441}{46.1} \times 100\% = 63.86\%$$

$C.V._A < C.V._B$ ; Hence group A is more consistent.

**10 CHAPTER - 2 | A Complete Solutions of Statistics I for BSc. CSIT**

23. The following data

Students	Section A ( $x_A$ )	Section B ( $x_B$ )	$x_A^2$	$x_B^2$
1	9	10	81	100
2	8	8	64	64
3	10	6	100	36
4	6	8	36	64
5	7	9	49	81
6	8	8	64	64
7	5	7	25	49
8	6	8	36	64
9	7	5	49	25
10	8	8	64	64
	$\sum x_A = 74$	$\sum x_B = 77$	$\sum x_A^2 = 568$	$\sum x_B^2 = 611$

$$\bar{x}_A = \frac{\sum x_A}{n} = \frac{74}{10} = 7.4$$

$$\sigma_A = \sqrt{\frac{\sum x_A^2}{n} - \left(\frac{\sum x_A}{n}\right)^2} = \sqrt{\frac{568}{10} - (7.4)^2} = 1.42$$

$$\bar{x}_B = \frac{\sum x_B}{n} = \frac{77}{10} = 7.7$$

$$\sigma_B = \sqrt{\frac{\sum x_B^2}{n} - \left(\frac{\sum x_B}{n}\right)^2} = \sqrt{\frac{611}{10} - (7.7)^2} = 1.34$$

$$C.V._A = \frac{\sigma_A}{\bar{x}_A} \times 100\% = \frac{1.42}{7.4} \times 100\% = 19.18\%$$

$$C.V._B = \frac{\sigma_B}{\bar{x}_B} \times 100\% = \frac{1.34}{7.7} \times 100\% = 17.4\%$$

$C.V._B < C.V._A$ . Hence, group B is more consistent.

24. The following data

Time for male ( $x_A$ )	Time for female ( $x_B$ )	$x_A^2$	$x_B^2$
5.70	7.52	32.49	56.55
6.80	8.20	46.24	67.24
7.25	8.32	52.56	69.22
8.20	6.90	67.24	47.61
8.10	6.80	65.61	46.24
7.20	8.30	51.84	68.89
6.88	7.45	47.33	55.50
7.20	9.00	51.84	81.00
7.35	10.50	54.02	110.25
7.45	7.20	55.50	51.84

6.90	10.20	47.61	104.04
7.22	8.26	52.12	68.22
6.85	8.50	46.92	72.25
6.40	8.32	40.96	69.22
6.20	10.00	38.44	100
$\sum x_A = 105.7$	$\sum x_B = 125.47$	$\sum x_A^2 = 750.72$	$\sum x_B^2 = 1068.07$

$$\bar{x}_A = \frac{\sum x_A}{n} = \frac{105.7}{15} = 7.046$$

$$\sigma_A = \sqrt{\frac{\sum x_A^2}{n} - \left(\frac{\sum x_A}{n}\right)^2} = \sqrt{\frac{750.72}{15} - \left(\frac{105.7}{15}\right)^2} = 0.673$$

$$\bar{x}_B = \frac{\sum x_B}{n} = \frac{125.47}{15} = 8.364$$

$$\sigma_B = \sqrt{\frac{\sum x_B^2}{n} - \left(\frac{\sum x_B}{n}\right)^2} = \sqrt{\frac{1068.07}{15} - \left(\frac{125.47}{15}\right)^2} = 1.117$$

$$C.V._A = \frac{\sigma_A}{\bar{x}_A} \times 100\% = \frac{0.673}{7.046} \times 100\% = 8.98\%$$

$$C.V._B = \frac{\sigma_B}{\bar{x}_B} \times 100\% = \frac{1.117}{8.364} \times 100\% = 13.35\%$$

$CV_A < CV_B$ . Hence male students is more consistent.

25. Solution.

$$\bar{x} = \frac{\sum x}{n} = \frac{552}{10} = 55.20$$

$$\sigma = \sqrt{\frac{\sum x^2}{n} - \left(\frac{\sum x}{n}\right)^2} = \sqrt{\frac{31700}{10} - \left(\frac{552}{10}\right)^2} = \sqrt{122.96} = 11.08$$

$$S_{KP} = \frac{\bar{x} - M_o}{\sigma} = \frac{55.2 - 46.2}{11.08} = 0.81$$

26. Solution.

$$n = 20, \sum x = 300, \sum x^2 = 5000, Md = 15, \bar{x} = \frac{300}{20} = 15,$$

$$\sigma = \sqrt{\frac{\sum x^2}{n} - \left(\frac{\sum x}{n}\right)^2} = \sqrt{\frac{5000}{20} - (15)^2} = 5$$

$$S_{KP} = 3 \frac{(\bar{x} - Md)}{\sigma} = 3 \times \frac{15 - 15}{5} = 0$$

$$CV = \frac{\sigma}{\bar{x}} \times 100 = \frac{5}{15} \times 100 = 33.3\%$$

27. Solution.

$$\bar{x} = 65, Md = 70, S_{KP} = -0.6, M_o = ?, C.V. = ?$$

$$M_o = 3Md - 2\bar{x} = 3 \times 70 - 2 \times 65 = 80$$

$$S_{KP} = \frac{\bar{x} - M_O}{\sigma}$$

$$\text{or } -0.6 = \frac{65 - 80}{\sigma}$$

$$\text{or } -0.6 = \frac{-15}{\sigma}$$

$$\text{or } \sigma = \frac{15}{0.6} = \frac{150}{6} = 25$$

$$CV = \frac{\sigma}{\bar{x}} \times 100\% = \frac{25}{65} \times 100\% = 38.40\%$$

28. Solution.

Given,  $\bar{x} = 200$ , C.V. = 8%,  $S_{KP} = 0.3$ ,  $M_O = ?$ ,  $M_d = ?$

$$C.V. = \frac{\sigma}{\bar{x}} \times 100\%$$

$$\text{or } 8\% = \frac{\sigma}{200} \times 100\%$$

$$\text{or } \sigma = \frac{200 \times 8}{100} = 16$$

$$S_{KP} = \frac{\bar{x} - M_O}{\sigma}$$

$$\text{or } 0.3 = \frac{200 - M_O}{16}$$

$$\text{or } 4.8 = 200 - M_O$$

$$\text{or } M_O = 200 - 4.8 = 195.20$$

$$M_O = 3M_d - 2\bar{x}$$

$$\text{or } 195.2 = 3M_d - 2 \times 200$$

$$\text{or } 3M_d = 400 + 195.20 = 595.2$$

$$\text{or } M_d = \frac{595.2}{3} = 198.40$$

29. Solution.

Size (x)	Frequency (f)	$fx$	$fx^2$
6	7	42	252
9	12	108	972
12	19	228	2736
15	10	150	2250
18	2	36	648
	$N = 50$	$\Sigma fx = 564$	$\Sigma fx^2 = 6858$

$$\bar{x} = \frac{\Sigma fx}{N} = \frac{564}{50} = 11.28$$

$$\sigma = \sqrt{\frac{\sum fx^2}{N} - \left(\frac{\sum fx}{N}\right)^2} = \sqrt{\frac{6858}{50} - \left(\frac{564}{50}\right)^2} = 3.149$$

$$M_O = 12$$

$$S_{KP} = \frac{\bar{x} - M_O}{\sigma} = \frac{11.28 - 12}{3.149} = -0.23$$

30. Solution.

Mid Value (x)	No. of workers (f)	$fx$	$fx^2$
150	80		
250	105		
350	120		
450	165		
550	100		
650	90		
750	60		
850	40		
	$N = 50$	$\sum fx = 347000$	$\sum fx^2 = 187400000$

$$\bar{x} = \frac{\sum fx}{N} = \frac{347000}{760} = 456.58$$

$$\sigma = \sqrt{\frac{\sum fx^2}{N} - \left(\frac{\sum fx}{N}\right)^2} = \sqrt{\frac{187400000}{760} - \left(\frac{347000}{760}\right)^2} = 195.23$$

$$\text{Model class} = (400 - 500)$$

$$f_0 = 120, f_1 = 165, f_2 = 100$$

$$\Delta_1 = f_1 - f_0 = 165 - 120 = 45 \quad \Delta_2 = f_1 - f_2 = 165 - 100 = 65$$

$$M_O = L + \frac{\Delta_1}{\Delta_1 + \Delta_2} \times h = 400 + \frac{45}{45 + 65} \times 100 = 400 + \frac{4500}{110} = 440.9$$

$$S_{KP} = \frac{\bar{x} - M_O}{\sigma} = \frac{456.57 - 440.90}{195.23} = 0.08$$

31.  $S_{KB} = 0.6$

$$Q_3 + Q_1 = 100 \quad \dots (1)$$

$$Md = 38$$

$$Q_3 = ?$$

$$S_{KB} = \frac{Q_3 + Q_1 - 2 Md}{Q_3 - Q_1}$$

$$\text{or } 0.6 = \frac{100 - 2 \times 38}{Q_3 - Q_1}$$

$$\text{or } Q_3 - Q_1 = \frac{100 - 76}{0.6}$$

$$\text{or } Q_3 - Q_1 = 40 \quad \dots (2)$$

$$\text{Add (1) and (2)}$$

$$2Q_3 = 140$$

$$\therefore Q_3 = 70$$

14 CHAPTER - 2 | A Complete Solutions of Statistics I for BSc. CSIT

32. Solution.

$$S_{KB} = -0.8, Q_1 = 44.10, Q_3 = 56.60, Md = ?$$

$$S_{KB} = \frac{Q_3 + Q_1 - 2Md}{Q_3 - Q_1}$$

$$\text{or } -0.8 = \frac{56.6 + 44.1 - 2Md}{56.6 - 44.1}$$

$$\text{or } -0.8 = \frac{100.70 - 2Md}{12.5}$$

$$\text{or } -0.8 \times 12.5 = 100.70 - 2Md$$

$$\text{or } 2Md = 10 + 100.70$$

$$\text{or } Md = \frac{110.70}{2} = 55.35$$

33. Solution.

✓

x	f	c.f.
20	3	3
23	6	9
24	10	19
26	12	31
28	11	42
30	9	51
40	4	55
$N = \sum f = 55$		

$$Q_1 = \frac{N+1}{4}^{\text{th}} \text{ item} = \frac{55+1}{4} = 14^{\text{th}} \text{ item} = 24$$

$$Md = \frac{2(N+1)}{4}^{\text{th}} \text{ item} = 2 \times 14 = 28^{\text{th}} \text{ item} = 26$$

$$Q_3 = \frac{3(N+1)}{4}^{\text{th}} \text{ item} = 3 \times 14 = 42^{\text{th}} \text{ item} = 28$$

$$S_{KB} = \frac{Q_3 + Q_1 - 2Md}{Q_3 - Q_1} = \frac{28 + 24 - 26}{28 - 24} = 0$$

34. Solution.

Hourly wage	No. of workers (f)	c.f.
23 - 27	22	22
28 - 32	16	38
33 - 37	9	47
38 - 42	4	51
43 - 47	3	54
48 - 52	1	55
$N = \sum f = 55$		

$$\frac{N}{4} = \frac{55}{4} = 13.75$$

$$Q_1 \text{ class} = (23 - 27)$$

$$\text{Adjusted } Q_1 \text{ class} = 22.5 - 27.5$$

$$Q_1 = L + \frac{\frac{N}{4} - c.f.}{f} \times h = 22.5 + \frac{13.75 - 0}{22} \times 5 = 25.62$$

$$\frac{2N}{4} = 2 \times 13.75 = 27.5$$

Md class = 28 - 32

Adjusted Md class = 27.5 - 32.5

$$Md = L + \frac{\frac{2N}{4} - c.f.}{f} \times h = 27.5 + \frac{27.5 - 22}{16} \times 5 = 27.5 + 1.718 = 29.218$$

$$\frac{3N}{4} = 3 \times 13.75 = 41.25$$

$Q_3$  class = (33 - 37)  $\therefore$  Adjusted class = 32.5 - 37.5

$$Q_3 = L + \frac{\frac{3N}{4} - c.f.}{f} \times h = 32.5 + \frac{41.25 - 38}{9} \times 5 = 34.305$$

$$S_{KB} = \frac{Q_3 + Q_1 - 2Md}{Q_3 - Q_1} = \frac{34.305 + 25.62 - 2 \times 29.218}{34.305 - 25.62} = 0.17$$

35. Solution.

Download speed (Mbps)	Time in min.	c.f.
Below 100	10	10
100 - 150	25	35
150 - 200	145	180
200 - 250	220	400
250 - 300	70	470
300 and above	30	500

$$\frac{N}{4} = \frac{500}{4} = 125 \quad \therefore Q_1 \text{ class} = (150 - 200)$$

$$Q_1 = L + \frac{\frac{N}{4} - c.f.}{f} \times h = 150 + \frac{125 - 35}{145} \times 50 = 181.034$$

$$\frac{2N}{4} = 2 \times 125 = 250 \quad \therefore Md \text{ class} = (200 - 250)$$

$$Md = L + \frac{\frac{2N}{4} - c.f.}{f} \times h = 200 + \frac{250 - 180}{220} \times 50 = 215.90$$

$$\frac{3N}{4} = 3 \times 125 = 375$$

$Q_3$  class = (200 - 250)

$$Q_3 = L + \frac{\frac{3N}{4} - c.f.}{f} \times h = 200 + \frac{375 - 180}{220} \times 50 = 244.32$$

$$S_{KB} = \frac{Q_3 + Q_1 - 2 Md}{Q_3 - Q_1} = \frac{244.32 + 181.03 - 2 \times 215.90}{244.32 - 181.03} = -0.102$$

x	f	c.f.
2	1	1
4	5	6
6	10	16
8	3	19
10	1	20
	$N = \sum f = 20$	

$$\text{For } P_{10}, \frac{10(N+1)}{100} = \frac{10 \times 21}{100} = 2.1 \quad \therefore P_{10} = 4$$

$$\text{For } P_{90}, \frac{90(N+1)}{100} = \frac{90 \times 21}{100} = 18.9 \quad \therefore P_{90} = 8$$

$$\text{For } Q_1, \frac{N+1}{4} = \frac{20+1}{4} = 5.25 \quad \therefore Q_1 = 4$$

$$\text{For } Q_3, \frac{3(N+1)}{4} = 3 \times 5.25 = 15.75 \quad \therefore Q_3 = 6$$

$$K = \frac{Q_3 - Q_1}{2(P_{90} - P_{10})} = \frac{6 - 4}{2(8 - 4)} = \frac{2}{2 \times 4} = 0.25$$

## 37. Solution.

Hourly remuneration	No. of person (f)	c.f.
100 - 110	10	10
110 - 120	14	24
120 - 130	18	42
130 - 140	24	66
140 - 150	16	82
150 - 160	12	94
160 - 170	6	100
	$N = \sum f = 100$	

$$\text{For } P_{10}, \frac{10N}{100} = \frac{10 \times 100}{100} = 10 \quad \therefore P_{10} \text{ class} = 100 - 110$$

$$P_{10} = L + \frac{\frac{10N}{100} - c.f.}{f} \times h = 100 + \frac{10 - 0}{10} \times 10 = 110$$

$$\text{For } P_{90}, \frac{90N}{100} = \frac{90 \times 100}{100} = 90 \quad \therefore P_{90} \text{ class} = 150 - 160$$

$$P_{90} = L + \frac{\frac{90N}{100} - c.f.}{f} \times h = 150 + \frac{90 - 82}{12} \times 10 = 156.66$$

$$\text{For } Q_1, \frac{N}{4} = \frac{100}{4} = 25 \quad \therefore Q_1 \text{ class} = (120 - 130)$$

$$Q_1 = L + \frac{\frac{N}{4} - c.f.}{f} \times h = 120 + \frac{25 - 24}{18} \times 10 = 120.55$$

For  $Q_3$ ,  $\frac{3N}{4} = 3 \times 25 = 75$

$\therefore Q_3$  class = (140 - 150)

$$Q_3 = L + \frac{\frac{3N}{4} - c.f.}{f} \times h = 140 + \frac{75 - 66}{16} \times 10 = 145.625$$

$$K = \frac{145.625 - 120.55}{2(156.66 - 110)} = \frac{25.075}{93.32} = 0.268$$

### 38. Solution.

Class interval	frequency (f)	Mid-value (x)	fx	$(x - \bar{x})$	$f(x - \bar{x})$	$f(x - \bar{x})^2$	$(x - \bar{x})^3$	$f(x - \bar{x})^4$
0 - 10	5	5	25	-20	-100	25	-20	-100
10 - 20	7	15	105	-10	-70	105	-10	-70
20 - 30	9	25	225	0	0	225	0	0
30 - 40	7	35	245	10	70	245	10	70
40 - 50	5	45	225	20	100	225	20	100
	33		825		0	5400	0	1740000

$$\bar{x} = \frac{\sum fx}{N} = \frac{825}{33} = 25$$

$$\mu_1 = \frac{\sum f(x - \bar{x})}{N} = \frac{0}{33} = 0$$

$$\mu_2 = \frac{\sum f(x - \bar{x})^2}{N} = \frac{5400}{33} = 163.63$$

$$\mu_3 = \frac{\sum f(x - \bar{x})^3}{N} = \frac{0}{33} = 0$$

$$\mu_4 = \frac{\sum f(x - \bar{x})^4}{N} = \frac{1740000}{33} = 52727.27$$

### 39. Solution.

x	f	$d = x - 4$	fd	$fd^2$	$fd^3$	$fd^4$
2	1	-2	-2	4	-8	16
3	3	-1	-3	3	-3	3
4	7	0	0	0	0	0
5	2	1	2	2	2	2
6	1	2	2	4	8	16
	$N = 14$		$\sum fd = -1$	$\sum fd^2 = 13$	$\sum fd^3 = -1$	$\sum fd^4 = 37$

$$\mu'_1 = \frac{\sum fd}{N} = \frac{-1}{14} = -0.071$$

$$\mu'_2 = \frac{\sum fd^2}{N} = \frac{13}{14} = 0.928$$

$$\mu'_3 = \frac{\sum fd^3}{N} = \frac{-1}{14} = -0.071$$

$$\mu'_4 = \frac{\sum fd^4}{N} = \frac{37}{14} = 2.64$$

### 40. Solution.

Marks	f	x	$d' = \frac{x - 45}{10}$	$fd'$	$fd'^2$	$fd'^3$
20 - 30	7	25	-2	-14	28	-56
30 - 40	10	35	-1	-10	10	-10
40 - 50	20	45	0	0	0	0
50 - 60	18	55	1	18	18	18
60 - 70	7	65	2	14	28	56
	$N = \sum f = 62$			$\sum fd' = 8$	$\sum fd'^2 = 84$	$\sum fd'^3 = 8$

$$\mu'_1 = h \times \frac{\sum fd}{N} = 10 \times \frac{8}{62} = 1.29 \quad \mu'_2 = h^2 \times \frac{\sum fd^2}{N} = 10^2 \times \frac{84}{62} = 135.483$$

$$\mu'_3 = h^3 \times \frac{\sum fd^3}{N} = 10^3 \times \frac{8}{62} = 129.03$$

$$\mu_1 = 0$$

$$\mu_2 = \mu'_2 - (\mu'_1)^2 = 135.483 - (1.29)^2 = 133.81$$

$$\mu_3 = \mu'_3 - 3\mu'_2 \mu'_1 + 2(\mu'_1)^3 = 129.03 - 3 \times 135.483 \times 1.29 + 2 \times (1.29)^3 = -390.98$$

41. given

$$\text{First arbitrary moment} = \mu'_1 = -2$$

$$\text{Second arbitrary moment} = \mu'_2 = 14$$

$$\text{Third arbitrary moment} = \mu'_3 = -20$$

$$\text{Fourth arbitrary moment} = \mu'_4 = 50$$

Now We have

$$\text{First central moment} = \mu_1 = \mu'_1 - \mu'_1 = 0$$

$$\text{Second central moment} = \mu'_2 = \mu'_2 - \mu'_1{}^2 = 14 - 2^2 = 10$$

$$\text{Third central moment} = \mu'_3 - 3\mu'_2 \mu'_1 + 2(\mu'_1)^3$$

$$= -20 - 3 \times 14 - 2 + 2 \times (-2)^3 = -20 + 84 - 16 = 48$$

$$\text{Fourth central moment} = \mu'_4 = \mu'_4 - 4\mu'_3 \mu'_1 + 6\mu'_2 \mu'_1{}^2 + 3(\mu'_1)^4$$

$$= 50 - 4 \times 200020 \times (-2) + 6 \times 14 \times (-2)^2 - 3 \times (-2)^4$$

$$= 50 - 160 + 336 - 48 = 178$$

42. Solution.

Given

$$\text{First arbitrary moment} = \mu_1 = 0$$

$$\text{Second arbitrary moment} = \mu_2 = 2.8$$

$$\text{Third arbitrary moment} = \mu_3 = -2$$

$$\text{Fourth arbitrary moment} = \mu_4 = 24.5$$

Now

We have

$$\text{Coefficient of skewness } (\gamma_1) = \frac{\mu_3}{\mu_2^{3/2}} = \frac{-2}{(2.8)^{3/2}} = -0.427$$

$\gamma_1 = -0.427 < 0$ , hence the distribution is negatively skewed.

$$\text{Now, coefficient of kurtosis } (\beta_2) = \frac{\mu_4}{\mu_2^2} = \frac{24.5}{(2.8)^2} = 3.125$$

$\beta_2 = 3.125 > 3$ , hence the distribution is leptokurtic.

43. Solution.

$$\text{First arbitrary moment about } x=5 = \mu'_1 = 2$$

$$\text{Second arbitrary moment about } x=5 = \mu'_2 = .20$$

$$\text{Third arbitrary moment about } x=5 = \mu'_3 = 40$$

Now, We have

$$\text{Central Tendency} = \text{AM} = \bar{X} = A + \mu_1' = 5+2 = 7$$

$$\text{Dispersion S.D.} = \sqrt{\mu_2} = \sqrt{\mu_2' - \mu_1'^2} = \sqrt{20 - 2^2} = 4$$

$$\begin{aligned}\text{Third central moment} &= \mu_3 = \mu_3' - 3\mu_2'\mu_1' + 2(\mu_1')^3 \\ &= 40 - 3 \times 20 \times 2 + 2 \times 2^3 = 40 - 120 + 16 = -64\end{aligned}$$

$$\text{Coefficient of skewness } (\gamma_1) = \frac{\mu_3}{\mu_2^{3/2}} = \frac{-64}{(16)^{3/2}} = \frac{-64}{64} = -1$$

$\gamma_1 = -1 < 0$ , hence the distribution is negatively skewed.

#### 44. Solution

Here,  $\sigma = 11$

$$\text{Or, } \mu_2 = \sigma^2 = 11^2 = 121$$

i) For leptokurtic distribution  $\beta_2 > 3$

$$\Rightarrow \frac{\mu_4}{\mu_2^2} > 3$$

$$\Rightarrow \frac{\mu_4}{121^2} > 3$$

$$\Rightarrow \mu_4 > 14641 \times 3$$

$$\Rightarrow \mu_4 > 43923$$

ii) For mesokurtic distribution,  $\beta_2 = 3$       (iii) For platykurtic distribution,  $\beta_2 < 3$

$$\Rightarrow \frac{\mu_4}{\mu_2^2} = 3$$

$$\Rightarrow \frac{\mu_4}{\mu_2^2} < 3$$

$$\Rightarrow \frac{\mu_4}{121^2} = 3$$

$$\Rightarrow \frac{\mu_4}{121^2} < 3$$

$$\Rightarrow \mu_4 = 16641 \times 3$$

$$\Rightarrow \mu_4 < 16641 \times 3$$

$$\Rightarrow \mu_4 = 43923$$

$$\Rightarrow \mu_4 < 43923$$

#### 45. Solution.

x	f	fx	d = x - \bar{x}	fd	fd <sup>2</sup>	fd <sup>3</sup>	fd <sup>4</sup>
0	5	0	-5	-25	125	-625	3125
1	10	10	-4	-40	160	-640	2560
2	30	60	-3	-90	270	-810	2430
3	70	210	-2	-140	280	-560	1120
4	140	560	-1	-140	140	-140	140
5	200	1000	0	0	0	0	0
6	140	840	1	140	140	140	140
7	70	490	2	140	280	560	1120
8	30	240	3	90	270	810	2430
9	10	90	4	40	160	640	2560
10	5	50	5	25	125	625	3125
	N = 710	$\Sigma fx = 35$		$\Sigma fd = 0$	$\Sigma fd^2 = 1950$	$\Sigma fd^3 = 0$	$\Sigma fd^4 = 18750$

$$\text{Mean} = \frac{\sum fm}{N} = \frac{3550}{710} = 5$$

$$\text{First central moment} = \mu_1 = 0$$

$$\text{Second central moment} = \mu_2 = \frac{\sum fd^2}{N} = \frac{1950}{710} = 2.75$$

$$\text{Third central moment} = \mu_3 = \frac{\sum fd^3}{N} = \frac{0}{710} = 0$$

$$\text{Fourth central moment} = \mu_4 = \frac{\sum fd^4}{N} = \frac{18750}{710} = 26.41$$

Now, we have

$$\text{Coefficient of skewness } (\gamma_1) = \frac{\mu_3}{\mu_2^{3/2}} = \frac{0}{(2.75)^{3/2}} = 0$$

$\gamma_1 = 0$ , hence the distribution is symmetrical.

$$\text{Now, coefficient of kurtosis } (\beta_2) = \frac{\mu_4}{\mu_2^2} = \frac{26.41}{(2.75)^2} = 3.49$$

$\beta_2 = 3.49 > 3$ , hence the distribution is leptokurtic.

#### 46. Solution.

For B Sc CSIT Students

Project conducted (x)	No of Students (f)	$\sum fx$	$(X - \bar{X})$	$f(X - \bar{X})$	$f(X - \bar{X})^2$	$f(X - \bar{X})^3$	$f(X - \bar{X})^4$
20	5	100	-3.26	-16.3	53.14	-173.23	564.73
22	7	154	-1.26	-8.82	11.11	-14.00	17.64
23	10	230	-0.26	-2.6	0.68	-0.18	0.05
25	8	200	1.74	13.92	24.22	42.14	73.33
26	5	130	2.74	13.7	37.54	102.85	281.82
	$N = \sum f = 35$	$\sum fx = 818$		$\sum f(X - \bar{X}) = 0$	$\sum f(X - \bar{X})^2 = 126.69$	$\sum f(X - \bar{X})^4 = -42.41$	$\sum f(X - \bar{X})^4 = 937.57$

$$\text{Mean Value } (\bar{X}) = \frac{\sum fx}{N} = \frac{818}{35} = 23.26$$

$$\text{Variance } \mu_2 = \frac{\sum f(X - \bar{X})^2}{N} = \frac{126.69}{35} = 3.62$$

$$\text{Standard Deviation} = \sqrt{\mu^2} = \sqrt{3.62} = 1.9$$

$$\text{Coefficient of Variation} = \frac{SD}{\bar{X}} \times 100 = \frac{1.9}{23.26} \times 100 = 8.18$$

$$\text{Third Central Moment } \mu_3 = \frac{\sum f(X - \bar{X})^3}{N} = \frac{-42.41}{35} = -1.21$$

$$\text{Fourth Central Moment } \mu_4 = \frac{\sum f(X - \bar{X})^4}{N} = \frac{937.57}{35} = 26.79$$

Now, We have

$$\text{Central Tendency} = \text{AM} = \bar{X} = A + \mu_1' = 5+2 = 7$$

$$\text{Dispersion S.D.} = \sqrt{\mu_2} = \sqrt{\mu_2' - \mu_1'^2} = \sqrt{20 - 2^2} = 4$$

$$\begin{aligned}\text{Third central moment} &= \mu_3 = \mu_3' - 3\mu_2' \mu_1' + 2(\mu_1')^3 \\ &= 40 - 3 \times 20 \times 2 + 2 \times 2^3 = 40 - 120 + 16 = -64\end{aligned}$$

$$\text{Coefficient of skewness } (\gamma_1) = \frac{\mu_3}{\mu_2^{3/2}} = \frac{-64}{(16)^{3/2}} = \frac{-64}{64} = -1$$

$\gamma_1 = -1 < 0$ , hence the distribution is negatively skewed.

#### 44. Solution

Here,  $\sigma = 11$

$$\text{Or, } \mu_2 = \sigma^2 = 11^2 = 121$$

i) For leptokurtic distribution  $\beta_2 > 3$

$$\Rightarrow \frac{\mu_4}{\mu_2^2} > 3$$

$$\Rightarrow \frac{\mu_4}{121^2} > 3$$

$$\Rightarrow \mu_4 > 14641 \times 3$$

$$\Rightarrow \mu_4 > 43923$$

ii) For mesokurtic distribution,  $\beta_2 = 3$     (iii) For platykurtic distribution,  $\beta_2 < 3$

$$\Rightarrow \frac{\mu_4}{\mu_2^2} = 3 \qquad \qquad \qquad \Rightarrow \frac{\mu_4}{\mu_2^2} < 3$$

$$\Rightarrow \frac{\mu_4}{121^2} = 3 \qquad \qquad \qquad \Rightarrow \frac{\mu_4}{121^2} < 3$$

$$\Rightarrow \mu_4 = 16641 \times 3 \qquad \qquad \qquad \Rightarrow \mu_4 < 16641 \times 3$$

$$\Rightarrow \mu_4 = 43923 \qquad \qquad \qquad \Rightarrow \mu_4 < 43923$$

#### 45. Solution.

x	f	fx	d=x-\bar{x}	fd	fd <sup>2</sup>	fd <sup>3</sup>	fd <sup>4</sup>
0	5	0	-5	-25	125	-625	3125
1	10	10	-4	-40	160	-640	2560
2	30	60	-3	-90	270	-810	2430
3	70	210	-2	-140	280	-560	1120
4	140	560	-1	-140	140	-140	140
5	200	1000	0	0	0	0	0
6	140	840	1	140	140	140	140
7	70	490	2	140	280	560	1120
8	30	240	3	90	270	810	2430
9	10	90	4	40	160	640	2560
10	5	50	5	25	125	625	3125
	N = 710	$\Sigma fx = 3550$		$\Sigma fd = 0$	$\Sigma fd^2 = 1950$	$\Sigma fd^3 = 0$	$\Sigma fd^4 = 18750$

**20 CHAPTER - 2 | A Complete Solutions of Statistics I for BSc. CSIT**

$$\text{Mean} = \frac{\sum f_m}{N} = \frac{3550}{710} = 5$$

$$\text{First central moment} = \mu_1 = 0$$

$$\text{Second central moment} = \mu_2 = \frac{\sum f d^2}{N} = \frac{1950}{710} = 2.75$$

$$\text{Third central moment} = \mu_3 = \frac{\sum f d^3}{N} = \frac{0}{710} = 0$$

$$\text{Fourth central moment} = \mu_4 = \frac{\sum f d^4}{N} = \frac{18750}{710} = 26.41$$

Now, we have

$$\text{Coefficient of skewness } (\gamma_1) = \frac{\mu_3}{\mu_2^{3/2}} = \frac{0}{(2.75)^{3/2}} = 0$$

$\gamma_1 = 0$ , hence the distribution is symmetrical.

$$\text{Now, coefficient of kurtosis } (\beta_2) = \frac{\mu_4}{\mu_2^2} = \frac{26.41}{(2.75)^2} = 3.49$$

$\beta_2 = 3.49 > 3$ , hence the distribution is leptokurtic.

#### 46. Solution.

For B Sc CSIT Students

Project conducted (x)	No of Students (f)	fX	(X- $\bar{X}$ )	f(X- $\bar{X}$ )	f(X- $\bar{X}$ ) <sup>2</sup>	f(X- $\bar{X}$ ) <sup>3</sup>	f(X- $\bar{X}$ ) <sup>4</sup>
20	5	100	-3.26	-16.3	53.14	-173.23	564.73
22	7	154	-1.26	-8.82	11.11	-14.00	17.64
23	10	230	-0.26	-2.6	0.68	-0.18	0.05
25	8	200	1.74	13.92	24.22	42.14	73.33
26	5	130	2.74	13.7	37.54	102.85	281.82
	N = $\sum f = 35$	$\sum fX = 818$		$\sum f(X - \bar{X}) = 0$	$\sum f(X - \bar{X})^2 = 126.69$	$\sum f(X - \bar{X})^4 = -42.41$	$\sum f(X - \bar{X})^4 = 937.57$

$$\text{Mean Value } (\bar{X}) = \frac{\sum f x}{N} = \frac{818}{35} = 23.26$$

$$\text{Variance} = \mu_2 = \frac{\sum f (X - \bar{X})^2}{N} = \frac{126.69}{35} = 3.62$$

$$\text{Standard Deviation} = \sqrt{\mu^2} = \sqrt{3.62} = 1.9$$

$$\text{Coefficient of Variation} = \frac{SD}{X} \times 100 = \frac{1.9}{23.26} \times 100 = 8.18$$

$$\text{Third Central Moment } \mu_3 = \frac{\sum f (X - \bar{X})^3}{N} = \frac{-42.41}{35} = -1.21$$

$$\text{Fourth Central Moment } \mu_4 = \frac{\sum f (X - \bar{X})^4}{N} = \frac{937.57}{35} = 26.79$$

$$P_{90} = 90N/100^{\text{th}} \text{ item} = 90 \times \frac{60}{100} = 54$$

Cf just greater than  $90N/100$  is 56 for which corresponding class is (118 - 122), hence  $P_{75}$  class is (117.5 - 122.5)

$$L = 117.5, h = 5 f = 6, cf = 50$$

$$\frac{90 N}{100} - c.f. \\ P_{90} = L + \frac{100 - c.f.}{f} \times h = 117.5 + \frac{54 - 50}{6} \times 5 = 120.83$$

$$\text{Percentile coefficient of kurtosis (K)} = \frac{P_{75} - P_{25}}{2(P_{90} - P_{10})} = \frac{115.71 - 105.83}{2(120.83 - 101.50)} = 0.255$$

$K = 0.255 < 0.263$ , hence the distribution is leptokurtic.

#### 48. Solution.

Arranging the data in ascending order of magnitude

0	0	1	1	1	2	2	2	2	3	3	3	3	3
3	4	4	4	4	4	5	5	6					

There are a total of twenty five points in the data set. The median is thus the 13<sup>th</sup> data values or:

$$\text{Here size of median} = \left(\frac{N+1}{2}\right)^{\text{th}} \text{ item} = \left(\frac{26}{2}\right)^{\text{th}} \text{ item} = 13^{\text{th}} \text{ item}$$

Or, Median = 3

$$\text{Similarly, size of } Q_1 = \left(\frac{N+1}{2}\right)^{\text{th}} \text{ item} = \left(\frac{26}{2}\right)^{\text{th}} \text{ item} = 6.5^{\text{th}} \text{ item}$$


$$Q_1 = 6^{\text{th}} + (7^{\text{th}} - 6^{\text{th}} \text{ item}) \times 0.5 = 2 + (2-2) \times 0.5 = 2$$

$$\text{And, size of } Q_3 = \left(3 \frac{N+1}{4}\right)^{\text{th}} = \left(\frac{78}{4}\right)^{\text{th}} = 19.5^{\text{th}} \text{ item}$$

$$Q_3 = 19^{\text{th}} + (20^{\text{th}} - 19^{\text{th}} \text{ item}) \times 0.5 = 4 + (4-4) \times 0.5 = 4$$

We assemble all of the above results together, and report that the five number summaries for the above set of data is 0, 2, 3, 4, 6.

Box and Whisker of Drive D



The data are not skewed, the data are symmetrical

## 49. Solution

Marks, x	No. of students, f	c f
35	3	3
38	5	8
45	10	18
60	8	26
72	4	30
80	2	32
85	1	33

Here size of median =  $\left(\frac{N+1}{2}\right)^{\text{th}}$  item =  $\left(\frac{34}{2}\right)^{\text{th}}$  item = 17<sup>th</sup> item

The cf just greater than 17 is 18 and the corresponding value is 45  
Therefore, Median = 45

Similarly, size of  $Q_1 = \left(3\frac{N+1}{4}\right)^{\text{th}} = \left(\frac{34}{4}\right)^{\text{th}} = 8.5^{\text{th}}$  item


$$Q_1 = 8^{\text{th}} + (9^{\text{th}} - 8^{\text{th}}) \times 0.5 = 38 + (45 - 38) \times 0.5 = 41.5$$

$$\text{And, size of } Q_3 = \left(3\frac{N+1}{4}\right)^{\text{th}} = \left(\frac{102}{4}\right)^{\text{th}} = 25.5^{\text{th}} \text{ item}$$

The cf just greater than 25.5 is 26 and the corresponding value is 60  
 $Q_3 = 60$

We assemble all of the above results together, and report that the five number summaries for the above set of data is 35, 41.5, 45, 60, 85.

Box and Whisker Plot



From the above figure we can say that the distribution is positively skewed.

50. Solution.

vehicles	No of days ( $f$ )	cf
0	10	3
10	20	14
20	30	53
30	40	20
40	50	10
		100

$$\text{Size of } M_d = \frac{N}{2} = \frac{100}{2} = 50^{\text{th}} \text{ item}$$

The c.f. just greater than 50 is 70. Hence the corresponding class 20 - 30 is the median class.

$$M_d = L + \frac{\frac{N}{2} - \text{c.f.}}{f} \times h = 20 + \frac{50 - 17}{53} \times 10 = 20 + 6.23 = 26.23$$

$$\text{Size of } Q_1 = \frac{N}{4} = \frac{100}{4} = 25$$


The c.f. just greater than 25 is 70. Hence,  $Q_1$  lies in class 20 - 30

$$Q_1 = L + \frac{\frac{N}{4} - \text{c.f.}}{f} \times h = 20 + \frac{25 - 17}{53} \times 10 = 20 + 1.51 = 21.51$$

$$\text{Size of } Q_3 = \frac{3N}{4} = \frac{3 \times 100}{4} = 75$$

The c.f. just greater than 75 is 90. Hence,  $Q_3$  lies in class 30-40

$$Q_3 = L + \frac{\frac{3N}{4} - \text{c.f.}}{f} \times h = 30 + \frac{75 - 70}{20} \times 10 = 30 + 2.5 = 32.5$$



The five number summary is  $\{0, 21.51, 26.23, 32.5, 50\}$ , since box and whisker of right side is larger than left so the distribution is positively skewed.

51. Solution. Stem leaf display of above data are as follows

First week	stem	Second week
	2	8
	3	9 9
7 7 2 1	4	0 2 3 4 5 6 6
8 4 4 3 1	5	0 1 2 3 5 6
4 3 2 0 0	6	0 0

52. Solution.

✓ Stem	leaf
11	9
12	0
13	1 3 4 5
14	8 8
15	0 2 4 8 8 9
16	1 2 2 3 7 9
17	1 1 2 2 5 6 7
18	5 6 7 8
19	0 3 4 5 7 8 9
20	2 7
21	6 7 7 9
22	1 2 8
23	9 9
24	1

**Practical work****1. Solution**

Here Mean =  $\frac{\text{Sum of all observation}}{\text{Number of observation}} = \frac{2152}{50} = 43.04$

Arranging the data in descending order of magnitude

1	2	4	4	5	5	5	10	11	12	13	13	14	19	20
29	29	29	30	31	31	32	33	35	35	36	52	54	61	68
2	2	2	3	2	5	2	6	2	6	2	7	2	7	2
7	4	8	1	9	4	1	1	0	1	1	0	1	2	3
12	6	2	3	2	5	2	6	2	6	2	7	2	7	2
1	0	1	0	1	0	1	0	1	2	3	1	2	6	5

Median = 25.5<sup>th</sup> item = average of 25<sup>th</sup> and 26<sup>th</sup> item =  $\frac{28 + 29}{2} = 28.5$

Mode = Since, mode is ill defined because items 5, 26, 27, 29 are repeated 3 times

First quartile =  $Q_1 = 51/4$  th item = 12.75<sup>th</sup> item

$$= 12^{\text{th}} \text{ item} + (13-12)^{\text{th}} \text{ item} \times 0.75 = 13 + (14-13) \times 0.75 = 13.75$$

Third quartile =  $Q_3 = 3 \times 51/4$  th item = 38.25<sup>th</sup> item

$$= 38^{\text{th}} \text{ item} + (39-38)^{\text{th}} \text{ item} \times 0.25 = 54 + (61-54) \times 0.25 = 55.75$$

Range = Largest item - Smallest item = 165 - 1 = 164

Inter-quartile range =  $Q_3 - Q_1 = 55.75 - 13.75 = 42$


$$\text{Variance} = \frac{\sum x^2}{n} - \left( \frac{\sum x}{n} \right)^2 = \frac{178754}{50} - \left( \frac{2152}{50} \right)^2 = 3575.08 - 1852.44 = 1722.64$$

Standard deviation = SD =  $\sqrt{1722.64} = 41.5$


$$\text{Coefficient of variation} = CV = \frac{SD}{\text{Mean}} \times 100 = \frac{41.5}{43.04} \times 100 = 96.42$$

**Boxplot**

(1, 13.75, 28.5, 55.75, 165)



## Boxplot



### Chapter 3

#### 1. On throwing dice

Sample space  $\{(1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6), (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6), (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6), (5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6)\}$

Let A = sum greater than 9, B = Neither 8 nor 10

$$N = 36$$

$$m(A) = 6$$

$$m(B) = 36 - 7 = 28$$

$$P(A) = \frac{m(A)}{N} = \frac{6}{36} = \frac{1}{6}; \quad P(B) = \frac{m(B)}{N} = \frac{28}{36} = \frac{7}{9}$$

#### 2. A secretary

$$N = {}^5P_5 = 5! = 120$$

Let event of putting letters in correct envelop = A

$$m(A) = 1$$

$$P(\bar{A}) = 1 - P(A) = 1 - \frac{m(A)}{N} = 1 - \frac{1}{120} = \frac{119}{120}$$

#### 3. Solution.

Leap year contains 366 days

A year has 52 complete weeks =  $52 \times 7 = 364$  days

Remaining days = 2

$\therefore$  Sample space of remaining 2 days

$\therefore$  (Sun - Mon, Mon - Tue, Tue - Wed, Wed - Thu, Thu - Fri, Fri - Sat, Sat - Sun)

Favourable no. of cases for Saturday (m) = 2

Total no. of cases (N) = 7

$$P = \frac{m}{N} = \frac{2}{7}$$

#### 4. A problem in

$$P(A) = \frac{1}{5}, P(B) = \frac{2}{5}, P(C) = \frac{3}{5}$$

$$P(A \cup B \cup C) = 1 - P(\overline{A \cup B \cup C})$$

$$= 1 - P(\bar{A} \cap \bar{B} \cap \bar{C})$$

$$= 1 - P(\bar{A}) P(\bar{B}) P(\bar{C})$$

$$= 1 - [1 - P(A)] [1 - P(B)] [1 - P(C)]$$

$$= 1 - \left[1 - \frac{1}{5}\right] \left[1 - \frac{2}{5}\right] \left[1 - \frac{3}{5}\right]$$

$$= 1 - \frac{4}{5} \times \frac{3}{5} \times \frac{2}{5} = 1 - \frac{24}{125} = \frac{121}{125}$$

5. The odds against

For B

$$\frac{N-m}{m} = \frac{4}{3}$$

$$\text{Or, } P(B) = \frac{m}{N} = \frac{3}{3+4} = \frac{3}{7}$$

For A

$$\frac{m}{N-m} = \frac{7}{6}$$

$$P(A) = \frac{m}{N} = \frac{7}{7+6} = \frac{7}{13}$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$= \frac{3}{7} + \frac{7}{13} - \frac{3}{7} \times \frac{7}{13} = \frac{39+49-21}{91} = \frac{67}{91}$$

6. From a pack of

Total cards = 52, selected = 2

Let king = K, Queen = Q, Face card = F, Spade = S, Club = C

$$N = {}^{52}C_2 = 1326$$

$$P(KQ) = \frac{m(KQ)}{N} = \frac{{}^4C_1 \times {}^4C_1}{1326} = \frac{4 \times 4}{1326} = 0.012$$

$$P(2F) = \frac{m(2F)}{N} = \frac{{}^{12}C_2}{1326} = \frac{66}{1326} = 0.049$$

$$P(SC) = \frac{m(SC)}{N} = \frac{{}^{13}C_1 \times {}^{13}C_1}{1326} = \frac{13 \times 13}{1326} = 0.127$$

7. Probability that a

Let Numerical method = M, Computer graphic = S

$$P(M) = 0.75, P(S) = 0.85$$

$$P(M \cap S) = P(M) P(S) = 0.75 \times 0.85 = 0.6375$$

$$P(M \cup S) = P(M) + P(S) - P(M \cap S) = 0.75 + 0.85 - 0.6375 = 0.962$$

8.

$$P(A) = 0.4, P(B) = 0.3, P(A \cup B) = 0.58$$

Now,

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\text{Or, } 0.58 = 0.4 + 0.3 - P(A \cap B)$$

$$\text{Or, } P(A \cap B) = 0.7 - 0.58 = 0.12$$

$$\text{Here, } P(A) P(B) = 0.4 \times 0.3 = 0.12$$

$$P(A \cap B) = P(A) P(B)$$

Hence, A and B are independent.

## 9. A and B play

Prob. of A's getting 2,  $P(A) = \frac{1}{6}$

Prob. not getting 2,  $P(\bar{A}) = 1 - P(A) = 1 - \frac{1}{6} = \frac{5}{6}$

Prob. of B's getting 2,  $P(B) = \frac{1}{6}$

Prob. of not getting 2,  $P(\bar{B}) = \frac{5}{6}$

If A begins the game then prob. of A's winning

$$= P(A \cup \bar{A}\bar{B} A \cup \bar{A}\bar{B} \bar{A} \cup \dots)$$

$$= P(A) + P(\bar{A}) P(\bar{B}) P(A) + P(\bar{A}) P(\bar{B}) P(\bar{A}) P(\bar{B}) P(A) + \dots$$

$$= \frac{1}{6} + \frac{5}{6} \times \frac{5}{6} \times \frac{1}{6} + \frac{5}{6} \times \frac{5}{6} \times \frac{5}{6} \times \frac{5}{6} \times \frac{1}{6} + \dots$$

$$= \frac{\frac{1}{6}}{1 - \frac{25}{36}} = \frac{1}{6} \times \frac{36}{11} = \frac{6}{11}$$

$$\text{Prob. of B's winning} = 1 - \text{Prob. of A's winning} = 1 - \frac{6}{11} = \frac{5}{11}$$

## 10. The probability that

Let defective itehing = A, crack defect = B

$$P(A) = 0.12, P(B) = 0.29, P(A \cap B) = 0.07$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.12 + 0.29 - 0.07 = 0.34$$

## 11. The probability that

Let 50 yrs old man alive at 60 = A

45 yrs woman alive at 55 = B

$$P(A) = 0.83, P(B) = 0.87$$

$$\text{i. } P(A \cap B) = P(A) P(B) = 0.83 \times 0.87 = 0.722$$

$$\text{ii. } P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.83 + 0.87 - 0.722 = 0.978$$

## 12. A card is drawn

Total cards = 52, Cards selected = 1

Let, King = k, Queen = Q, Jack = J, Club = C

$$N = 52, m(k) = 4, m(Q) = 4, m(J) = 4, m(C) = 13, m(J \cap C) = 1$$

$$\text{i. } P(k \cup Q) = P(k) + P(Q)$$

$$= \frac{m(k)}{N} + \frac{m(Q)}{N} = \frac{4}{52} + \frac{4}{52} = \frac{2}{13}$$

$$\text{ii. } P(J \cup C) = P(J) + P(C) - P(J \cap C)$$

$$= \frac{m(J)}{N} + \frac{m(C)}{N} - \frac{m(J \cap C)}{N} = \frac{4}{52} + \frac{13}{52} - \frac{1}{52} = \frac{16}{52} = \frac{4}{13}$$

13. Two cards are

Total cards = 52, Black cards = 26, Red cards = 26, Ace card = 4

Selected cards = 2 in which are after other

Let Black card = B, red card = R, Ace = A

$$\text{i. } P((B_1 \cap R_{II})) = P(B_1) P(R_{II}/B_1) = \frac{26}{52} \times \frac{26}{51} = \frac{13}{51}$$

$$\text{ii. } P(A_1 \cap A_{II}) = P(A_1) P(A_{II}/A_1) = \frac{4}{52} \times \frac{3}{51} = \frac{1}{221}$$

14. A book case contains

Total books = 6 DL + 9 MP = 15 books

Books selected = 4

$$\text{i. } P(4DL_I \cap 4MP_{II}) = P(4DL_I) P(4MP_{II}) = \frac{6C_4}{15C_4} \times \frac{9C_4}{15C_4} = \frac{15 \times 126}{1365 \times 1365} = 0.001$$

$$\text{ii. } P(4DL_I \cap 4MP_{II}) = P(4DL_I) \times P(4MP_{II}/4DL_I) = \frac{6C_4}{15C_4} \times \frac{9C_4}{11C_4} = \frac{15 \times 125}{1365 \times 330} = 0.004$$

15. The student body

Let girls = G, Boys = B, Interest in sports = S

$$P(G) = 60\% = 0.6, P(B) = 40\% = 0.4$$

$$P(G \cap S) = 40\% \text{ of } 60\% = 0.4 \times 0.6 = 0.24$$

$$P(B \cap S) = 60\% \text{ of } 40\% = 0.6 \times 0.4 = 0.24$$

$$P(S/G) = P(GS)/P(G) = \frac{0.24}{0.6} = \frac{2}{5}$$

16. A and B toss

$$\text{Prob. of A's getting head } P(A) = \frac{1}{2}$$

$$\text{Prob. of not getting head } P(\bar{A}) = 1 - P(A) = \frac{1}{2}$$

$$\text{Prob. of B's getting head } P(B) = \frac{1}{2}$$

$$\text{Prob. of not getting } P(\bar{B}) = 1 - P(B) = \frac{1}{2}$$

If A starts game

$$\text{Prob. of A's winning} = P(A \cup \bar{A} \bar{B} A \cup \bar{A} B \bar{A} B A \cup \dots)$$

$$= P(A) + P(\bar{A}) P(\bar{B}) P(A) + P(\bar{A}) P(\bar{B}) P(\bar{A}) P(\bar{B}) P(A) + \dots$$

$$= \frac{1}{2} + \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} + \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} + \dots$$

$$= \frac{\frac{1}{2}}{1 - \frac{1}{2} \times \frac{1}{2}} = \frac{1}{2} \times \frac{4}{3} = \frac{2}{3}$$

$$\text{Prob. of B's winning} = 1 - \text{Prob. of A's winning} = 1 - \frac{2}{3} = \frac{1}{3}$$

17. A person is known

Let a person A and another B

$$\text{Here } P(A) = \frac{4}{5}, P(B) = \frac{3}{4}$$

$$P(A \cap B) = P(A) \times P(B) = \frac{4}{5} \times \frac{3}{4} = \frac{3}{5}$$

18. A six faced dice

Let Odd = O and even = E

$$P(E) = 2/3, P(O) = 1/3$$

$$P(\text{Even}) = P(\text{EE or OO}) = P(E)P(E) + P(O)P(O) = \frac{2}{3} \times \frac{2}{3} + \frac{1}{3} \times \frac{1}{3} = \frac{4}{9} + \frac{1}{9} = \frac{5}{9}$$

19. In tossing a coin

Let head = H, tail = T

Sample space = {HHH, HHT, HTH, THH, HTT, THT, TTH, TTT}

$$N = 8$$

$$P(\text{No. of heads}) = \frac{1}{8}$$

$$P(\text{Two heads}) = \frac{3}{8}$$

$$P(\text{At least two heads}) = \frac{4}{8} = \frac{1}{2}$$

20. There are three traffic

$$P(G) = 0.7, P(R) = 1 - P(G) = 1 - 0.7 = 0.3$$

Sample space = {RRR, RRG, GRR, GRG, RGG, GGR, GGG}

$$P(\text{stop no. more than one}) = P(G R G \cup R G G \cup G G R \cup G G G)$$

$$= P(G) P(R) P(G) + P(R) P(G) P(G) + P(G) P(G) P(R) + P(G) P(G) P(G)$$

$$= 0.7 \times 0.3 \times 0.7 + 0.3 \times 0.7 \times 0.7 + 0.7 \times 0.7 \times 0.3 + 0.7 \times 0.7 \times 0.7$$

$$= 0.441 + 0.343 = 0.784$$

21. The odds that

Let first critics A, second B and third C.

$$\text{For } A, \frac{m}{N-m} = \frac{3}{2}$$

$$\text{For } B, \frac{m}{N-m} = \frac{4}{3}$$

$$P(A) = \frac{m}{N} = \frac{3}{3+2} = \frac{3}{5}$$

$$P(A) = \frac{m}{N} = \frac{4}{3} = \frac{4}{7}$$

$$\text{For } C, \frac{m}{N-m} = \frac{2}{3}$$

$$P(C) = \frac{m}{N} = \frac{2}{2+3} = \frac{2}{15}$$

$$P(\text{majority will be favourable}) = P(\bar{A}\bar{B}\bar{C} \cup \bar{A}\bar{B}C \cup A\bar{B}\bar{C} \cup ABC)$$

$$= P(A) P(B) P(\bar{C}) + P(\bar{A}) P(B) P(C) + P(A) P(\bar{B}) P(C) + P(A) P(B) P(C)$$

$$= \frac{3}{5} \times \frac{4}{7} \times \left(1 - \frac{2}{5}\right) + \left(1 - \frac{3}{5}\right) \times \frac{4}{7} \times \frac{2}{5} + \frac{3}{5} \times \left(1 - \frac{4}{7}\right) \times \frac{2}{5} + \frac{3}{5} \times \frac{4}{7} \times \frac{2}{5}$$

$$= \frac{36}{175} + \frac{16}{175} + \frac{18}{175} + \frac{24}{175} = \frac{94}{175} = 0.537$$

22. Solution.

Bank Credit card		Travel & Entertainment Credit card		Total
	Yes	No		{ } 200 }
Yes	60	60	120	
No	15	65	80	
Total	75	125	200	

- a.  $P(\text{the student has a bank credit card}) = \frac{120}{200} = \frac{3}{5} = 0.6$
- b.  $P(\text{the student has a bank credit card and a travel and entertainment card}) = \frac{60}{200} = 0.3$
- c.  $P(\text{the student has a bank credit card or a travel and entertainment card}) = P(B \text{ or } T) = P(B) + P(T) - P(B \& T) = \frac{120}{200} + \frac{75}{200} - \frac{60}{200} = \frac{135}{200} = 0.675$
- d.  $P(\text{he or she has a travel and entertainment card}) = \frac{125}{200} = 0.625$

23. Solution

a.

Gender	Health Club Facility		Total
	Used	Not used	
Male	65	105	170
Female	45	35	80
total	110	140	250

- b.  $P(\text{an employee chosen at random is a female and has used the health club facilities}) = \frac{45}{250} = 0.18$
- c.  $P(\text{an employee chosen at random is a male}) = \frac{170}{250} = 0.68$
- d.  $P(\text{an employee chosen at random is a male or has not used the health club facilities}) = P(M \text{ or } NH) = P(M) + P(NH) - P(M \& NH)$   
 $= \frac{170}{250} + \frac{140}{250} - \frac{105}{250} = \frac{100}{250} = 0.82$

24. Solution.

$$\text{Total IC chips} = 5000$$

$$\text{IC chips manufactured by company } x = 1000,$$

$$y = 5000 - 1000 = 4000$$

Let  $A_1$  = IC chips manufactured by company  $x$

$A_2$  = IC chips manufactured by company  $y$

$$P(A_2/D) = \frac{P(A_2) P(D/A_2)}{P(A_1) P(D/A_1) + P(A_2) P(D/A_2) + P(A_3) P(D/A_3)}$$

$$= \frac{0.35 \times 0.04}{0.0345} = 0.405$$

$$P(A_3/D) = \frac{P(A_3) P(D/A_3)}{P(A_1) P(D/A_1) + P(A_2) P(D/A_2) + P(A_3) P(D/A_3)}$$

$$= \frac{0.4 \times 0.02}{0.0345} = \frac{0.008}{0.0345} = 0.231$$

## 31. Solution.

Denote the events,

$A = \{\text{errors in module I}\}$ ,  $B = \{\text{errors in module II}\}$ ,  $C = \{\text{crash}\}$

Further,  $\{\text{errors in module I alone}\} = A \setminus B = A \setminus (A \cap B)$

$\{\text{errors in module II alone}\} = B \setminus A = B \setminus (A \cap B)$ .

It is given that  $P\{A\} = 0.2$ ,  $P\{B\} = 0.4$ ,  $P(A \cap B) = (0.2)(0.4) = 0.08$ ,

By independence,

$P\{C|A \setminus B\} = 0.5$ ,  $P\{C|B \setminus A\} = 0.8$ , and  $P\{C|A \cap B\} = 0.9$

We need to compute  $P\{A \cap B|C\}$ . Since  $A$  is a union of disjoint events  $A \setminus B$  and  $A \cap B$ , we compute

$$P\{A \setminus B\} = P\{A\} - P\{A \cap B\} = 0.2 - 0.08 = 0.12.$$

Similarly,

$$P\{B \setminus A\} = 0.4 - 0.08 = 0.32$$

Events  $(A \setminus B)$ ,  $(B \setminus A)$ ,  $A \cap B$ , and  $(\overline{A \cup B})$  form a partition of  $\Omega$ , because they are mutually exclusive and exhaustive. The last of them is the event of no errors in the entire program. Given this event, the probability of a crash is 0. Notice that  $A$ ,  $B$ , and  $(A \cap B)$  are neither mutually exclusive nor exhaustive, so they cannot be used for the Bayes Rule. Now organize the data.

Location of errors	Probability of a crash
$P\{A \setminus B\} = 0.12$	$P\{C A \setminus B\} = 0.5$
$P\{B \setminus A\} = 0.32$	$P\{C B \setminus A\} = 0.8$
$P\{A \cap B\} = 0.08$	$P\{C A \cap B\} = 0.9$
$P\{\overline{A \cup B}\} = 0.48$	$P\{C \overline{A \cup B}\} = 0$

Combining the Bayes Rule and the Law of Total Probability,

$$P\{A \cap B|C\} = \frac{P\{A|A \cap B\} P\{A \cap B\}}{P\{C\}},$$

Where

$$P\{C\} = P\{C|A \setminus B\} P\{A \setminus B\} + P\{C|B \setminus A\} P\{B \setminus A\} + P\{C|A \cap B\} P\{A \cap B\} + P\{C|\overline{A \cup B}\} P\{\overline{A \cup B}\}$$

Then

$$P\{A \cap B|C\} = \frac{(0.9)(0.08)}{(0.5)(0.12) + (0.8)(0.32) + (0.9)(0.08) + 0} = 0.1856$$

## 32. Solution.

Let urn I =  $A_1$ , urn II =  $A_2$ , urn III =  $A_3$

$$P(A_1) = P(A_2) = P(A_3) = \frac{1}{3}$$

Let R = Real chips

$$P(R/A_1) = \frac{6}{10}; \quad P(R/A_2) = \frac{2}{8}; \quad P(R/A_3) = \frac{1}{9}$$

$$P(A_1/R) = ?$$

$$\begin{aligned} P(A_1/R) &= \frac{P(A_1) P(R/A_1)}{P(A_1) P(R/A_1) + P(A_2) P(R/A_2) + P(A_3) P(R/A_3)} \\ &= \frac{\frac{1}{3} \times \frac{6}{10}}{\frac{1}{3} \times \frac{6}{10} + \frac{1}{3} \times \frac{2}{8} + \frac{1}{3} \times \frac{1}{9}} = 0.624 \end{aligned}$$

Let the chips drawn are red without replacement =  $R_w$

$$P(R_w/A_1) = \frac{6}{10} \times \frac{5}{9} = \frac{1}{3} \quad P(R_w/A_2) = \frac{2}{8} \times \frac{1}{7} = \frac{1}{28}$$

$$P(R_w/A_3) = \frac{1}{9} \times \frac{0}{8} = 0 \quad P(A_1/R_w) = ?$$

$$\begin{aligned} P(A_1/R_w) &= \frac{P(A_1) P(R_w/A_1)}{P(A_1) P(R_w/A_1) + P(A_2) P(R_w/A_2) + P(A_3) P(R_w/A_3)} \\ &= \frac{\frac{1}{3} \times \frac{1}{3}}{\frac{1}{3} \times \frac{1}{3} + \frac{1}{3} \times \frac{1}{28} + \frac{1}{3} \times 0} = 0.903 \end{aligned}$$

$$\begin{aligned} P(A_3/R_w) &= \frac{P(A_3) P(R_w/A_3)}{P(A_1) P(R_w/A_1) + P(A_2) P(R_w/A_2) + P(A_3) P(R_w/A_3)} \\ &= \frac{\frac{1}{3} \times 0}{\frac{1}{3} \times \frac{1}{3} + \frac{1}{3} \times \frac{1}{28} + \frac{1}{3} \times 0} = 0 \end{aligned}$$

## Chapter 4

### 1. Solution

a. Here for Population Mean  $\bar{Y} = \frac{\sum Y}{N} = \frac{0 + 1 + 3 + 8 + 9}{5} = 4.2$

$$\begin{aligned}\text{Population Variance, } S^2 &= \frac{1}{N-1} \sum (Y_i - \bar{Y})^2 \\ &= \frac{1}{N-1} \{ \sum Y_i^2 - N\bar{Y}^2 \} \\ &= \frac{1}{4} \{ 0^2 + 1^2 + 3^2 + 8^2 + 9^2 - 5(4.2)^2 \} = 16.7\end{aligned}$$

- b. Here the sample of size 2 from population of size 6 can be drawn in  $C(5,2)$  ways = 10 ways

Hence the samples are: (0,1) (0,3) (0,8) (0,9) (1,3) (1,8) (1,9) (3,8) (3,9) (8,9)

Sample No.	Sample values ( $y_i$ )	Sample mean ( $\bar{y}_i$ )	$\{\bar{y}_i - E(\bar{y})\}^2$	$s_i^2 = \frac{1}{n-1} \sum (y_i^2 - \bar{y})^2$
1	(0,1)	0.5	13.69	0.5
2	(0,3)	1.5	7.29	4.5
3	(0,8)	4	0.04	32
4	(0,9)	4.5	0.09	40.5
5	(1,3)	2	4.84	2
6	(1,8)	4.5	0.09	24.5
7	(1,9)	5	0.64	32
8	(3,8)	5.5	1.69	12.5
9	(3,9)	6	3.24	18
10	(8,9)	8.5	18.49	0.5
Total		42	50.1	167

c.  $E(\bar{y}) = \frac{1}{C(N,n)} \sum_i y_i = \frac{42}{10} = 4.2$

d.  $V(\bar{y}) = \frac{1}{C(N,n)} \sum_i \bar{y}_i - E(\bar{y})^2 = \frac{50.1}{10} = 5.01$

e. Yes,  $E(\bar{y}) = \bar{Y}$  verified

Here, sample mean is unbiased estimate of the population mean

$$E(s^2) = \frac{1}{C(N,n)} \sum_i s_i^2 = \frac{167}{10} = 16.7. \text{ Hence, } E(s^2) = S^2$$

f. Now,  $V(\bar{y}) = \frac{(1-f)S^2}{n} = \left(1 - \frac{2}{5}\right) \times \frac{16.7}{2} = 5.01$  Hence, Formula verified.

**40 CHAPTER - 4 | A Complete Solutions of Statistics I for BSc. CSIT**

**2. Solution.**

Here, Population size ( $N$ ) = 5, Sample size ( $n$ ) = 2.

- a. Calculation of population mean and population variance:

$Y_i$	$Y_i - \bar{Y}$	$(Y - \bar{Y})^2$
0	-3.2	10.24
1	-2.2	4.84
3	-0.2	0.04
5	1.8	3.24
7	3.8	14.44
$\sum Y = 16$		$\sum(Y - \bar{Y})^2 = 32.8$

$$\text{Population mean } (\mu \text{ or } \bar{Y}) = \frac{\sum Y_i}{N} = \frac{16}{5} = 3.2$$

$$\text{Population variance } (\sigma^2) = \frac{\sum(Y_i - \bar{Y})^2}{N} = \frac{32.8}{5} = 6.56$$

- b. Number of possible samples of size 2 that can be drawn from the population of size  $N = 5$  by using with replacement technique is: Number of possible samples =  $N^n = 5^2 = 25$

Thus the possible samples are: (0,0), (0,1), (0,3), (0,5), (0,7), (1,0), (1,1), (1,3), (1,5), (1,7), (3,0), (3,1), (3,3), (3,5), (3,7), (5,0), (5,1), (5,3), (5,5), (5,7), (7,0), (7,1), (7,3), (7,5), (7,7).

- c. Calculation of sample means and variance of the sampling distribution means:

S. No.	Sample Values ( $Y_i$ )	Sample Means ( $\bar{y}_i$ )	$(\bar{y}_i - \bar{y})$	$(\bar{y}_i - \bar{y})^2$	$s^2 = \frac{1}{n-1} \{ \sum y_i^2 - n\bar{y}^2 \}$
1	(0,0)	0	-3.2	10.2	0
2	(0,1)	0.5	-2.7	7.29	0.5
3	(0,3)	1.5	-1.7	2.89	4.5
4	(0,5)	2.5	-0.7	0.49	12.5
5	(0,7)	3.5	0.3	0.09	24.5
6	(1,0)	0.5	-2.7	7.29	0.5
7	(1,1)	1	-2.2	4.84	0
8	(1,3)	2	-1.2	1.44	2
9	(1,5)	3	-0.2	0.04	8
10	(1,7)	4	0.8	0.64	18
11	(3,0)	1.5	-1.7	2.89	4.5
12	(3,1)	2	-1.2	1.44	2
13	(3,3)	3	-0.2	0.04	0
14	(3,5)	4	0.8	0.64	2
15	(3,7)	5	1.8	3.24	8
16	(5,0)	2.5	-0.7	0.49	12.5

17	(5,1)	3	-0.2	0.04	8
18	(5,3)	4	0.8	0.64	2
19	(5,5)	5	1.8	3.24	0
20	(5,7)	6	2.8	7.84	2
21	(7,0)	3.5	0.3	0.09	24.5
22	(7,1)	4	0.8	0.64	18
23	(7,3)	5	1.8	3.24	8
24	(7,5)	6	2.8	7.84	2
25	(7,7)	7	3.8	14.4	0
		80		82	

$$\text{Mean of the sample means } (\bar{y}) = \frac{\sum \bar{y}_i}{\text{No. of samples (N^n)}} = \frac{80}{25} = 3.2$$

Since the mean of the sample means ( $\bar{y}$ ) = 3.2 is equal to the population mean ( $\bar{Y}$ ) = 3.2, so we can conclude that the mean of the sampling distribution of the sample means is equal to the population mean.

d. Variance of the sample means is

$$V(\bar{y}) = \frac{\sum (\bar{y}_i - \bar{y})^2}{\text{No. of samples}} = \frac{82}{25} = 3.28 \text{ or}$$

$$V(\bar{y}) = \frac{\sigma^2}{n} = \frac{6.56}{2} = 3.28$$

e. variance of sample means gives the same value as calculated from (d)

3. Solution:

Here  $N = 800$ ,  $n = 120$

$$\begin{aligned} \text{Var}(\bar{y}_{st})_{\text{prop}} &= \left( \frac{1}{n} - \frac{1}{N} \right) \sum W_i S_i^2 \\ &= \left( \frac{1}{n N} - \frac{1}{N^2} \right) \sum N_i S_i^2 \quad \left\{ W_i = \frac{N_i}{N} \right\} \\ &= \left( \frac{1}{120 \times 800} - \frac{1}{800^2} \right) \{200 \times 36 + 300 \times 64 + 300 \times 144\} \\ &= 0.616 \end{aligned}$$

$$\begin{aligned} \text{Var}(\bar{y}_{st})_{\text{opt}} &= \frac{1}{n} (\sum W_i S_i)^2 - \frac{1}{N} \sum W_i S_i^2 \\ &= \frac{1}{n N^2} (\sum N_i S_i)^2 - \frac{1}{N^2} \sum N_i S_i^2 \\ &= \frac{1}{120 \times 800^2} (200 \times 6 + 300 \times 8 + 300 \times 12)^2 - \frac{1}{800^2} (200 \times 36 + 300 \times 64 + 300 \times 144) \\ &= 0.675 - 0.1087 \\ &= 0.566 \end{aligned}$$

## 4. Solution.

Here,  $N = 800$ ,  $N_1 = 200$ ,  $N_2 = 300$ ,  $N_3 = 300$   $S_1 = 6$ ,  $S_2 = 8$ ,  $S_3 = 12$ ,  $n = 120$   
 $N_1S_1 = 200 \times 6 = 1200$ ,  $N_2S_2 = 300 \times 8 = 2400$ ,  $N_3S_3 = 300 \times 12 = 3600$ ,  
 $\sum N_i S_i = 1200 + 2400 + 3600 = 7200$

## (i) Under proportional allocation

$$\frac{n_i}{N_i} = \frac{n}{N}$$

$$n_i = \frac{n}{N} N_i, \quad i = 1, 2, 3$$

$$n_1 = \frac{120}{800} \times 200 = 30$$

$$n_2 = \frac{120}{800} \times 300 = 45$$

$$n_3 = \frac{120}{800} \times 300 = 45$$

## ii) Under Neyman's optimum allocation

$$\frac{n_i}{n} = \frac{N_i S_i}{\sum N_i S_i}$$

$$n_i = \frac{n N_i S_i}{\sum N_i S_i}$$

$$n_1 = \frac{120 \times 1200}{7200} = 20$$

$$n_2 = \frac{120 \times 240}{7200} = 40$$

$$n_3 = \frac{120 \times 3600}{7200} = 60$$

In the proportional allocation the sample fraction  $n_i/N_i$  is constant for each stratum

$$\frac{n_i}{N_i} = \frac{n}{N} = C$$

$n_i$  proportional to  $N_i$

In Neyman's optimum allocation  $V(\bar{y}_{st})$  is minimum for fixed  $n$  if  $n_i$  proportional to  $N_i S_i$ .

Hence in former one  $n_i$  depends on  $N_i$  but in later one  $n_i$  also depends on  $S_i$

## Chapter 5

### 1. Solution.

x	P(x)
0	a
1	3a
2	5a
3	7a
4	9a
5	11a
6	13a
7	15a
8	17a
$\Sigma P(x) = 81a$	

Since,  $\Sigma P(x) = 1$

$$\Rightarrow 81a = 1 \quad \Rightarrow a = \frac{1}{81}$$

$$P(x < 3) = P(x = 0) + P(x = 1) + P(x = 2)$$

$$= a + 3a + 5a = 9a = 9 \times \frac{1}{81} = \frac{1}{9}$$

$$P(x \geq 3) = P(x = 3) + P(x = 4) + P(x = 5) + P(x = 6) + P(x = 7) + P(x = 8)$$

$$= 7a + 9a + 11a + 13a + 15a + 17a$$

$$= 72a$$

$$= 72 \times \frac{1}{81} = \frac{8}{9}$$

$$P(0 < x < 5) = P(x = 1) + P(x = 2) + P(x = 3) + P(x = 4)$$

$$= 3a + 5a + 7a + 9a$$

$$= 24a = 24 \times \frac{1}{81} = \frac{8}{27}$$

2. Let x be the number of toss required to get tail.

Let T denotes out tail and H denotes outcome head.

Outcomes	No. of TBS	P(x)	x P(x)
T	1	1/2	1/2
HT	2	1/4	2/4
HHT	3	1/8	3/8
HHHT	4	1/16	4/16
HHHHT	5	1/32	5/32

$$E(x) = \Sigma x P(x) = \frac{1}{2} + \frac{2}{4} + \frac{3}{8} + \frac{4}{16} + \frac{5}{32} + \dots$$

$$\text{Let } S = \frac{1}{2} + \frac{2}{4} + \frac{3}{8} + \frac{4}{16} + \frac{5}{32} \dots$$

$$\frac{S}{2} = \frac{1}{4} + \frac{2}{8} + \frac{3}{16} + \frac{4}{32} \dots \dots \dots$$

$$5 - \frac{S}{2} = \frac{1}{2} + \left(\frac{2}{4} - \frac{1}{4}\right) + \left(\frac{4}{16} - \frac{3}{16}\right) + \dots$$

$$\frac{S}{2} = \frac{\frac{1}{2}}{1 - \frac{1}{2}}$$

$$\frac{S}{2} = 1$$

$$\Rightarrow S = 2$$

$$\therefore E(x) = 2$$

### 3. Solution.

On throwing two dice possible outcomes are (1,1), (1,2), (1,3), (1,4), (1,5), (1,6), (2,1), (2,2), (2,3), (2,4), (2,5), (2,6), (3,1), (3,2), (3,3), (3,4), (3,5), (3,6), (4,1), (4,2), (4,3), (4,4), (4,5), (4,6), (5,1), (5,2), (5,3), (5,4), (5,5), (5,6), (6,1), (6,2), (6,3), (6,4), (6,5), (6,6).

Let  $x$  = Sum of numbers

$x$	$P(x)$	$x P(x)$
2	1/36	2/36
3	2/36	6/36
4	3/36	12/36
5	4/36	20/36
6	5/36	30/36
7	6/36	42/36
8	5/36	40/36
9	4/36	36/36
10	3/36	30/36
11	2/36	22/36
12	1/36	12/36
		$\Sigma x P(x) = \frac{252}{36} = 7$

$$E(x) = \Sigma x P(x) = 7$$

### 4. Solution.

$x$	$P(x)$	$x P(x)$
-2	0.2	- 0.4
-1	$k$	- $k$
0	0.4	0
1	$2k$	$2k$
2	$k$	$2k$
	$\Sigma P(x) = 4k + 0.6$	$\Sigma x P(x) = 3k - 0.4$

Since,  $\sum_{x} x P(x) = 1$

$$\text{or } 4k + 0.6 = 1$$

$$\text{or } 4k = 0.4 \Rightarrow k = 0.1$$

$$(ii) P(x \geq 0) = P(x = 0) + P(x = 1) + P(x = 2) = 0.4 + 2k + k = 0.4 + 3 \times 0.1 = 0.7$$

$$(iii) E(x) = 3k - 0.4 = 3 \times 0.1 - 0.4 = -0.1$$

$$(iv) E(4x + 5) = 4 E(x) + 5 = 4 \times (-0.1) + 5 = 4.6$$

5. Solution.

x	P(x)	$x P(x)$	$x^2 P(x)$
0	0.05	0	0
1	0.35	0.35	0.35
2	0.2	0.40	0.80
3	0.25	0.75	2.25
4	0.15	0.60	2.40
		$\Sigma x P(x) = 2.1$	$\Sigma x^2 P(x) = 5.8$

$$E(x) = \Sigma x P(x) = 2.1$$

$$E(x^2) = \Sigma x^2 P(x) = 5.8$$

$$V(x) = E(x^2) - [E(x)]^2 = 5.8 - (2.1)^2 = 1.39$$

$$S.D. = \sqrt{V(x)} = \sqrt{1.39} = 1.178$$

6. Solution.

x	P(x)	$x P(x)$
700	1/12	700/12
900	1/12	900/12
1100	3/12	3300/12
1300	3/12	3900/12
1500	2/12	3000/12
1700	2/12	3400/12
		$\Sigma x P(x) = 1266.67$

Expected earning = Rs. 1266.67

7. Solution.

Let  $x$  = Profit

x	P(x)	$x P(x)$
$180,000 - 140,000 = 40,000$	0.32	12800
$170,000 - 140,000 = 30,000$	0.55	16500
$150,000 - 140,000 = 10,000$	0.13	1300
		$\Sigma x P(x) = 30600$

$$E(x) = \Sigma x P(x) = 30,600$$

**46 CHAPTER - 5 | A Complete Solutions of Statistics I for BSc. CSIT**

8. Let  $x$  = Rupees

$x$	$P(x)$	$x P(x)$
5000	0.001	5
3000	0.003	9
1000	0.005	5
		$\Sigma x P(x) = 19$

$$\text{Fairy price} = \Sigma x P(x) = 19$$

9. Solution.

Let  $x$  = No. of Video card,  $V$  = Video card,  $N$  = Network card from each urn selected as  $N_I N_{II} N_{III}$ ,  $V_I V_{II} V_{III}$ ,  $N_I V_{II} N_{III}$ ,  $N_I N_{II} V_{III}$ ,  $V_I V_{II} N_{III}$ ,  $V_I N_{II} V_{III}$ ,  $N_I V_{II} V_{III}$ ,  $V_I V_{II} V_{III}$ .

$$I: 3N + 2V, II: 5N + 6V, III: 2N + 4V$$

$x$	$P(x)$	$x P(x)$
0	$\frac{3}{5} \times \frac{5}{11} \times \frac{2}{6} = \frac{30}{330}$	0
1	$\frac{2}{5} \times \frac{5}{11} \times \frac{2}{6} + \frac{3}{5} \times \frac{6}{11} \times \frac{2}{6} + \frac{3}{5} \times \frac{5}{11} \times \frac{4}{6} = \frac{116}{330}$	$\frac{116}{330}$
2	$\frac{2}{5} \times \frac{6}{11} \times \frac{2}{6} + \frac{2}{5} \times \frac{5}{11} \times \frac{4}{6} + \frac{3}{5} \times \frac{6}{11} \times \frac{4}{6} = \frac{136}{330}$	$\frac{272}{330}$
3.	$\frac{2}{5} \times \frac{6}{11} \times \frac{4}{6} = \frac{48}{330}$	$\frac{144}{330}$
		$\Sigma x P(x) = 532/330$

$$E(x) = \Sigma x P(x) = 532/330 = 1.61$$

10. Solution. Let  $x$  = Profit

$x$	$P(x)$	$x P(x)$
5000	0.18	900
8000	0.22	1760
0	0.33	0
-3000	0.27	-810
		$\Sigma x P(x) = 1850$

$$E(x) = \Sigma x P(x) = 1850$$

11. Solution.

$$\text{Since, } \int_{-\infty}^{+\infty} f(x) dx = 1$$

$$\Rightarrow \int_0^1 k(1-x)^2 dx = 1$$

$$\Rightarrow k \left[ \frac{(1-x)^{2+1}}{(2+1)(-1)} \right]_0^1 = 1$$

$$\Rightarrow k \left[ \frac{(1-x)^3}{-3} \right]_0^1 = 1$$

$$\Rightarrow k \left[ 0 + \frac{1}{3} \right] = 1$$

$$\Rightarrow k = 3$$

$$\begin{aligned} P(0 < x < 0.5) &= \int_0^{0.5} f(x) dx = \int_0^{0.5} 3(1-x)^2 dx \\ &= 3 \left[ \frac{(1-x)^3}{-3} \right]_0^{0.5} \\ &= -[(1-x)^3]_0^{0.5} \\ &= -[(1-0.5)^3 - (1-0)^3] = 0.875 \end{aligned}$$

$$\begin{aligned} E(x) &= \int_{-\infty}^{+\infty} x f(x) dx \\ &= \int_0^1 x \cdot 3(1-x)^2 dx \\ &= 3 \int_0^1 x(1-2x+x^2) dx = 3 \int_0^1 (x-2x^2+x^3) dx \\ &= 3 \left[ \frac{x^2}{2} - 2 \frac{x^3}{3} + \frac{x^4}{4} \right]_0^1 \\ &= 3 \left[ \frac{1}{2} - \frac{2}{3} + \frac{1}{4} \right] = 3 \frac{6-8+3}{12} = \frac{1}{4} \end{aligned}$$

12. Solution.

$$\int_{-\infty}^{+\infty} f(x) dx = 1$$

$$\Rightarrow \int_0^1 k x^2 dx = 1$$

$$\Rightarrow k \left[ \frac{x^3}{3} \right]_0^1 = 1$$

$$\Rightarrow \frac{k}{3} = 1$$

$$\Rightarrow k = 3$$

$$P(0 < x < 0.5) = \int_0^{0.5} f(x) dx = \int_0^{0.5} 3x^2 dx = 3 \left[ \frac{x^3}{3} \right]_0^{0.5} = \frac{3}{3} [(0.5)^3] = 0.125$$

$$\begin{aligned} E(x) &= \int_0^1 x f(x) dx = \int_0^1 x \cdot 3x^2 dx \\ &= \int_0^1 x \cdot 3x^2 dx = 3 \int_0^1 x^3 dx = 3 \left[ \frac{x^4}{4} \right]_0^1 = \frac{3}{4} [1^4] = 0.75 \end{aligned}$$

13. Solution.

$$f(x) = \begin{cases} c(10-x)^2 & 0 < x < 10 \\ 0 & \text{otherwise} \end{cases}$$

$$P(1 < x < 2) = ?$$

$$\text{Since, } \int_{-\infty}^{+\infty} f(x) dx = 1$$

$$\text{Or } \int_0^{10} C(10-x)^2 dx = 1$$

$$\text{Or } C \left[ \frac{(10-x)^3}{3(-1)} \right]_0^{10} = 1$$

$$\text{Or } C \left( 0 + \frac{10^3}{3} \right) = 1$$

$$\text{Or } C = \frac{3}{1000}$$

$$P(1 < x < 2)$$

$$= \int_1^2 \frac{3}{1000} (10-x)^2 dx$$

$$= \frac{3}{1000} \left[ \frac{(10-x)^3}{3(-1)} \right]_0^2$$

$$= 0.217$$

14. Solution.

$$f(x) = x e^{-x}, x \geq 0$$

$$E(x) = \int_{-\infty}^{+\infty} x f(x) dx$$

$$= \int_0^{\infty} x x e^{-x} dx$$

$$= \int_0^{\infty} e^{-x} x^2 dx$$

$$= \int_0^{\infty} e^{-x} x^{3-1} dx$$

$$= \sqrt{3} = (3-1)! = 2! = 2$$

15. Solution.

$$f(x) = \begin{cases} 12x(1-x)^2 & 0 < x < 1 \\ 0 & \text{otherwise} \end{cases}$$

$$E(x) = \int_{-\infty}^{+\infty} x f(x) dx$$

$$= \int_0^1 12x(1-x^2) dx$$

$$= 12 \int_0^1 x^2 (1-2x+x^2) dx$$

$$= 12 \int_0^1 (x^2 - 2x^3 + x^4) dx$$

$$= 12 \left[ \frac{x^3}{3} - \frac{2x^4}{4} + \frac{x^5}{5} \right]_0^1$$

$$= 12 \left[ \frac{1}{3} - \frac{1}{2} + \frac{1}{5} \right] = 12 \times \frac{10 - 15 + 6}{30} = 2/5$$

$$E(x^2) = \int x^2 f(x) dx$$

$$\begin{aligned}
 &= \int_0^1 x^2 \cdot 12x(1-x)^2 dx \\
 &= 12 \int x^3 (1-2x+x^2) dx \\
 &= 12 \left[ \frac{x^4}{4} - \frac{2x^5}{5} + \frac{x^6}{6} \right]_0^1 = 1/5
 \end{aligned}$$

16. Solution.

$$f(x) = \begin{cases} 2(1-x) & 0 < x < 1 \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned}
 E(x) &= x f(x) dx \\
 &= \int_0^1 x \cdot 2(1-x) dx \\
 &= 2 \left[ \frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 = 2 \frac{3-2}{6} = \frac{1}{3}
 \end{aligned}$$

$$\begin{aligned}
 E(x^2) &= \int_{-\infty}^{\infty} x^2 f(x) dx \\
 &= \int_0^1 x^2 \cdot 2(1-x) dx \\
 &= 2 \int_0^1 (x^2 - x^3) dx \\
 &= 2 \left[ \frac{x^3}{3} - \frac{x^4}{4} \right]_0^1 \\
 &= 2 \left[ \frac{1}{3} - \frac{1}{4} \right] = 2 \times \frac{4-3}{12} = \frac{1}{6}
 \end{aligned}$$

$$\begin{aligned}
 V(x) &= E(x^2) - [E(x)]^2 \\
 &= \frac{1}{6} - \left(\frac{1}{3}\right)^2 = \frac{3-2}{18} = \frac{1}{18}
 \end{aligned}$$

$$\text{S.D.} = \sqrt{V(x)} = \sqrt{\frac{1}{18}} = 0.235$$

17. Solution.

$$f(x) = 3x^2; 0 \leq x \leq 1$$

$$a = ?, b = ?$$

$$(i) P(x \leq a) = P(x > a)$$

$$\Rightarrow \int_0^a f(x) dx = \int_1^1 f(x) dx$$

$$\Rightarrow \int_0^a 3x^2 dx = \int_a^1 3x^2 dx$$

$$\Rightarrow 3 \left[ \frac{x^3}{3} \right]_0^a = \left[ \frac{x^3}{3} \right]_a^1$$

$$\Rightarrow a^3 = 1 - a^3$$

$$\Rightarrow 2a^3 = 1$$

$$\Rightarrow a^3 = \frac{1}{2}$$

$$\Rightarrow a = \sqrt[3]{0.5} = 0.793$$

$$(ii) P(x > b) = 0.05$$

$$\Rightarrow \int_b^1 f(x) dx = 0.05$$

$$\Rightarrow \int_b^1 3x^2 dx = 0.05$$

$$\Rightarrow \left[ 3 \frac{x^3}{3} \right]_b^1 = 0.05$$

$$\Rightarrow 1 - b^3 = 0.05$$

$$\Rightarrow b^3 = 0.95$$

$$\Rightarrow b = \sqrt[3]{0.95} = 0.98$$

18. Solution.

$$f(x) = ke^{-x/\sigma} \quad 0 < x < \infty, \sigma > 0$$

$$\text{Since, } \int_{-\infty}^{+\infty} f(x) dx = 1$$

$$\Rightarrow \int_0^{\infty} k e^{-x/\sigma} dx = 1$$

$$\text{Let } \frac{x}{\sigma} = y \text{ then } dx = \sigma dy$$

Where  $x = 0, y = 0$  when  $x = \infty, y = \infty$

$$\therefore \int_0^{\infty} k e^{-x/\sigma} dx = 1$$

$$\Rightarrow k \sigma \int_0^{\infty} e^{-y} y^{1-1} dy = 1$$

$$\Rightarrow k \sigma \times 1 = 1$$

$$\Rightarrow k = \frac{1}{\sigma}$$

$$\therefore f(x) = \frac{1}{\sigma} e^{-x/\sigma}$$

$$E(x) = \int x f(x) dx$$

$$= \int_0^{\infty} x \frac{1}{\sigma} e^{-x/\sigma} dx$$

$$= \int_0^{\infty} y e^{-y} \sigma dy$$

$$= \sigma \int_0^{\infty} e^{-y} y^{2-1} dy$$

$$= \sigma \sqrt{2}$$

$$= \sigma 1 1$$

$$= \sigma$$

$$E(x^2) = \int x^2 f(x) dx$$

$$\begin{aligned}
 &= \int_0^{\infty} x^2 \frac{1}{\sigma} e^{-x/\sigma} dx \\
 &= \int_0^{\infty} x \frac{x}{\sigma} e^{-x/\sigma} \sigma dy \\
 &= \int_0^{\infty} \sigma_y \cdot y e^{-y} \sigma dy \\
 &= \sigma^2 \int_0^{\infty} e^{-y} y^2 dy \\
 &= \sigma^2 \int_0^{\infty} e^{-y} y^{3-1} dy \\
 &= \sigma^2 \sqrt{3} \\
 &= \sigma^2 21 \\
 &= 2 \sigma^2
 \end{aligned}$$

$$\begin{aligned}
 v(x) &= E(x^2) = [E(x)]^2 \\
 &= 2\sigma^2 - \sigma^2 = \sigma^2
 \end{aligned}$$

19. Solution.

x	P(x)	x P(x)	x <sup>2</sup> P(x)	x <sup>3</sup> P(x)	x <sup>4</sup> P(x)
-3	1/9	-3/9	9/9	-27/9	81/9
-2	1/9	-2/9	4/9	-8/9	16/9
-1	1/9	-1/9	1/9	-1/9	1/9
0	1/3	0	0	0	0
1	1/9	1/9	1/9	1/9	1/9
2	1/9	2/9	4/9	8/9	16/9
3	1/9	3/9	9/9	27/9	81/9
	$\Sigma P(x) = 0$	$\Sigma x P(x) = 28/9$	$\Sigma x^2 P(x) = 28/9$	$\Sigma x^3 P(x) = 0$	$\Sigma x^4 P(x) = 196/9$

First four moments about origin,

$$\mu_1^1 = E(x) = \Sigma x P(x) = 0$$

$$\mu_2^1 = E(x^2) = \Sigma x^2 P(x) = 28/9 = 3.11$$

$$\mu_3^1 = E(x^3) = \Sigma x^3 P(x) = 0$$

$$\mu_4^1 = E(x^4) = \Sigma x^4 P(x) = 196/9 = 21.77$$

First four moments about mean

$$\mu_1 = 0$$

$$\mu_2 = \mu_2^1 - \mu_1^1 \cdot \mu_1^1 = 3.11 - 0^2 = 3.11$$

$$\mu_3 = \mu_3^1 - 3\mu_2^1 \mu_1^1 + 2\mu_1^1 \cdot \mu_1^1 = 0 - 3 \times 3.11 \times 0 + 2 \times 0^3 = 0$$

$$\mu_4 = \mu_4^1 - 4\mu_3^1 \mu_1^1 + 6\mu_2^1 \mu_1^1 \cdot \mu_1^1 + 3\mu_1^1 \cdot \mu_1^1 = 21.77 - 0 + 0 - 0 = 21.77$$

Measure of central tendency,  $E(x) = 0$

Measure of skewness

$$\beta_1 = \frac{\mu_3^2}{\mu_2^3} = \frac{0^2}{(3.11)^3} = 0$$

Measure of kurtosis

$$\beta_2 = \frac{\mu_4}{\mu_2^2} = \frac{21.77}{(3.11)^2} = 2.25$$

## 20. Solution.

$$f(x) = kx(2-x) \quad 0 \leq x \leq 2$$

$$\text{Since, } \int_{-x}^x f(x) dx = 1$$

$$\Rightarrow \int_0^2 kx(2-x) dx = 1$$

$$\Rightarrow k \int_0^2 (2x - x^2) dx = 1$$

$$\Rightarrow k \left[ \frac{2x^2}{2} - \frac{x^3}{3} \right]_0^2 = 1$$

$$\Rightarrow k \left[ 2^2 - \frac{2^3}{3} \right] = 1$$

$$\Rightarrow k \left( 4 - \frac{8}{3} \right) = 1$$

$$\Rightarrow k = 3/4$$

$$\therefore f(x) = \frac{3}{4}x(2-x)$$

$$\begin{aligned}\mu'_2 &= \int_{-\infty}^{+\infty} x f(x) dx = \int_0^2 x \frac{3}{4}x(2-x) dx \\ &= \frac{3}{4} \int_0^2 (2x^2 - x^3) dx \\ &= \frac{3}{4} \left[ 2 \frac{x^3}{3} - \frac{x^4}{4} \right]_0^2 \\ &= \frac{3}{4} \left[ 2 \times \frac{8}{3} - \frac{16}{4} \right] = \frac{3}{4} \left[ \frac{64 - 48}{12} \right] = 1\end{aligned}$$

$$\begin{aligned}\mu'_2 &= E(x^2) = \int_{-\infty}^{+\infty} x^2 f(x) dx = \int_0^2 x^2 \frac{3}{4}x(2-x) dx \\ &= \frac{3}{4} \int_0^2 (2x^2 - x^3) dx \\ &= \frac{3}{4} \left[ \frac{2x^4}{4} - \frac{x^5}{5} \right]_0^2 \\ &= \frac{3}{4} \left[ \frac{2^4}{2} - \frac{2^5}{5} \right] = \frac{6}{5} = 1.2\end{aligned}$$

$$\begin{aligned}\mu'_3 &= E(x^3) = \int_{-\infty}^{+\infty} x^3 f(x) dx = \int_0^2 x^3 \frac{3}{4}x(2-x) dx \\ &= \frac{3}{4} \int_0^2 (2x^4 - x^5) dx \\ &= \frac{3}{4} \left[ 2 \frac{x^5}{5} - \frac{x^6}{6} \right]_0^2 \\ &= \frac{3}{4} \left[ 2 \times \frac{2^5}{5} - \frac{2^6}{6} \right]\end{aligned}$$

$$= \frac{3}{4} \times \frac{384 - 320}{30} = \frac{3}{4} \times \frac{64}{30}$$

$$\begin{aligned}\mu_4' &= E(x^4) = \int_{-\infty}^{+\infty} x^4 f(x) dx = \int_0^2 x^4 \frac{3}{4} x (2-x) dx \\ &= \frac{3}{4} \int_0^2 (2x^5 - x^6) dx \\ &= \frac{3}{4} \left[ \frac{2x^6}{6} - \frac{x^7}{7} \right]_0^2 \\ &= \frac{3}{4} \left[ 2 \times \frac{2^6}{6} - \frac{2^7}{7} \right] = \frac{3}{4} \left[ \frac{128}{6} - \frac{128}{7} \right] = \frac{3}{4} \times \frac{128}{42} = 2.28\end{aligned}$$

$$\mu_2 = \mu_2' - \mu_1'^2 = 1.2 - 1^2 = 0.2$$

$$\mu_3 = \mu_3' - 3 \mu_2' \mu_1' + 2 \mu_1'^3 = 1.6 - 3 \times 1.2 \times 1 + 2 \times 1^3 = 1.6 - 3.6 + 2 = 0$$

$$\mu_4 = \mu_4' - 3 \mu_3' \mu_1' + 6 \mu_2' \mu_1'^2 - 3 \mu_1'^4 = 2.28 - 4 \times 1.6 \times 1 + 6 \times 1.2 \times 1^2 - 3 \times 1^4 = 0.08$$

### Measures

$$\text{Central tendency} = E(x) = \mu_1' = 1$$

$$\text{Dispersion} = \sigma = \sqrt{V(x)} = \sqrt{\mu_2} = \sqrt{0.2} = 0.44$$

$$\text{Skewness} = \beta_1 = \frac{\mu_3^2}{\mu_2^3} = \frac{0}{(0.2)^3} = 0$$

$$\text{Kurtosis} = \beta_2 = \frac{\mu_4}{\mu_2^2} = \frac{0.08}{(0.2)^2} = 2$$

$$21. f(x) = k x^2 e^{-x} \quad 0 < x < \infty$$

$$\text{Since, } \int_{-\infty}^{+\infty} f(x) dx = 1$$

$$\Rightarrow \int_0^{\infty} kx^2 e^{-x} dx = 1$$

$$\Rightarrow k \int_0^{\infty} x^2 e^{-x} x^{3-1} dx = 1$$

$$\Rightarrow k \sqrt{3} = 1$$

$$\Rightarrow k 2! = 1$$

$$\Rightarrow k = \frac{1}{2}$$

$$\therefore f(x) = \frac{1}{2} x^2 e^{-x} dx$$

Now,

$$\mu_1' = E(x) = \int_{-\infty}^{+\infty} x f(x) dx = \int_{-\infty}^{+\infty} x \frac{1}{2} x^2 e^{-x} dx = \frac{1}{2} \int_0^{\infty} e^{-x} x^3 dx = \frac{3!}{2} = 3$$

$$\mu_2' = E(x^2) = \int_0^{\infty} x^2 e^{-x} dx = \frac{1}{2} \int_0^{\infty} e^{-x} x^4 dx = \frac{4!}{2} = 12$$

$$\mu_3' = E(x^3) = \int_0^{\infty} x^3 f(x) dx = \int_0^{\infty} x^3 \frac{1}{2} x^2 e^{-x} dx = \frac{1}{2} \int_0^{\infty} e^{-x} x^5 dx = \frac{5!}{2} = 60$$

$$\mu_4' = E(x^4) = \int_0^{\infty} x^4 f(x) dx = \int_0^{\infty} x^4 \frac{1}{2} x^2 e^{-x} dx = \frac{1}{2} \int_0^{\infty} e^{-x} x^6 dx = \frac{6!}{2} = 360$$

$$\mu_2 = \mu_2' - \mu_1'^2 = 12 - 3^2 = 3$$

$$\mu_3 = \mu_3' - 3 \mu_2' \mu_1' + 2 \mu_1'^3 = 60 - 2 \times 12 \times 3 + 2 \times 3^3 = 6$$

**54 CHAPTER - 5 | A Complete Solutions of Statistics I for BSc. CSIT**

$$\mu_4 = \mu_4' - 4\mu_3'\mu_1 + 6\mu_2'\mu_1^2 - 3\mu_1^4 = 360 - 4 \times 60 \times 3 + 6 \times 12 \times 3^2 - 3 \times 3^4 = 45$$

$$\text{Mean} = E(x) = \mu_1 = 3$$

$$\text{S.D.} = \sqrt{V(x)} = \sqrt{\mu_2} = \sqrt{3} = 1.73$$

$$\text{Skewness} = \beta_1 = \frac{\mu_3^2}{\mu_2^3} = \frac{6^2}{3^3} = \frac{36}{27} = 1$$

$$\text{Kurtosis} = \beta_2 = \frac{\mu_4}{\mu_2^2} = \frac{45}{3^2} = 5$$

$$22. P(x, y) = \frac{2x + y}{28} \quad x = 1, 3 \\ y = 2, 4$$

$$P(y) = \frac{y+4}{14}$$

$$E(x|y=2) = \sum_x x P(x|y=2)$$

$$= \sum_x x \frac{\frac{2x+2}{28}}{\frac{2+4}{14}}$$

$$= \sum_x x \frac{2x+2}{28} \times \frac{14}{6}$$

$$= \frac{1}{12} \sum_x 2x^2 + 2x = \frac{1}{12} [(2 \times 1^2 + 2 \times 1) + (2 \times 3^2 + 2 \times 3)] = \frac{7}{3}$$

$$E(x^2|y=2) = \sum_x x^2 P(x|y=2)$$

$$= \sum_x x^2 \frac{P(x, y)}{P(y)}$$

$$= \sum_x x^2 \frac{\frac{2x+2}{28}}{\frac{2+4}{14}}$$

$$= \sum_x x^2 \frac{2x+2}{28} \times \frac{14}{6}$$

$$= \frac{1}{12} \sum_x 2x^3 + 2x^2 = \frac{1}{12} [(2 \times 1^3 + 2 \times 1^2) + (2 \times 3^3 + 2 \times 3^2)]$$

$$= \frac{1}{12} [4 + 54 + 18] = \frac{76}{12}$$

$$V(x|y=2) = E(x^2|y=2) - [E(x|y=2)]^2$$

$$= \frac{76}{12} - \left(\frac{7}{3}\right)^2 = \frac{8}{9}$$

## Chapter 6

1.  $f(x, y) = \frac{6}{5} (x + y^2) \quad 0 < x < 1, 0 < y < 1$

(i)  $P(0.2 < x < 0.5, 0.4 < y < 0.6)$

$$\begin{aligned}
 &= \int_{0.2}^{0.5} \left\{ \int_{0.4}^{0.6} f(x, y) dy \right\} dx \\
 &= \int_{0.2}^{0.5} \left\{ \int_{0.4}^{0.6} \frac{6}{5} (x + y^2) dy \right\} dx \\
 &= \frac{6}{5} \int_{0.2}^{0.5} \left[ xy + \frac{y^3}{3} \right]_{0.4}^{0.6} dx \\
 &= \frac{6}{5} \int_{0.2}^{0.5} \left[ 0.6x + \frac{(0.6)^3}{3} - 0.4x - \frac{(0.4)^3}{3} \right] dx \\
 &= \frac{6}{5} \int_{0.2}^{0.5} \left( 0.2x + \frac{0.152}{3} \right) dx \\
 &= \frac{6}{5} \int_{0.2}^{0.5} \left[ 0.2 \frac{x^2}{2} + \frac{0.152}{3} x \right]_{0.2}^{0.5} \\
 &= \frac{6}{5} \left[ \frac{0.2}{2} \times (0.5)^2 + \frac{0.152}{3} \times 0.5 - \frac{0.2}{2} \times (0.2)^2 - \frac{0.152}{3} \times (0.2) \right] \\
 &= \frac{6}{5} [0.025 + 0.0253 - 0.004 - 0.0103] = 0.043
 \end{aligned}$$

$$\begin{aligned}
 P(x > 0.4, y < 0.5) &= \int_{0.4}^1 \left[ \int_0^{0.5} f(x, y) \right] dy dx \\
 &= \int_{0.4}^1 \left[ \int_0^{0.5} \frac{6}{5} (x + y^2) dy \right] dx \\
 &= \frac{6}{5} \int_{0.4}^1 \left[ xy + \frac{y^3}{3} \right]_0^{0.5} dx \\
 &= \frac{6}{5} \int_{0.4}^1 \left[ 0.5x + \frac{(0.5)^3}{3} \right] dx \\
 &= \frac{6}{5} \int_{0.4}^1 \left[ 0.5x + \frac{0.125}{3} \right] dx \\
 &= \frac{6}{5} \left[ 0.5 \frac{x^2}{2} + \frac{0.125}{3} x \right]_{0.4}^1 \\
 &= \frac{6}{5} \left[ \frac{0.5}{2} (1)^2 + \frac{0.125}{3} (1) - \frac{0.5}{2} (0.4)^2 - \frac{0.125}{3} (0.4) \right] = 0.28
 \end{aligned}$$

### 2. Solution.

$$f(x, y) = \begin{cases} x + y & 0 < x, y < 1 \\ 0 & \text{Otherwise} \end{cases}$$

$$f(x) = \int_0^1 f(x, y) dy = \int_0^1 (x + y) dy = \left[ xy + \frac{y^2}{2} \right]_0^1 = x + \frac{1}{2}$$

$$f(y) = \int_0^1 f(x, y) dx = \int_0^1 (x + y) dx = \left[ \frac{x^2}{2} + xy \right]_0^1 = \frac{1}{2} + y$$

$$E(xy) = \int_0^1 \int_0^1 xu f(x, y) dx dy$$

$$= \int_0^1 x \left[ \int_0^1 xy f(x, y) dy \right]$$

$$= \int_0^1 x \left[ x \frac{y^2}{2} + \frac{y^3}{3} \right]_0^1 dx$$

$$= \int_0^1 x \left( \frac{x}{2} + \frac{1}{3} \right) dx$$

$$= \int_0^1 \left( \frac{x^2}{2} + \frac{x}{3} \right) dx$$

$$= \left[ \frac{1}{2} \frac{x^3}{3} + \frac{1}{3} \frac{x^2}{2} \right]_0^1 = \left[ \frac{x^3}{6} + \frac{x^2}{6} \right]_0^1 = \frac{1}{6} + \frac{1}{6} = \frac{1}{3}$$

$$E(x) = \int_0^1 x f(x) dx = \int_0^1 x \left( x + \frac{1}{2} \right) dx = \left[ \frac{x^3}{3} + \frac{1}{2}x^2 - \frac{x^2}{2} \right]_0^1 = \frac{1}{3} + \frac{1}{4} = \frac{7}{12}$$

$$E(y) = \int_0^1 y f(y) dy = \frac{7}{12}$$

$$\text{Cov}(x, y) = E(xy) - E(x) E(y) = \frac{1}{3} - \frac{7}{12} \times \frac{7}{12} = -0.0069$$

### 3. Solution.

$$f(x_1, x_2) = \frac{2}{3} (x_1 + 2x_2) \quad 0 < x_1 < 1, 0 < x_2 < 1$$

$$f(x_1) = \int_0^1 f(x_1, x_2) dx_2 = \int_0^1 \frac{2}{3} (x_1 + 2x_2) dx_2 = \frac{2}{3} \left[ x_1 x_2 + \frac{2x_2^2}{2} \right]_0^1 = \frac{2}{3} (x_1 + 1)$$

$$f(x_2) = \int_0^1 f(x_1, x_2) dx_1 = \int_0^1 \frac{2}{3} (x_1 + 2x_2) dx_1 = \frac{2}{3} \left[ \frac{x_1^2}{2} + 2x_1 x_2 \right]_0^1 = \frac{1 + 4x_2}{3}$$

$$f(x_1/x_2) = \frac{f(x_1, x_2)}{f(x_2)} = \frac{\frac{2}{3} (x_1 + 2x_2)}{\frac{1 + 4x_2}{3}} = \frac{2 (x_1 + 2x_2)}{1 + 4x_2}$$

$$f(x_2/x_1) = \frac{f(x_1, x_2)}{f(x_1)} = \frac{\frac{2}{3} (x_1 + 2x_2)}{\frac{2}{3} (x_1 + 1)} = \frac{x_1 + 2x_2}{x_1 + 1}$$

### 4. Solution.

$$P(x, y) = \frac{x - y + 3}{48}, \quad x = 0, 1, 2, 3 \\ y = 0, 1, 2, 3$$

$$P(x) = \sum_y P(x, y) = \sum_{y=0}^3 \frac{x - y + 3}{48} = \frac{x - 0 + 3}{48} + \frac{x - 1 + 3}{48} + \frac{x - 2 + 3}{48} + \frac{x - 3 + 3}{48}$$

$$= \frac{4x - 6 + 12}{48} = \frac{4x + 6}{48} = \frac{2x + 3}{24}$$

$$P(y) = \sum_x P(x, y) = \sum_{x=0}^3 \frac{x - y + 3}{48}$$

$$= \frac{0 - y + 3}{48} + \frac{1 - y + 3}{48} + \frac{2 - y + 3}{48} + \frac{3 - y + 3}{48} = \frac{9 - 2y}{24}$$

$$P(x/y = 1) = \frac{P(x, y = 1)}{P(y = 1)} = \frac{\frac{x - 1 + 3}{48}}{\frac{9 - 2}{24}} = \frac{x + 2}{14}$$

$$P(y/x = 1) = \frac{P(x = 2, y)}{P(x = x)} = \frac{\frac{2 - y + 3}{48}}{\frac{4 + 3}{24}} = \frac{5 - y}{14}$$

5. Solution.

	x	-1	0	1
y				
0		1/15	2/15	1/15
1		3/15	2/15	1/15
2		2/15	1/15	2/15

$$(i) P(x = 1, y \leq 1) = P(x > 1, y = 0) + P(x = 1, y = 1) = \frac{1}{15} + \frac{1}{15} = \frac{2}{15}$$

$$(ii) P(y \leq 1) = P(y = 0) + P(y = 1) = \left(\frac{1}{15} + \frac{2}{15} + \frac{1}{15}\right) + \left(\frac{3}{15} + \frac{2}{15} + \frac{1}{15}\right) = \frac{4}{15} + \frac{6}{15} = \frac{2}{3}$$

$$(iii) P(x = -1) = \frac{1}{15} + \frac{3}{15} + \frac{2}{15} = \frac{2}{15}$$

6. Solution.

Let hard ware = H and soft ware = S

$$\text{Total} = 3H + 4S = 7$$

$$\text{Selected} = 2$$

Here, x = Prob. of Hardware, y = Prob. of Software

$$P(x = 0, y = 0) = 0$$

$$P(x = 0, y = 1) = 0$$

$$P(x = 0, y = 2) = \frac{^4C_2}{^7C_2} = \frac{6}{21}$$

$$P(x = 1, y = 0) = 0$$

$$P(x = 1, y = 1) = \frac{^3C_1 \times ^4C_1}{^7C_2} = \frac{12}{21}$$

$$P(x = 1, y = 2) = 0$$

$$P(x = 2, y = 0) = \frac{^3C_2}{^7C_2} = \frac{3}{21}$$

$$P(x = 2, y = 1) = 0$$

$$P(x = 2, y = 2) = 0$$

x \ y	0	1	2	P(4)
x	0	1	2	
y	0	0	3/21	3/21
0	0	0	0	12/21
1	0	12/21	0	12/21
2	6/21	0	0	6/21

$$P(x = 0/y = 0) = 0, P(x = 0/y = 1) = 0$$

$$P(x = 0/y = 2) = \frac{P(x = 0, y = 2)}{P(y = 2)} = \frac{6/21}{6/21} = 1$$

$$P(x = 1/y = 0) = 0; P(x = 1/y = 1) = \frac{P(x = 1, y = 1)}{P(y = 1)} = \frac{\frac{12}{21}}{\frac{21}{21}} = 1$$

$$P(x = 1/y = 2) = 0$$

$$P(x = 2/y = 0) = \frac{P(x = 2, y = 0)}{P(y = 0)} = \frac{3/21}{3/21} = 1$$

$$P(y = 2/y = 1) = 0 \quad P(x = 2/y = 2) = 0$$

7. Solution.

x \ y	0	1	2	P(4)
x	0	1	2	
y	0.03	0.02	0.05	0.1
-1	0.01	0.25	0.45	0.72
0	0	0.03	0.05	0.18
1				

$$P(x = 0/y = -1) = \frac{0.03}{0.1} = 0.3$$

$$P(x = 0/y = 0) = \frac{0.01}{0.72} = 0.013$$

$$P(x = 0/y = 1) = \frac{0}{0.18} = 0$$

$$P(x = 1/y = -1) = \frac{0.02}{0.1} = 0.2$$

$$P(x = 1/y = 0) = \frac{0.25}{0.72} = 0.347$$

$$P(x = 1/y = 1) = \frac{0.03}{0.18} = 0.166$$

$$P(x = 2/y = -1) = \frac{0.05}{0.1} = 0.5$$

$$P(x = 2/y = 0) = \frac{0.45}{0.72} = 0.625$$

$$P(x = 2/y = 1) = \frac{0.05}{0.18} = 0.277$$

$$P(x = 3/y = -1) = \frac{0}{0.1} = 0$$

$$P(x = 3/y = 0) = \frac{0.01}{0.72} = 0.013$$

$$P(x = 3/y = 1) = \frac{0.1}{0.18} = 0.555$$

$$P(y = -1/x = 0) = \frac{0.03}{0.04} = 0.75$$

$$P(y = -1/x = 1) = \frac{0.02}{0.3} = 0.067$$

$$P(y = -1/x = 2) = \frac{0.05}{0.55} = 0.109$$

$$P(y = -1/x = 3) = \frac{0}{0.11} = 0$$

$$P(y = 0/x = 0) = \frac{0.01}{0.04} = 0.25$$

$$P(y = 0/x = 1) = \frac{0.25}{0.3} = 0.833$$

$$P(y = 0/x = 2) = \frac{0.45}{0.55} = 0.818$$

$$P(y = 0/x = 3) = \frac{0.01}{0.11} = 0.09$$

$$P(y = 1/x = 0) = \frac{0}{0.04} = 0$$

$$P(y = 1/x = 1) = \frac{0.03}{0.3} = 0.1$$

$$P(y = 1/x = 2) = \frac{0.05}{0.55} = 0.09$$

$$P(y = 1/x = 3) = \frac{0.1}{0.11} = 0.909$$

$$\begin{aligned} P(x = 0, y \leq 1) &= P(x = 0, y = -1) + P(x = 0, y = 0) + P(x = 0, y = 1) \\ &= 0.03 + 0.01 + 0 \\ &= 0.04 \end{aligned}$$

$$P(x \geq 0, y < 1) = P(y = 1) + P(y = 0) = 0.1 + 0.72 = 0.82$$

$$P(x \leq 0/y = 1) = \frac{P(x \leq 0, y = 1)}{P(y = 1)} = \frac{0}{0.18} = 0$$

### 8. Solution.

$$f(x, y) = \begin{cases} x^2 + kxy & 0 \leq x \leq 1 \\ 0 & \text{elsewhere} \end{cases}$$

$$= \int_{-\infty}^{+\infty} f(x, y) dy = \int_0^1 2x dy = [2xy]_0^1 = 2x$$

$$f(y) = \int_{-\infty}^{+\infty} f(x, y) dx = \int_0^1 2x dx = \left[ 2 \frac{x^2}{2} \right]_0^1 = 1$$

Here,  $f(x, y) = 2x$

$$f(x) f(y) = 2x \cdot 1 = 2x$$

$$f(x, y) = f(x) f(y)$$

Hence, x and y are independent.

9.  $f(x, y) = 2x$

$$f(x) = \int_0^1 f(x, y) dy = \int_0^1 2x dy = 2x [y]_0^1 = 2x$$

$$f(y) = \int_0^1 f(x, y) dx = \int_0^1 2x dx = 2 \left[ \frac{x^2}{2} \right]_0^1 = 1$$

$$f(x) f(y) = 2x \cdot 1 = 2x$$

Hence, x and y are independent.

10.  $f(x, y) = \begin{cases} k(6-x-y) & 0 \leq x \leq 2, 0 \leq y \leq 4 \\ 0 & \text{elsewhere} \end{cases}$

Since,  $\int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} f(x, y) dy dx = 1$

$$\Rightarrow \int_0^2 \left[ \int_2^4 k(6-x-y) dy \right] dx = 1$$

$$\Rightarrow k \int_0^2 \left[ 6y - xy - \frac{y^2}{2} \right]_2^4 dx = 1$$

$$\Rightarrow k \int_0^2 \left( 6 \times 4 - 4x - \frac{16}{2} - 6 \times 2 + 2x + \frac{4}{2} \right) dx = 1$$

$$\Rightarrow k \int_0^2 (6 - 2x) dx = 1$$

$$\Rightarrow k \left[ 6x - \frac{2x^2}{2} \right]_0^2 = 1$$

$$\Rightarrow k(12 - 4) = 1$$

$$\Rightarrow k = \frac{1}{8}$$

$$\therefore f(x, y) = \frac{6-x-y}{8}$$

$$P(x \leq 1 \cap y < 3) = \int_0^1 \left[ \int_2^3 \frac{6-x-y}{8} dy \right] dx$$

$$= \frac{1}{8} \int_0^1 \left( 18 - 3x - \frac{9}{2} - 12 + 2x + \frac{4}{2} \right) dx$$

$$= \frac{1}{8} \int_0^1 (3.5 - x) dx = \frac{1}{8} \left[ 3.5x - \frac{x^2}{2} \right]_0^1 = \frac{1}{8} \left( 3.5 \times 1 - \frac{1}{2} \right) = \frac{3}{8}$$

$$f(y) = \int_{-\infty}^{\infty} f(x, y) dx = \int_0^2 \frac{6-x-y}{8} dx = \frac{1}{8} \left[ 6x - \frac{x^2}{2} - xy \right]_0^2 \\ = \frac{1}{8} [12 - 2 - 2y] = \frac{5-y}{4}$$

$$P(y \leq 3) = \int_2^3 f(y) dy = \int_2^3 \frac{5-y}{4} dy = \frac{1}{4} \left[ 5y - \frac{y^2}{2} \right]_2^3 = \frac{1}{4} \left[ 15 - \frac{9}{2} - 10 + \frac{4}{2} \right] = \frac{5}{8}$$

$$P(x \leq 1/y \leq 3) = \frac{P(x \leq 1, y \leq 3)}{P(y \leq 3)} = \frac{\int_0^1 \left[ \int_2^3 f(x, y) dy \right] dx}{\int_2^3 f(y) dy} = \frac{3/8}{5/8} = \frac{3}{5}$$

11.  $f(x, y) = 4xy e^{-(x^2+y^2)}$   $x \geq 0, y \geq 0$

$$f(x) = \int_0^{\infty} f(x, y) dy = \int_0^{\infty} 4xy e^{-(x^2+y^2)} dy = 4x e^{-x^2} \int_0^{\infty} y e^{-y^2} dy$$

Let  $y^2 = z$ ;

$$2y dy = dz$$

When  $y = 0, z = 0$  and when  $y = \infty, z = \infty$

$$= 4x e^{-x^2} \int_0^{\infty} e^{-z} \frac{dz}{z} \\ = \frac{4x e^{-x^2}}{2} \int_0^{\infty} e^{-z} z^{1-1} dz \\ = z x e^{-x^2}$$

$$f(y) = \int_0^{\infty} f(x, y) dx = 2y e^{-y^2}$$

$$f(x/y) = \frac{f(x, y)}{f(y)} = \frac{4xy e^{-(x^2+y^2)}}{2y e^{-y^2}} = 2x e^{-x^2}$$

$$\text{Here, } f(x) f(y) = 2x e^{-x^2} 2y e^{-y^2} = 4xy e^{-(x^2+y^2)} = f(x, y)$$

Hence,  $x$  and  $y$  are independent.

12.  $f(x, y) = k(2x + 3y)$   $0 \leq x \leq 1, 0 \leq y \leq 1$

Since,

$$\int_{-\infty}^{\infty} \left[ \int_{-\infty}^{\infty} f(x, y) dy \right] dx = 1$$

$$\Rightarrow \int_0^1 \left[ \int_0^1 k(2x + 3y) dy \right] dx = 1$$

$$\Rightarrow k \int_0^1 \left[ 2xy + \frac{3y^2}{2} \right]_0^1 dx = 1$$

$$\Rightarrow k \int_0^1 \left( 2x + \frac{3}{2} \right) dx = 1$$

$$\Rightarrow k \left[ \frac{2x^2}{2} + \frac{3}{2} x \right]_0^1 = 1$$

$$\Rightarrow k \left[ 1 + \frac{3}{2} \right] = 1$$

$$\Rightarrow k = \frac{2}{5} \quad \therefore f(x, y) = \frac{2}{5}(2x + 3y)$$

$$f(x) = \int_{-\infty}^{+\infty} f(x, y) dy = \int_0^1 \frac{2}{5}(2x + 3y) dy \\ = \frac{2}{5} \left[ 2xy + \frac{3y^2}{2} \right]_0^1 = \frac{2}{5} \left[ 2x + \frac{3}{2} \right] = \frac{4x + 3}{5}$$

$$f(y) = \int_{-\infty}^{+\infty} f(x, y) dx = \int_0^1 \frac{2}{5}(2x + 3y) dx \\ = \frac{2}{5} \left[ \frac{2x^2}{2} + 3xy \right]_0^1 = \frac{2}{5}[1 + 3y]$$

$$f(x/y) = \frac{f(x, y)}{f(y)} = \frac{\frac{2}{5}(2x + 3y)}{\frac{2}{5}(1 + 3y)} = \frac{2x + 3y}{1 + 3y}$$

$$f(y/x) = \frac{f(x, y)}{f(x)} = \frac{\frac{2}{5}(2x + 3y)}{\frac{4x + 3}{5}} = \frac{4x + 6y}{4x + 3}$$

$$f(x) f(y) = \frac{f(x, y)}{f(x)} = \frac{4x + 3}{5} \times \frac{2}{5}(1 + 3y) = \frac{2}{25}(4x + 12xy + 3 + 9y) \neq f(x, y)$$

Hence, x and y are dependent.

$$E(x) = \int_0^1 x f(x) dx = \int_0^1 x \frac{(4x + 3)}{5} dx \\ = \frac{1}{5} \int_0^1 (4x^2 + 3x) dx \\ = \frac{1}{5} \left[ \frac{4x^3}{3} + \frac{3x^2}{2} \right]_0^1 = \frac{1}{5} \left[ \frac{4}{3} + \frac{3}{2} \right] = \frac{17}{30}$$

$$E(x^2) = \int_0^1 x^2 f(x) dx = \int_0^1 x^2 \frac{4x + 3}{5} dx \\ = \frac{1}{5} \int_0^1 (4x^3 + 3x^2) dx = \frac{1}{5} \left[ \frac{4x^4}{4} + \frac{3x^3}{3} \right]_0^1 = \frac{1}{5}[1+1] = \frac{2}{5}$$

$$E(y) = \int_0^1 y f(y) dy = \int_0^1 y \frac{2}{5}(1 + 3y) dy \\ = \frac{2}{5} \int_0^1 (y + 3y^2) dy \\ = \frac{2}{5} \left[ \frac{y^2}{2} + \frac{3y^3}{3} \right]_0^1 = \frac{2}{5} \left[ \frac{1}{2} + 1 \right] = \frac{3}{5}$$

$$E(y^2) = \int_0^1 y^2 f(y) dy = \int_0^1 y^2 \frac{2}{5}(1 + 3y) dy \\ = \frac{2}{5} \int_0^1 (y^2 + 3y^3) dy = \frac{2}{5} \left[ \frac{y^3}{3} + \frac{3y^4}{4} \right]_0^1 = \frac{2}{5} \left[ \frac{1}{3} + \frac{3}{4} \right] = \frac{13}{30}$$

$$\begin{aligned}
 E(x_4) &= \int_0^1 \left[ \int_0^1 xy f(x, y) dy \right] dx = \int_0^1 \left[ \int_0^1 xy \frac{2}{5} (2x + 3y) dy \right] dx \\
 &= \frac{2}{5} \int_0^1 \left[ \int_0^1 (2x^2 y + 3xy^2) dy \right] dx \\
 &= \frac{2}{5} \int_0^1 \left[ \frac{2x^2 y^2}{2} + \frac{3xy^3}{3} \right]_0^1 dx \\
 &= \frac{2}{5} \int_0^1 (x^2 + x) dx = \frac{2}{5} \left[ \frac{x^3}{3} + \frac{x^2}{2} \right]_0^1 = \frac{2}{5} \left[ \frac{1}{3} + \frac{1}{2} \right] = \frac{1}{3}
 \end{aligned}$$

$$\begin{aligned}
 E(x^2y^2) &= \int_0^1 \left[ \int_0^1 x^2y^2 f(x, y) dy \right] dx = \int_0^1 \left[ \int_0^1 x^2y^2 \frac{2}{5} (2x + 3y) dy \right] dx \\
 &= \frac{2}{5} \int_0^1 \left[ \int_0^1 (2x^3 y^2 + 3x^2 y^3) dy \right] dx \\
 &= \frac{2}{5} \int_0^1 \left[ \frac{2x^3 y^3}{3} + \frac{3x^2 y^4}{4} \right]_0^1 dx \\
 &= \frac{2}{5} \int_0^1 \left( \frac{2}{3} x^3 + \frac{3}{4} x^2 \right) dx \\
 &= \frac{2}{5} \left[ \frac{2}{3} \frac{x^4}{4} + \frac{3}{4} \frac{x^3}{3} \right]_0^1 = \frac{2}{5} \left[ \frac{1}{6} + \frac{1}{4} \right] = \frac{1}{6}
 \end{aligned}$$

$$V(x) = E(x^2) = [E(x)]^2 = \frac{2}{5} - \left(\frac{17}{30}\right)^2 = 0.4 - (0.566)^2 = 0.079$$

$$V(y) = E(y^2) - [E(y)]^2 = \frac{1}{6} - \left(\frac{1}{3}\right)^2 = \frac{1}{6} - \frac{1}{9} = 0.055$$

$$\text{Cov}(xy) = E(xy) - E(x)E(y) = \frac{1}{3} - \frac{17}{30} \times \frac{3}{5} = -\frac{1}{50}$$

$$r(x, y) = \frac{\text{Cov}(x, y)}{\sqrt{V(x)} \sqrt{V(y)}} = \frac{-\frac{1}{50}}{\sqrt{0.079} \sqrt{0.073}} = -0.087$$

$$E(2x + 3y) = 2E(x) + 3E(y) = 2 \times \frac{17}{30} + 3 \times \frac{3}{5} = 2.9$$

$$\begin{aligned}
 V(3x - 2y) &= 9V(x) + 4V(y) - 12\text{Cov}(xy) \\
 &= 9 \times 0.079 + 4 \times 0.073 - 12(-0.0066) = 1.08
 \end{aligned}$$

$$13. P(x, y) = k(x + y), \begin{matrix} x \leq 1, 2, 3 \\ y = 1, 2 \end{matrix}$$

Since,

$$\sum_x \sum_y P(x, y) = 1$$

$$\Rightarrow \sum_{x=1}^3 \sum_{y=1}^2 k(x + y) = 1$$

$$\Rightarrow k [(1+1) + (1+2) + (2+1) + (2+2) + (3+1) + (3+2)] = 1$$

$$\Rightarrow k = \frac{1}{21}$$

$$\therefore P(x, y) = \frac{1}{21} P(x, y)$$

$$P(x) = \sum_y P(x, y)$$

$$= \sum_{y=1}^2 \frac{1}{21} (x+y) = \frac{1}{21} [(x+1) + (x+2)] = \frac{2x+3}{21}$$

$$P(y) = \sum_x P(x, y)$$

$$= \sum_{x=1}^2 \frac{1}{21} (x+y) = \frac{1}{21} (1+y+2+y+3+y) = \frac{6+3y}{21}$$

$$P(x, y) = \frac{P(x, y)}{P(y)} = \frac{\frac{1}{21}(x+y)}{\frac{6+3y}{21}} = \frac{x+y}{6+3y} = \frac{x+y}{3y+6}$$

$$P(y/x) = \frac{P(x, y)}{P(x)} = \frac{\frac{1}{21}(x+y)}{\frac{2x+3}{21}} = \frac{x+y}{2x+3}$$

$$P(x, y) = \frac{1}{21} (x+y)$$

$$P(x) P(y) = \frac{2x+3}{21} \times \frac{3y+6}{21} = \frac{6xy+12x+9y+18}{441}$$

$$P(x, y) = P(x), P(y)$$

Hence, x and y are dependent

$$E(x) = \sum_{x=1}^3 2x^2 + 3x = \frac{1}{21} [(2 \times 1^2 + 3 \times 1) + (2 \times 2^2 + 3 \times 2) + (2 \times 3^2 + 3 \times 3)] = \frac{46}{21}$$

$$E(x^2) = \sum_x x^2 P(x) = \sum_{x=1}^3 x^2 \frac{2x+3}{21}$$

$$= \frac{1}{21} \sum_{x=1}^3 (2x^3 + 3x^2)$$

$$= \frac{1}{21} [(2 \times 1^3 + 3 \times 1^2) + (2 \times 2^3 + 3 \times 2^2) + (2 \times 3^3 + 3 \times 3^2)]$$

$$= \frac{114}{21}$$

$$\begin{aligned}
 E(y) &= \sum_x y P(y) = \sum_{y=1}^2 y \frac{6+3y}{21} \\
 &= \frac{1}{21} \sum_{y=1}^2 6y + 3y^2 \\
 &= \frac{1}{21} [(6 \times 1 + 3 \times 1^2) + (6 \times 2 \times 3 \times 2^2)] = \frac{33}{21}
 \end{aligned}$$

$$\begin{aligned}
 E(y^2) &= \sum_y y^2 P(y) = \sum_{y=1}^2 y^2 \frac{(6+3y)}{21} \\
 &= \frac{1}{21} \sum_{y=1}^2 6y^2 + 3y^3 \\
 &= \frac{1}{21} [(6 \times 1^2 + 3 \times 1^3) + (6 \times 2^2 + 3 \times 2^3)] = \frac{57}{21}
 \end{aligned}$$

$$\begin{aligned}
 E(xy) &= \sum_x \sum_y xy P(x, y) = \sum_{x=1}^3 \sum_{y=1}^2 xy \frac{x+y}{21} \\
 &= \frac{1}{21} \sum_{x=1}^3 \sum_{y=1}^2 x^2 y + xy^2 \\
 &= \frac{1}{21} \sum_{x=1}^3 (x^2 \times 1 + x \times 1^2) + (x^2 \times 2 + x \times 2^2) \\
 &= \frac{1}{21} \sum_{x=1}^3 (x^2 + x + 2x^2 + 4x) \\
 &= \frac{1}{21} \sum_{x=1}^3 (3x^2 + 5x) \\
 &= \frac{1}{21} [(3 \times 1^2 + 5 \times 1) + (3 \times 2^2 + 5 \times 2) + (3 \times 3^2 + 5 \times 3)] \\
 &= \frac{306}{21}
 \end{aligned}$$

$$V(x) = E(x^2) - [E(x)]^2 = \frac{114}{21} - \left(\frac{46}{21}\right)^2 = 5.42 - 4.79 = 0.63$$

$$V(y) = E(y^2) - [E(y)]^2 = \frac{57}{21} - \left(\frac{33}{21}\right)^2 = 2.71 - 2.46 = 0.25$$

$$V(xy) = E(x^2y^2) - [E(xy)]^2 = \frac{306}{21} - \left(\frac{72}{21}\right)^2 = 14.57 - 11.75 = 2.82$$

$$\text{Cov}(xy) = E(xy) - E(x) E(y) = \frac{72}{21} - \frac{46}{21} \times \frac{33}{21} = 3.428 - 3.442 = -0.017$$

$$r = \frac{\text{Cov}(x, y)}{\sqrt{V(x)} \sqrt{V(y)}} = \frac{-0.17}{\sqrt{0.63} \sqrt{0.25}} = -0.043$$

$$E(2x - 3y) = 2 E(x) - 3 E(y) = 2 \times \frac{46}{21} - 3 \times \frac{33}{21} = \frac{92 - 99}{21} = -0.33$$

$$\begin{aligned} V(2x + 3y) &= 4 V(x) + 9 V(y) + 12 \text{Cov}(x, y) \\ &= 4 \times 0.63 + 9 \times 0.25 + 12 \times (-0.017) = 4.566 \end{aligned}$$

14.  $f(x, y) = \begin{cases} k e^{-(x+y)} & 0 \leq x \leq \infty, 0 \leq y \leq \infty \\ 0 & \text{otherwise} \end{cases}$

Since,  $\int_{-\infty}^{+\infty} \left[ \int_{-\infty}^{\infty} f(x, y) dy \right] dx = 1$

$$\Rightarrow \int_0^{\infty} \left[ \int_0^{\infty} k e^{-(x+y)} dy \right] dx = 1$$

$$\Rightarrow k \int_0^{\infty} e^{-x} \left[ \int_0^{\infty} e^{-y} dy \right] dx = 1$$

$$\Rightarrow k \int_0^{\infty} e^{-x} \cdot 1 dx = 1$$

$$\Rightarrow k \cdot 1 = 1$$

$$\Rightarrow k = 1 \quad \therefore f(x, y) = e^{-(x+y)}$$

$$f(x) = \int_{-\infty}^{+\infty} f(x, y) dy$$

$$= \int_0^{\infty} e^{-(x+y)} dy$$

$$= \int_0^{\infty} e^{-x} e^{-y} dy$$

$$= e^{-x} \int_0^{\infty} e^{-y} dy = e^{-x} \times 1 = e^{-x}$$

$$f(y) = \int_{-\infty}^{+\infty} f(x, y) dx$$

$$= \int_0^{\infty} e^{-(x+y)} dx$$

$$= e^{-y} \int_0^{\infty} e^{-x} dx = e^{-y} \cdot 1 = e^{-y}$$

$$f(x/y) = \frac{f(x, y)}{f(y)} = \frac{e^{-(x+y)}}{e^{-y}} = e^{-x}$$

$$f(y/x) = \frac{f(x, y)}{f(x)} = \frac{e^{-(x+y)}}{e^{-x}} = e^{-y}$$

$$\text{Here, } f(x, y) = e^{-(x+y)}$$

$$f(x) f(y) = e^{-x} e^{-y} = e^{-(x+y)}$$

$$f(x, y) = f(x) f(y)$$

Hence, x and y are independent

$$E(x) = \int_{-\infty}^{+\infty} x f(x) dx = \int_0^{\infty} x e^{-x} dx = 1$$

$$E(x^2) = \int_{-\infty}^{+\infty} x^2 f(x) dx = \int_0^{\infty} x^2 e^{-x} dx = 2$$

$$E(y) = \int_0^{\infty} y e^{-y} dy = 1$$

$$E(y^2) = \int_0^{\infty} y^2 e^{-y} dy = 2$$

$$\begin{aligned} E(xy) &= \int_{-\infty}^{\infty} \left[ \int_{-\infty}^{\infty} xy f(x, y) dy \right] dx \\ &= \int_0^{\infty} \left[ \int_{-\infty}^{\infty} xy e^{-(x+y)} dy \right] dx \\ &= \int_0^{\infty} x e^{-x} \left[ \int_0^{\infty} y e^{-y} dy \right] dx \\ &= \int_0^{\infty} x e^{-x} \cdot 1 dx \\ &= 1 \end{aligned}$$

$$\begin{aligned} E(x^2 y^2) &= \int_0^{\infty} \left[ \int_0^{\infty} x^2 y^2 f(x, y) dy \right] dx \\ &= \int_0^{\infty} \left[ \int_0^{\infty} x^2 y^2 e^{-(x+y)} dy \right] dx \\ &= \int_0^{\infty} x^2 e^{-x} \left[ \int_0^{\infty} y^2 e^{-y} dy \right] dx \\ &= \int_0^{\infty} x^2 e^{-x} \cdot 2 dx = 2 \int_0^{\infty} e^{-x} x^2 dx = 2 \times 2 = 4 \end{aligned}$$

$$V(x) = E(x^2) - [E(x)]^2 = 2 - 1^2 = 1$$

$$V(y) = E(y^2) - [E(y)]^2 = 2 - 1^2 = 1$$

$$V(xy) = E(x^2 y^2) - [E(xy)]^2 = 4 - 1^2 = 3$$

$$\text{Cov}(xy) = E(xy) - E(x) E(y) = 1 - 1 \cdot 1 = 0$$

$$r(x, y) = \frac{\text{Cov}(x, y)}{\sqrt{V(x)} \sqrt{V(y)}} = \frac{0}{\sqrt{1} \sqrt{1}} = 0$$

$$E(3x - 2y) = 3 E(x) - 2E(y) = 3 \times 1 - 2 \times 1 = 1$$

$$V(3x + 2y) = 9 V(x) + 4 V(y) + 12 \text{Cov}(xy) = 9 \times 1 + 4 \times 1 + 12 \times 0 = 13$$

15.  $f(x, y) = \begin{cases} k(x+2)e^{-y} & 0 \leq x \leq 1, y \geq 0 \\ 0 & \text{elsewhere} \end{cases}$

Since,  $\int_{-\infty}^{+\infty} \left[ \int_{-\infty}^{\infty} f(x, y) dy \right] dx = 1$

$$\Rightarrow \int_0^1 \left[ \int_0^{\infty} k(x+2)e^{-y} dy \right] dx = 1$$

$$\Rightarrow k \int_0^1 (x+2) \left[ \int_0^{\infty} e^{-y} dy \right] dx = 1$$

$$\Rightarrow k \int_0^1 (x+2) 1 dx = 1$$

$$\Rightarrow k \left[ \frac{x^2}{2} + 2x \right]_0^1 = 1$$

$$\Rightarrow k \left( \frac{1}{2} + 2 \right) = 1$$

$$\Rightarrow k = \frac{2}{5} \quad \therefore f(x, y) = \frac{2}{5} (x + 2) e^{-y}$$

$$f(x) = \int_{-\infty}^{\infty} f(x, y) dy$$

$$= \int_0^{\infty} \frac{2}{5} (x + 2) e^{-y} dy$$

$$= \frac{2}{5} (x + 2) \int_0^{\infty} e^{-y} dy$$

$$= \frac{2}{5} (x + 2)$$

$$f(y) = \int_{-\infty}^{\infty} f(x, y) dx$$

$$= \int_0^1 \frac{2}{5} (x + 2) e^{-y} dx$$

$$= \frac{2}{5} e^{-y} \left[ \frac{x^2}{2} + 2x \right]_0^1$$

$$= \frac{2}{5} e^{-y} \left[ \frac{1}{2} + 2 \right] = \frac{2}{5} e^{-y} + \frac{5}{2} = e^{-y}$$

$$f(x/y) = \frac{f(x, y)}{f(y)} = \frac{\frac{2}{5} (x + 2) e^{-y}}{e^{-y}}$$

$$f(y/x) = \frac{f(x, y)}{f(x)} = \frac{\frac{2}{5} (x + 2) e^{-y}}{\frac{2}{5} (x + 2)} = e^{-y}$$

Here,  $f(x, y) = \frac{2}{5} (x + 2) e^{-y}$

$$f(x) f(y) = \frac{2}{5} (x + 2) e^{-y}$$

Hence,  $x$  and  $y$  are independent.

$$E(x) = \int_{-\infty}^{\infty} x f(x) dx$$

$$= \int_0^1 x \frac{2}{5} (x + 2) dx$$

$$= \frac{2}{5} \left[ \frac{x^3}{3} + \frac{2x^2}{2} \right]_0^1$$

$$= \frac{2}{5} \left[ \frac{1}{3} + 1 \right] = \frac{2}{5} \times \frac{4}{3} = \frac{8}{15}$$

$$\begin{aligned}
 E(x^2) &= \int_{-\infty}^{\infty} x^2 f(x) dx \\
 &= \int_0^1 x^2 \frac{2}{5} (x+2) dx \\
 &= \frac{2}{5} \int_0^1 (x^3 + 2x^2) dx \\
 &= \frac{2}{5} \left[ \frac{x^4}{4} + \frac{2x^3}{3} \right]_0^1 \\
 &= \frac{2}{5} \left[ \frac{1}{4} + \frac{2}{3} \right] = \frac{2}{5} \times \frac{3+8}{12} = \frac{11}{30}
 \end{aligned}$$

$$\begin{aligned}
 E(y) &= \int_{-\infty}^{+\infty} y f(y) dy = \int_0^{\infty} y e^{-y} dy = 1 \\
 E(y^2) &= \int_{-\infty}^{+\infty} y^2 f(y) dy = \int_0^{\infty} y^2 e^{-y} dy = 2 \\
 E(xy) &= \int_{-\infty}^{+\infty} \left[ \int_{-\infty}^{\infty} xy f(x, y) dy \right] dx \\
 &= \frac{2}{5} \int_0^1 x \left[ \int_0^{\infty} y e^{-y} dy \right] dx \\
 &= \frac{2}{5} \int_0^1 (x^2 + 2) \cdot 1 dx \\
 &= \frac{2}{5} \left[ \frac{x^3}{3} + \frac{2x^2}{2} \right]_0^1 = \frac{2}{5} \left( \frac{1}{3} + 1 \right) = \frac{8}{15}
 \end{aligned}$$

$$\begin{aligned}
 E(x^2y^2) &= \int_{-\infty}^{\infty} \left[ \int_{-\infty}^{\infty} x^2 y^2 f(x, y) dy \right] dx \\
 &= \int_0^1 x^2 (x+2) \left[ \int_0^{\infty} y^2 e^{-y} dy \right] dx \\
 &= \frac{2}{5} \int_0^1 (x^3 + 2x^2) 2 dx \\
 &= \frac{4}{5} \left[ \frac{x^4}{4} + \frac{2x^3}{3} \right]_0^1 \\
 &= \frac{4}{5} \left[ \frac{1}{4} + \frac{2}{3} \right] = \frac{11}{15}
 \end{aligned}$$

$$V(x) = E(x^2) - [E(x)]^2 = \frac{11}{30} - \left( \frac{8}{15} \right)^2 = 0.082$$

$$V(y) = E(y^2) - [E(y)]^2 = 2 - 1^2 = 1$$

$$V(xy) = E(x^2y^2) - [E(xy)]^2 = \frac{11}{15} - \left( \frac{8}{15} \right)^2 = 0.449$$

$$\text{Cov}(xy) = E(xy) - E(x) E(y) = \frac{8}{15} - \frac{8}{15} \times 1 = 0$$

$$r(x, y) = \frac{\text{Cov}(x, y)}{\sqrt{V(x)} \sqrt{V(y)}} = \frac{0}{\sqrt{0.082} \sqrt{1}} = 0$$

16.  $f(x, y) = \begin{cases} 2 & 0 < x < y < 1 \\ 0 & \text{otherwise} \end{cases}$

$$f(x) = \int_x^1 f(x, y) dy$$

$$= \int_x^1 2 dy$$

$$= 2 [y]_x^1 = 2 [1 - x]$$

$$f(y) = \int_0^y x f(x) dx$$

$$= \int_0^y 2 dx = 2 [x]_0^y = 2y$$

$$E(x) = \int_0^1 x f(x) dx$$

$$= \int_0^1 x \cdot 2(1-x) dx = 2 \int_0^1 (x - x^2) dx = 2 \left[ \frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 = 2 \left[ \frac{1}{2} - \frac{1}{3} \right] = \frac{1}{3}$$

$$E(x^2) = \int_0^1 x^2 f(x) dx$$

$$= \int_0^1 x^2 \cdot 2(1-x) dx = 2 \int_0^1 (x^2 - x^3) dx = 2 \left[ \frac{x^3}{3} - \frac{x^4}{4} \right]_0^1 = 2 \left[ \frac{1}{3} - \frac{1}{4} \right] = \frac{1}{6}$$

$$E(y) = \int_0^1 y f(y) dy = \int_0^1 y \cdot 2y dy = 2 \int_0^1 y^2 dy = 2 \left[ \frac{y^3}{3} \right]_0^1 = \frac{2}{3}$$

$$E(y^2) = \int_0^1 y^2 f(y) dy = \int_0^1 y^2 \cdot 2y dy = 2 \int_0^1 y^3 dy = 2 \left[ \frac{y^4}{4} \right]_0^1 = \frac{2}{4} = \frac{1}{2}$$

$$E(xy) = \int_0^1 \left[ \int_0^1 xy f(x, y) dy \right] dx$$

$$= \int_0^1 \left[ \int_0^1 xy \cdot z dy \right] dx$$

$$= \int_0^1 2x \left[ \int_0^1 y dy \right] dx = \int_0^1 2x \left[ \frac{y^2}{2} \right]_0^1 dx = \int_0^1 x dx = \left[ \frac{x^2}{2} \right]_0^1 = \frac{1}{2}$$

$$V(x) = E(x^2) - [E(x)]^2 = \frac{1}{6} - \left(\frac{1}{3}\right)^2 = \frac{1}{18}$$

$$V(y) = E(y^2) - [E(y)]^2 = \frac{1}{2} - \left(\frac{2}{3}\right)^2 = \frac{1}{18}$$

$$\text{Cov}(xy) = E(xy) - E(x) E(y) = \frac{1}{2} - \frac{1}{3} \times \frac{2}{3} = \frac{5}{18}$$

17.  $f(x, y) = \frac{9(1+x+y)}{2(1+x)^4(1+y)^4} \quad 0 < x < \infty$   
 $0 < y < \infty$

$$\begin{aligned} f(x) &= \int_{-\infty}^{+\infty} f(x, y) dy = \frac{9}{2} \int_0^{\infty} \frac{1+x+y}{(1+x)^4(1+y)^4} dy \\ &= \frac{9}{2} \int_0^{\infty} \frac{x+1+y}{(1+x)^4(1+y)^4} dy \\ &= \frac{9}{2} \left[ \int_0^{\infty} \frac{x}{(1+x)^4(1+y)^4} dy + \int_0^{\infty} \frac{1+y}{(1+y)^4(1+x)^4} dy \right] \\ &= \frac{9}{2} \left[ \frac{x}{(1+x)^4} \int_0^{\infty} (1+y)^{-4} dy + \frac{1}{(1+x)^4} \int_0^{\infty} (1+y)^{-3} dy \right] \\ &= \frac{9}{2} \left[ \frac{x}{3(1+x)^4} + \frac{1}{2(1+x)^4} \right] \\ &= \frac{9}{2} \left[ \frac{2x+3}{(1+x)^4} \right] = \frac{3}{4} \frac{(2x+3)}{(1+x)^4} \\ f(y) &= \int_{-\infty}^{+\infty} f(x, y) dx = \frac{9}{2} \times \frac{2y+3}{6(1+y)^4} = \frac{3}{4} \times \frac{(2y+3)}{(1+y)^4} \\ f(y/x) &= \frac{f(x/y)}{f(x)} = \frac{\frac{9(1+x+y)}{2(1+x)^4(1+y)^4}}{\frac{3(2x+3)}{4(1+x)^4}} = \frac{6(1+x+y)}{(2x+3)(1-y)^4} \end{aligned}$$

18. Comment on

Let  $x \sim B(n, p)$

$E(x) = 7$

$V(x) = 11$

Now,

$E(x) = np = 7 \dots \text{(i)}$

$V(x) = n p q = 11 \dots \text{(ii)}$

Substitute the value of  $np$  from (i) in (ii)

$7q = 11$

$\text{Or, } q = \frac{11}{7} = 1.57 > 1, \text{ Which is impossible.}$

Hence, given information is incorrect.

19. Find  $p$  if

$n = 6, 9p(x=4) = p(x=2), p = ?$

$\text{Here, } 9p(x=4) = p(x=2)$

$\text{Or, } 9c(6, 4)p^4q^2 = c(6, 2)p^2q^4$

$\text{Or, } 9 \times 15 p^4 q^2 = 15 p^2 q^4$

$\text{Or, } 9p^2 = q^2$

$\text{Or, } 9p^2 - q^2 = 0$

$\text{Or, } 9p^2 - (1-p)^2 = 0$

$$\text{Or, } 9p^2 - (1 - 2p + p^2) = 0$$

$$\text{Or, } 9p^2 - 1 + 2p - p^2 = 0$$

$$\text{Or, } 8p^2 + 2p - 1 = 0$$

$$\text{Or, } 8p^2 + 4p - 2p - 1 = 0$$

$$\text{Or, } 4p(2p + 1) - 1(2p + 1) = 0$$

$$\text{Or, } (2p + 1)(4p - 1) = 0$$

$$\therefore p = -\frac{1}{2}, \frac{1}{4}$$

$$p = -\frac{1}{2} \text{ is not possible} \quad \therefore p = \frac{1}{4}$$

20. In a binomial distribution

Let  $x \sim B(n, p)$  where  $x$  = no of success

$$n = 6$$

$$p(x = 3) = 0.2457$$

$$p(x = 4) = 0.0818$$

Now,

$$p(x = 3) = c(6, 3) p^3 q^3 = 0.2457 \quad \dots \quad (\text{i})$$

$$p(x = 4) = c(6, 4) p^4 q^2 = 0.0818 \quad \dots \quad (\text{ii})$$

Divide (i) by (ii)

$$\frac{C(6, 3) p^3 q^3}{C(6, 4) p^4 q^2} = \frac{0.2457}{0.0818}$$

$$\text{Or, } \frac{20q}{15p} = 3.003$$

$$\text{Or, } 4q = 3p \times 3.003$$

$$\text{Or, } 4(1-p) = 9.009p$$

$$\text{Or, } 4 - 4p = 9.009p$$

$$\text{Or, } 4 = 13.009p$$

$$\text{Or, } p = \frac{4}{13.009}$$

$$\therefore p = 0.307$$

$$q = 1 - p = 1 - 0.307 = 0.619$$

$$\text{Mean} = np = 6 \times 0.307 = 1.842$$

$$\text{Variance} = npq = 6 \times 0.307 \times 0.619 = 1.14$$

21. The mean and variance

Let  $x \sim B(n, p)$

$$E(x) = np = 3 \quad (\text{i})$$

$$V(x) = npq = 2 \quad (\text{ii})$$

Substitute value of np from (i) in (ii)  
 $3q = 2$       Or,  $q = \frac{2}{3}$

$$p = 1 - q = 1 - \frac{2}{3} = \frac{1}{3}$$

substitute the value of p in (i)

$$n \times \frac{1}{3} = 3 \quad \text{Or, } n = 9$$

$$\begin{aligned} \text{(i)} \quad p(x \leq 2) &= p(x = 0) + p(x = 1) + p(x = 2) \\ &= C(9, 0) p^0 q^9 + C(9, 1) p^1 q^8 + C(9, 2) p^2 q^7 \\ &= 1 \times \left(\frac{2}{3}\right)^9 + 9 \times \frac{1}{3} \times \left(\frac{2}{3}\right)^8 + 3 C \times \left(\frac{1}{3}\right)^2 \times \left(\frac{2}{3}\right)^7 \end{aligned}$$

$$= 0.026 + 0.117 + 0.234 = 0.377$$

$$\begin{aligned} \text{(ii)} \quad p(x \geq 7) &= p(x = 7) + p(x = 8) + p(x = 9) \\ &= C(9, 7) p^7 q^2 + C(9, 8) p^8 q^1 + C(9, 9) p^9 q^0 \\ &= 36 \times \left(\frac{1}{3}\right)^7 \left(\frac{2}{3}\right)^2 + 9 \times \left(\frac{1}{3}\right)^8 \left(\frac{2}{3}\right) + 1 \times \left(\frac{1}{3}\right)^9 \\ &= \left(\frac{1}{3}\right)^7 \left[ 16 + 2 + \frac{1}{9} \right] = \frac{18.111}{2187} = 0.0082 \end{aligned}$$

22. 12% of the items

Let, defective items = x

Prob. of defective  $p = 12\% = 0.12$ ,  $q = 1 - p = 0.88$

$n = 20$

$$p(x = 5) = c(20, 5) p^5 q^{15} = c(20, 5) (0.12)^5 (0.88)^{15} = 0.056$$

23. The average no.

Let  $x$  = defective pieces

$$\text{Prob. Of defective pieces (p)} = \frac{1}{10}, q = \frac{9}{10}$$

$n = 10$

$$p(x = 3) = c(10, 3) p^3 q^7 = c(10, 3) \left(\frac{1}{10}\right)^3 \left(\frac{9}{10}\right)^9 = 120 \times 0.001 \times 0.4782 = 0.057$$

24. Let  $x$  = defective computers

$$P = 5\% = 0.05, q = 0.95$$

$n = 20$ ,

$$P(X = 3) = {}^{20}C_3 (0.05)^3 (0.95)^{17} = 1140 \times 0.0000522 = 0.059$$

25. In a bombing

Prob. of hitting target  $p = 50\% = 0.5$

$$q = 1 - p = 0.5$$

No. of times hitting target =  $x$

No. of trial  $n = ?$

**74 CHAPTER - 6 | A Complete Solutions of Statistics I for BSc. CSIT**

$$(i) P(x \geq 1) \geq 99.9\%$$

$$\text{Or, } 1 - P(x < 1) \geq 99.9\%$$

$$\text{Or, } 1 - P(x = 0) \geq 0.999$$

$$\text{Or, } 1 - C(n, 0) p^0 q^n \geq 0.999$$

$$\text{Or, } 1 - (0.5)^n \geq 0.999$$

$$\text{Or, } 1 - 0.999 \geq (0.5)^n$$

$$\text{Or, } 0.001 \geq (0.5)^n$$

By trial method  $n = 10$

$$(ii) P(x \geq 2) \geq 99.9\%$$

$$\text{Or, } 1 - P(x < 2) \geq 99.9\%$$

$$\text{Or, } 1 - [P(x = 0) + P(x = 1)] \geq 0.999$$

$$\text{Or, } 1 - [C(n, 0) p^0 q^n + C(n - 1) p^n q^{n-1}] \geq 0.99$$

$$\text{Or, } 1 - [q^n + n p q^{n-1}] \geq 0.999$$

$$\text{Or, } 1 - [(0.5)^n + n (0.5) (0.5)^{n-1}] \geq 0.999$$

$$\text{Or, } 1 - [(0.5)^n + n (0.5)^n] \geq 0.999$$

$$\text{Or, } 1 - (0.5)^n [1 + n] \geq 0.999$$

$$\text{Or, } 1 - 0.999 \geq (0.5)^n [1 + n]$$

$$\text{Or, } 0.001 \geq (0.5)^n (1 + n)$$

By trial method  $n = 14$

26. Let  $x$  = No. of times success

$p$  = probability of hitting target

$$p = \frac{1}{3}, q = \frac{2}{3}$$

$$P(x \geq 1) > 90\%$$

$$\text{Or, } 1 - P(X < 1) > 0.9$$

$$\text{Or, } 0.1 > C(n, 0) \left(\frac{1}{3}\right)^0 \left(\frac{2}{3}\right)^n$$

$$\text{Or, } 0.1 > \left(\frac{2}{3}\right)^n$$

By trial method  $n = 6$

27. The probability that

Let,  $x$  = No. of students graduate

Prob. of students graduate  $p = 0.4, q = 1 - p = 1 - 0.4 = 0.6$

$$(i) P(x = 0) = C(n, x) p^x q^{n-x} = C(5, 0) (0.4)^0 (0.6)^{5-0} = 0.077$$

$$(ii) P(x = 1) = C(n, x) p^x q^{n-x} = C(5, 1) (0.4)^1 (0.6)^{5-1} = 0.25$$

$$(iii) P(x \geq 1) = 1 - P(x < 1) = 1 - P(x = 0) = 1 - 0.077 = 0.923$$

$$(iv) P(x = 5) = C(n, x) p^x q^{n-x} = C(5, 5) (0.4)^5 (0.6)^0 = 0.0102$$

28. At a particular university

Let  $x$  = No. of students with draw without computing course

$p$  = prob. of students withdraw without computing course

$$p = 20\% = 0.2, q = 1 - p = 0.8, n = 18$$

36. A car hire firm have

$$\text{Average } (\lambda) = 1.5$$

Let,  $x$  = No. of cars

$$P(x=0) = \frac{e^{-\lambda} \lambda^x}{x!} = \frac{e^{-1.5} (1.5)^0}{0!} = 0.223$$

$$P(x > 2) = 1 - P(x \leq 2)$$

$$= 1 - [P(x=0) + P(x=1) + P(x=2)]$$

$$= 1 - \left[ \frac{e^{-1.5} (1.5)^0}{0!} + \frac{e^{-1.5} (1.5)^1}{1!} + \frac{e^{-1.5} (1.5)^2}{2!} \right]$$

$$= 1 - [0.223 + 0.3495 + 0.2508] = 1 - 0.823 = 0.17$$

37. An automatic machine

Let,  $x$  = no. of defectives

$$\text{Prob. of defective } (p) = \frac{1}{400}$$

$$n = 100$$

$$\text{Average defective } (\lambda) = n p = 100 \times \frac{1}{400} = 0.25$$

$$\text{i. } P(x=0) = \frac{e^{-\lambda} \lambda^x}{x!} = \frac{e^{-0.25} (0.25)^0}{0!} = 0.778$$

$$\text{ii. } P(x \geq 1) = 1 - P(x < 1) = 1 - P(x=0) = 1 - 0.778 = 0.221$$

$$\text{iii. } P(x < 2) = P(x \leq 1) = P(x=0) + P(x=1)$$

$$= 0.778 + \frac{e^{-0.25} (0.25)^1}{1!} = 0.778 + 0.194 = 0.972$$

38. The chance of

Let  $x$  = No. of traffic accidents

Probability of traffic accident in a street  $(p) = 0.0005$

No. of street  $(n) = 1000$

Now,

$$\lambda = n p = 1000 \times 0.0005 = 0.5$$

$$\text{i. } P(x=0) = \frac{e^{-0.5} (0.5)^0}{0!} = 0.606$$

Consider in a year

$$\begin{aligned} \text{No. of days with no accidents} &= 365 \times P(x=0) \\ &= 365 \times 0.606 = 221.38 = 221 \end{aligned}$$

$$\text{iii. } P(x > 3) = 1 - P(x \leq 3)$$

$$= 1 - [P(x=0) + P(x=1) + P(x=2) + P(x=3)]$$

$$= 1 - \left[ \frac{e^{-0.5} (0.5)^0}{0!} + \frac{e^{-0.5} (0.5)^1}{1!} + \frac{e^{-0.5} (0.5)^2}{2!} \right] + e^{-0.5} (0.5)^3 / 3! = 1 - 0.606 - 0.125 - 0.0208 = 0.268$$

$$= 1 - e^{-0.5} (1 + 0.5 + 0.125 + 0.0208) = 1 - 1.645 e^{-0.5} = 0.003$$

No. of days with more than three accidents =  $365 \times P(x > 3)$

$$= 365 \times 0.003 = 1.14 = 1$$

39. Let  $\lambda$  = No. of telephone calls

$$\lambda = 3 \text{ per minute}$$

$$P(X = 0) = \frac{e^{-3} 3^0}{0!} = 0.049$$

$$\lambda = 3 \times 3 = 9 \text{ per three minute}$$

$$P(X \leq 4) = P(X = 0) + P(X = 1) + P(X = 2) + P(X = 3) + P(X = 4)$$

$$= \frac{e^{-9} 9^0}{0!} + \frac{e^{-9} 9^1}{1!} + \frac{e^{-9} 9^2}{2!} + \frac{e^{-9} 9^3}{3!} + \frac{e^{-9} 9^4}{4!}$$

$$= 0.945$$

40. The average

Let  $x$  = No. of network error

$$\text{Average no. network error } (\lambda) = 2.4$$

$$\text{i. } P(x = 0) = \frac{e^{-2.4} (2.4)^0}{0!} = 0.09$$

$$\text{ii. } P(x \geq 1) = 1 - P(x < 1) = 1 - P(x = 0) = 0.91$$

$$\text{iii. } P(x = 1) = \frac{e^{-2.4} (2.4)^1}{1!} = 0.21$$

41. Solution.

Let  $x$  = No. of messages arises

$$\lambda = 9 \text{ per hour}$$

$$\text{(a) } P(x \geq 3) = 1 - P(x < 3) = 1 - \left[ \frac{e^{-9} 9^0}{0!} + \frac{e^{-9} 9^1}{1!} + \frac{e^{-9} 9^2}{2!} \right]$$

$$= 0.997$$

$$\text{(b) } \lambda = 9 \times 2 = 18 \text{ per hour}$$

$$P(x = 5) = \frac{e^{-18} 18^5}{5!} = 0.066$$

42. A hospital has

$x$  = no. of emergency calls

$$\text{Average no. of emergency calls } (\lambda) = 4 \text{ per 10 min.}$$

$$\text{i. } \lambda = \frac{4}{10} = 0.4 \text{ per min.}$$

$$P(x \leq 2) = P(x = 0) + P(x = 1) + P(x = 2)$$

$$= \frac{e^{-0.4} (0.4)^0}{0!} + \frac{e^{-0.4} (0.4)^1}{1!} + \frac{e^{-0.4} (0.4)^2}{2!} = 0.67 + 0.268 + 0.053 = 0.99$$

$$\text{ii. } \lambda = 4 \text{ calls per 10 min.}$$

$$P(x = 3) = \frac{e^{-\lambda} \lambda^x}{x!} = \frac{e^{-4} 4^3}{3!} = 0.195$$

51. Suppose that the

Let  $x$  = no. of calls between 10 a. m. and 11 a. m.

$y$  = no. of calls between 11 a. m. and 12 noon.

$$x \sim P(\lambda); \lambda = 2$$

$$y \sim P(\lambda); \lambda = 4$$

$z = x + y$  = no. of calls between 10 a. m. and 12 noon

$$z \sim P(\lambda) \quad \lambda = 2 + 4 = 6$$

$$P(z \geq 5) = 1 - P(z < 5)$$

$$= 1 - [P(z = 0) + P(z = 1) + P(z = 2) + P(z = 3) + P(z = 4)]$$

$$= 1 - \left[ \frac{e^{-6} 6^0}{0!} + \frac{e^{-6} 6^1}{1!} + \frac{e^{-6} 6^2}{2!} + \frac{e^{-6} 6^3}{3!} + \frac{e^{-6} 6^4}{4!} \right]$$

$$= 1 - [0.0024 + 0.0148 + 0.0446 + 0.0892 + 0.133]$$

$$= 1 - 0.2848 = 0.7151$$

52. A source of

Mean no. of bacteria per c.c. ( $\lambda$ ) = 3

No. of test tube of capacity 1 c. c. ( $n$ ) = 10

Here,  $\lambda = n p$

$$\text{Or, } p = \frac{\lambda}{n} = \frac{3}{10}, q = 1 - p = \frac{7}{10}$$

Let  $x$  = No. of bacteria


$$\text{i. } P(x \geq 1) = 1 - P(x < 1) = 1 - P(x = 0) \\ = 1 - C(10, 0) (0.3)^0 (0.7)^{10-0} = 1 - 0.7^{10} = 0.971$$

$$\text{ii. } P(x = 7) = C(10, 7) (0.3)^7 (0.7)^{10-7} = 120 \times (0.3)^7 \times (0.7)^3 = 0.009$$

53. If  $Z$  is standard

$$Z \sim N(0, 1)$$


$$\text{a. } P(0 < Z < 1.34) = 0.4099$$



$$\text{b. } P(Z < 1.34) = 0.5 + P(0 < Z < 1.34)$$

$$= 0.5 + 0.4099$$


$$= 0.9099$$



$$\text{c. } P(Z > 1.96) = 0.5 - P(0 < Z < 1.96)$$

$$= 0.5 - 0.4750$$


$$= 0.025$$



$$\text{d. } P(1.34 < Z < 0)$$

$$= P(0 < Z < 1.34)$$

$$= 0.4099$$



82


## CHAPTER - 6 | A Complete Solutions of Statistics I for BSc. CSIT

c.  $P(Z < -1.34)$

$= 0.5 - P(-1.34 < Z < 0)$

$= 0.5 - P(0 < Z < 1.34)$

$= 0.5 - 0.4099 = 0.0901$




f.  $P(Z > -2.04)$

$= P(-2.04 < Z < 0) + 0.5$

$= P(0 < Z < 2.04) + 0.5$

$= 0.4793 + 0.5 = 0.9793$




g.  $P(1.96 < Z < 2.84)$

$= P(0 < Z < 2.84) - P(0 < Z < 1.96)$

$= 0.4977 - 0.4750$

$= 0.0227$




h.  $P(-1.96 < Z < 2)$

$= P(-1.96 < Z < 0) + P(0 < Z < 2)$

$= P(0 < Z < 1.96) + P(0 < Z < 2)$

$= 0.4750 + 0.4772$

$= 0.9522$




i.  $P(-2.48 < Z < -1.96)$

$= P(-2.48 < Z < 0) - P(-1.96 < Z < 0)$

$= P(0 < Z < 2.84) - P(0 < Z < 1.96)$

$= 0.4977 - 0.4750$

$= 0.0227$




j.  $P(Z < -1.96 \text{ or } Z > 1.96)$

$= 0.5 - P(-1.96 < Z < 0) + 0.5 - P(0 < Z < 1.96)$

$= 1 - 2P(0 < Z < 1.96)$

$= 1 - 2 \times 0.4750$

$= 1 - 0.95 = 0.05$

56. Suppose  $X$  follows the


$\mu = 100, \sigma = 10$

$X \sim N(\mu, \sigma^2)$

a.  $P(100 < X < 110) = ?$

Define  $Z = \frac{X - \mu}{\sigma} = \frac{X - 100}{10}$

When  $X = 100, Z = \frac{100 - 100}{10} = 0$




When  $X = 110, Z = \frac{110 - 100}{10} = 1$

$$\begin{aligned} P(100 < X < 110) &= P(0 < Z < 1) \\ &= 0.3413 \end{aligned}$$

b.  $P(X > 120)$   
When  $X = 120$ ,  
 $Z = \frac{120 - 100}{10} = 2$


$$\begin{aligned}P(X > 120) &= P(Z > 2) \\&= 0.5 - P(0 < Z < 2) \\&= 0.5 - 0.4772 = 0.0228\end{aligned}$$



c.  $P(X < 115)$   
When,  $X = 115$

$$Z = \frac{115 - 100}{10} = 1.5$$

$$\begin{aligned}P(X < 115) &= P(Z < 1.5) \\&= 0.5 + P(0 < Z < 1.5) \\&= 0.5 + 0.4332 = 0.8332\end{aligned}$$




d.  $P(80 < X < 100)$   
When  $X = 80$ ,

$$Z = \frac{80 - 100}{10} = -2$$


When  $X = 100$

$$Z = \frac{X - 100}{100} = \frac{100 - 100}{100} = 0$$

$$\begin{aligned}P(80 < X < 100) &= P(-2 < Z < 0) \\&= P(0 < Z < 2) = 0.4772\end{aligned}$$



e.  $P(X < 80) = P(Z < -2)$   
=  $0.5 - P(-2 < Z < 0)$   
=  $0.5 - P(0 < Z < 2)$   
=  $0.5 - 0.4772 = 0.028$




f.  $P(X > 90)$

When  $X = 90$

$$Z = \frac{90 - 100}{10} = -1$$


$$\begin{aligned}P(X > 90) &= P(Z > -1) \\&= P(-1 < Z < 0) + 0.5 \\&= P(0 < Z < 1) + 0.5 \\&= 0.3413 + 0.5 = 0.8413\end{aligned}$$



g.  $P(112 < X < 130)$

$$\text{When } X = 112, Z = \frac{112 - 100}{10} = 1.2$$

$$\text{When } X = 130, Z = \frac{130 - 100}{10} = 3.0$$



## 84 CHAPTER - 6 | A Complete Solutions of Statistics I for BSc. CSIT


$$\begin{aligned}
 P(112 < X < 130) &= P(1.2 < Z < 3) \\
 &= P(0 < Z < 3) - P(0 < Z < 1.2) \\
 &= 0.49865 - 0.3849 \\
 &= 0.11375
 \end{aligned}$$

h.  $P(85 < X < 125)$

$$\text{When } X = 85, Z = \frac{85 - 100}{10} = -1.5$$

$$\text{When } X = 125, Z = \frac{125 - 100}{10} = 2.5$$

$$\begin{aligned}
 P(85 < X < 125) &= P(-1.5 < Z < 2.5) \\
 &= P(-1.5 < Z < 0) + P(0 < Z < 2.5) \\
 &= 0.4332 + 0.4938 = 0.9270
 \end{aligned}$$




i.  $P(75 < X < 95)$

$$\text{When } X = 75, Z = \frac{75 - 100}{10} = -2.5$$

$$\text{When } X = 95, Z = \frac{95 - 100}{10} = -0.5$$

$$\begin{aligned}
 P(75 < X < 95) &= P(-2.5 < Z < -0.5) \\
 &= P(-2.5 < Z < 0) - P(-0 < Z < 0) \\
 &= P(0 < Z < 2.5) - P(0 < Z < 0.5) \\
 &= 0.4938 - 0.1915 = 0.3023
 \end{aligned}$$



### 57. Solution.

Let  $x$  = monthly production of computer parts  $x \sim N(\mu, \sigma^2)$


$$\mu = 100,000$$

$$\sigma = 20,000$$

(i)  $P(x > 120,000)$

$$\text{Define } Z = \frac{X - \mu}{\sigma} = \frac{X - 100000}{20000}$$

$$\text{When } X = 120,000, Z = \frac{120000 - 100000}{20000} = 1$$



$$\therefore P(X > 120,000) = P(Z > 1)$$

$$= 0.5 - P(0 < Z < 1)$$

$$= 0.5 - 0.3413 = 0.1587$$

(ii)  $P(x < 125000)$

$$\text{When } X = 125,000$$

$$Z = \frac{125000 - 100000}{20000} = 1.25$$


$$\therefore P(X < 125000)$$

$$= P(Z < 1.25)$$

$$= 0.5 + P(0 < Z < 1.25)$$


$$= 0.5 + 0.3944$$

$$= 0.8944$$



(iii)  $P(X > 60,000)$   
 When  $X = 60,000$   
 $Z = \frac{60000 - 100000}{20000} = -2$


$$\begin{aligned}P(X > 60,000) &= P(Z > -2) \\&= P(-2 < Z < 0) + 0.5 \\&= 0.4772 + 0.5 \\&= 0.9772\end{aligned}$$



(iv)  $P(X < 80,000)$   
 When  $X = 80,000$

$$Z = \frac{80000 - 100000}{20000} = -1$$

$$\begin{aligned}\therefore P(X < 80,000) &= P(Z < -1) \\&= 0.5 - P(-1 < Z < 0) \\&= 0.5 - P(0 < Z < 1) \\&= 0.5 - 0.3413 = 0.1587\end{aligned}$$




(v)  $P(1050 < X < 130000)$

$$\text{When } X = 1050, Z_1 = \frac{1050 - 100000}{20000} = -4.9$$

$$\text{When } X = 130000, Z_2 = \frac{130000 - 100000}{20000} = 1.5$$

$$\begin{aligned}P(1050 < X < 130000) &= P(-4.9 < Z < 1.5) + P(0 < Z < 1.5) \\&= P(0 < Z < 4.9) + P(0 < Z < 1.5) \\&= 0.5 + 0.4332 \\&= 0.9332\end{aligned}$$




(vi)  $P(70,000 < X < 140,000)$

$$\text{When } X = 70,000; Z_1 = \frac{70000 - 100000}{20000} = -1.5$$

$$\text{When } X = 140,000; Z_2 = \frac{140000 - 100000}{20000} = 2$$

$$\begin{aligned}P(70,000 < X < 140,000) &= P(-1.5 < Z < 2) \\&= P(-1.5 < Z < 0) + P(0 < Z < 2) \\&= P(0 < Z < 1.5) + P(0 < Z < 2) \\&= 0.4332 + 0.4772 \\&= 0.9104\end{aligned}$$




(vii)  $P(75,000 < X < 95,000)$

$$\text{When } X = 75,000; Z_1 = \frac{75000 - 100000}{20000} = -1.25$$

$$\text{When } X = 95,000; Z_2 = \frac{95000 - 100000}{20000} = -0.25$$

$$\begin{aligned}
 & P(75,000 < X < 95,000) \\
 & = P(-1.25 < Z < -0.25) \\
 & = P(-1.25 < Z < 0) - P(-0.25 < Z < 0) \\
 & = P(0 < Z < 1.25) - P(0 < Z < 0.25) \\
 & = 0.3944 - 0.0987 \\
 & = 0.2957
 \end{aligned}$$



58. The mean yield

Let,  $x$  = yield in kilos


$$x \sim N(\mu, \sigma^2), \mu = 662, \sigma = 32, N = 1000$$

i.  $P(x > 700)$

$$\text{Define, } Z = \frac{x - \mu}{\sigma} = \frac{x - 662}{32}$$

$$\text{When, } x = 700, z = \frac{700 - 662}{32} = 1.87$$

$$\begin{aligned}
 P(x > 700) &= P(z > 1.87) \\
 &= 0.5 - P(0 < z < 1.87) \\
 &= 0.5 - 0.381 = 0.119
 \end{aligned}$$




$$\therefore \text{No. of plots} = N P(x > 700) = 1000 \times 0.119 = 119$$

ii.  $P(x < 650)$

$$\text{When, } x = 650, z = \frac{x - 662}{32} = \frac{650 - 662}{32} = -0.375$$

$$\begin{aligned}
 P(x < 650) &= P(z < -0.375) \\
 &= 0.5 - P(-0.375 < z < 0.375) \\
 &= 0.5 - 0.144 \\
 &= 0.356
 \end{aligned}$$



$$\therefore \text{No. of puts} = N P(x < 650) = 1000 \times 0.356 = 356$$

ii. Let, lowest yield =  $x_1$

$$P(x \geq x_1) = \frac{100}{1000}$$

$$\text{Or, } P(x \geq x_1) = 0.1$$

$$\text{When } x = x_1$$

$$z = \frac{x_1 - 662}{32} = z_1 \text{ (say)} \quad \dots \dots \dots \text{ (i)}$$

Then,

$$P(x \geq x_1) = 0.1$$

$$\Rightarrow P(z \geq z_1) = 0.1$$

$$\Rightarrow 0.5 - P(0 \leq z \leq z_1) = 0.1$$

$$\Rightarrow 0.5 - 0.1 = P(0 \leq z \leq z_1)$$

$$\Rightarrow P(0 \leq z \leq z_1) = 0.4$$

Then  $z_1 = 1.28$

Substitute  $z_1$  in (i)

$$\frac{x_1 - 662}{32} = 1.28$$

$$\text{Or, } x_1 = 32 \times 1.28 + 662 = 702.96 \approx 703$$

Hence, lowest yield of best 100 plots is 703 kilos.

59. Let  $x$  = time taken to download,  $n = 95$

For one file

$$\mu = 16, \sigma = 5, \sigma^2 = 25$$

For 95 file

$$\mu = 16 \times 95 = 1520$$

$$\sigma^2 = 25 \times 95 = 2375 \quad \therefore \sigma = 48.73$$

$$\mu = 1520, \sigma = 48.73$$


$$P(x < 1500) = P\left(z \leq \frac{1500 - 1520}{48.73}\right)$$

$$= P(Z < -0.41)$$

$$= 0.5 - P(-0.41 < z < 0)$$

$$= 0.5 - 0.1591$$

$$= 0.3409$$



60. The local authorities

$$\text{No. of lamps (N)} = 1000$$

$$\text{Average life of lamp } (\mu) = 1000 \text{ hrs}$$

$$\text{S.D. } (\sigma) = 200 \text{ hrs}$$

Let,  $X$  = no. of burning hrs of lamp

$$\text{Define } Z = \frac{X - \mu}{\sigma} = \frac{X - 1000}{200}$$


- i.  $P(X < 800)$

$$\text{When, } X = 800, Z = \frac{800 - 1000}{200} = -1$$

$$P(X < 800) = P(Z < -1)$$

$$= 0.5 - P(-1 < Z < 0)$$

$$= 0.5 - 0.3413 = 0.1587$$



$$\text{No. of bulb burn fail in 800 hrs} = N P(X < 800) = 1000 \times 0.1587 = 158.7 \approx 159$$

- ii.  $P(800 < X < 1200)$

$$\text{When } X = 1200, Z = \frac{1200 - 1000}{200} = 1$$


$$P(800 < X < 1200) = P(-1 < Z < 1)$$

$$= P(-1 < Z < 0) + P(0 < Z < 1)$$

$$= 2 P(0 < Z < 1)$$

$$= 2 \times 0.3413$$

$$= 0.6826$$



No. of bulb burn between 800 and 1200 hrs

$$= N P(800 < X < 1200) = 1000 \times 0.6826 = 682.6 \approx 683$$

Let, after  $x = x_1$  hrs 10% lamp fail

$$P(x < x_1) = 10\%$$

When,  $x = x_1$ 

$$Z = \frac{x_1 - 1000}{200} = -Z_1 \text{ (say)} \quad \dots \dots \dots \text{(i)}$$

$$P(x < x_1) = 0.1$$

$$\text{Or, } P(z < -Z_1) = 0.1$$

$$\text{Or, } 0.5 - P(-Z_1 < z < 0) = 0.1$$

$$\text{Or, } 0.5 - 0.1 = P(0 < z < Z_1)$$

$$\text{Or, } P(0 < Z < Z_1) = 0.4$$

$$Z_1 = 1.28$$

Substitute  $Z_1$  in (i)

$$\frac{x_1 - 1000}{200} = -1.28$$

$$\text{Or, } x_1 - 1000 = -256$$

$$\text{Or, } x_1 = 1000 - 256 = 744$$

$\therefore$  10% lamps fails on burning 744 hrs

Let after  $x = x_2$  hrs. 10% lamp continue burning

$$P(x > x_2) = 10\%$$

$$\text{When } x = x_2, Z = \frac{x_2 - 1000}{200} = Z_2 \text{ (say)} \quad \dots \dots \dots \text{(ii)}$$

$$P(x > x_2) = 0.1$$

$$\text{Or, } P(z > Z_2) = 0.1$$

$$\text{Or, } 0.5 - 0.1 = P(0 < z < Z_2)$$

$$\text{Or, } P(0 < z < Z_2) = 0.4$$

$$Z_2 = 1.28$$

Substitute  $Z_2$  in (ii)

$$\frac{x_2 - 1000}{200} = 1.28$$

$$\text{Or, } x_2 - 1000 = 256$$

$$\text{Or, } x_2 = 1000 + 256 = 1256$$

$\therefore$  10% of lamps continue burning after 1256 hrs.

61. Let  $x$  = Time spent on training  $X \sim N(\mu, \sigma^2)$




$$\mu = 500, \sigma = 100$$

$$P(X < 420) = ?$$

$$\text{Let } Z = \frac{X - \mu}{\sigma} = \frac{X - 200}{100}$$

$$\text{When } X = 4200, Z = \frac{420 - 200}{100} = \frac{220}{100} = 2.2$$

$$P(X < 420) = P(Z < 2.2) = 0.5 + P(0 < Z < 2.2) = 0.5 + 0.4861 = 0.9861$$



62. Solution.

Let  $x$  = life time of electronic component

$$x \sim N(\mu, \sigma^2)$$

$$\mu = 5000, \sigma = 100$$

$$P(X < 5012) = ?$$

$$P(4000 < X < 6000) = ?$$

$$P(X > 7000) = ?$$

$$\text{Define } Z = \frac{X - \mu}{\sigma} = \frac{X - 5000}{100}$$

When  $X = 5012$ 

$$Z = \frac{5012 - 5000}{100} = 0.12$$

$$P(X < 5012) = P(Z < 0.12) = 0.452$$

When  $X = 4000$ 

$$Z = \frac{4000 - 5000}{100} = -10$$

When  $X = 6000$ 

$$Z = \frac{6000 - 5000}{100} = 10$$

$$P(4000 < X < 6000) = P(-10 < Z < 10) = 0.998$$

When  $X = 7000$ 

$$Z = \frac{7000 - 5000}{100} = 20$$

$$P(X > 7000) = P(Z > 20) = 0$$

63. Let,  $y$  = length of rod

$$X \sim N(\mu, \sigma^2)$$

$$\mu = 50, \sigma = 0.06$$

$$\text{Prob. of rod accepted} = P(49.85 \leq X \leq 50.15)$$

$$\text{Let } Z = \frac{X - \mu}{\sigma} = \frac{X - 50}{0.06}$$

$$\text{When } X = 49.85, Z = \frac{49.85 - 50}{0.06} = -2.5$$

$$\text{When } X = 50.15, Z = \frac{50.15 - 50}{0.06} = 2.5$$

$$P(49.85 \leq X \leq 50.15)$$

$$= P(-2.5 \leq Z \leq 2.5)$$



$$= P(-2.5 \leq Z \leq 0) + P(0 \leq Z \leq 2.5)$$

$$= 2P(0 \leq Z \leq 2.5)$$

$$= 2 \times 0.4938$$

$$= 0.9876$$

Hence 98.76% rods are accepted.



$$\text{Prob. of rod rejected} = P(X < 49.85) + P(x > 50.15)$$

$$= P(Z < -2.5) + P(Z > 2.5)$$

$$= 1 - 0.9876$$

$$= 0.0124$$

Hence, 1.24% rods are rejected.

64. Let  $X$  = Size of battery

$$X \sim N(\mu, \sigma^2)$$

$$\mu = 40, \sigma = 5$$

$$N = 1000$$

$$P(X > 35) = ?$$

$$\text{Let } Z = \frac{X - \mu}{\sigma} = \frac{X - 40}{5}$$

$$\text{When } X = 35, Z = \frac{35 - 40}{5} = -1$$

$$P(X > 35)$$

$$= P(Z > -1)$$

$$= P(-1 < Z < 0) + 0.5$$

$$= P(0 < Z < 1) + 0.5$$


$$= 0.3413 + 0.5$$

$$= 0.8413$$

$$\text{No. of cylinder need replacement after 35 days} = N P(X > 35)$$

$$= 1000 \times 0.8413$$

$$= 841.3 \approx 841$$



65. Income of a computer operator

Let  $X$  = income

$$\mu = 1520, \sigma = 160, X \sim N(\mu, \sigma^2)$$

$$N = 10,000$$

$$\% \text{ of poor} = \frac{2000}{10000} \times 100\% = 20\%$$

$$\% \text{ of rich} = \frac{1000}{10000} \times 100\% = 10\%$$

Let highest income of poorest 2000 =  $x_1$

Lowest income of richest 1000 =  $x_2$

$$\text{i. } P(X < x_1) = 20\%$$

$$\text{or, } P(X < x_1) = 0.2$$

$$\text{Define } Z = \frac{X - \mu}{\sigma} = \frac{X - 1520}{160}$$

$$\text{When, } X = x_1, Z = \frac{x_1 - 1520}{160} = -z_1 \text{ (say)} \quad \dots \dots \dots \text{(i)}$$

$$P(X < x_1) = 0.2$$

$$\text{Or, } P(Z < -Z_1) = 0.2$$

$$\text{Or, } 0.5 - P(-Z_1 < Z < 0) = 0.2$$

$$\text{Or, } 0.5 - 0.2 = P(-Z_1 < Z < 0)$$

$$\text{Or, } P(0 < Z < Z_1) = 0.3$$

$$\therefore Z_1 = 0.84$$

Substitute  $Z_1$  in (i)

$$\frac{x_1 - 1520}{160} = -0.84$$

$$\text{Or, } x_1 = -134.4 + 1520 = 1385.6$$

ii.  $P(X > x_2) = 10\%$

$$\text{Or, } P(X > x_2) = 0.1$$

$$\text{When } X = x_2, Z = \frac{x_2 - 1520}{160} = z_2 \text{ (say)} \quad \dots \quad (\text{ii})$$

$$P(X > x_2) = 0.1$$

$$\text{Or, } P(Z > z_2) = 0.1$$

$$\text{Or, } 0.5 - P(0 < Z < z_2) = 0.1$$

$$\text{Or, } 0.5 - 0.1 = P(0 < Z < z_2)$$

$$\text{Or, } P(0 < Z < z_2) = 0.4$$

$$\text{Or, } z_2 = 1.28$$

Substitute  $z_2$  in (ii)

$$\frac{x_2 - 1520}{160} = 1.28$$

$$x_2 = 204.80 + 1520 = 1724.80$$

66. Let life time of calculator =  $x$

$$X \sim N(\mu_1, \sigma^2), \mu = 54, \sigma = 8$$

Let warranty life time =  $x_1$

$$P(X \leq X_1) = 22\% = 0.22$$

$$\text{Define } Z = \frac{X - \mu}{\sigma} = \frac{X - 54}{8}$$

$$\text{When } X = X_1; Z = \frac{X_1 - 54}{8} = -Z_1 \text{ (say)} \quad \dots \quad (1)$$


$$P(X \leq X_1) = 0.22$$

$$\Rightarrow P(Z \leq -Z_1) = 0.22$$

$$\Rightarrow 0.5 - 0.22 = P(0 < Z < Z_1)$$

$$\Rightarrow 0.28 = P(0 < Z < Z_1)$$

$$\Rightarrow Z_1 = 0.77$$



Substitute  $Z_1$  in equation (1)

$$\frac{X_1 - 54}{8} = -0.77$$

$$\text{or } x_1 - 54 = -6.16$$

$$\text{or } x_1 = 54 - 6.16 = 47.84$$

Hence, warranty period should be 47.84 months.

67. Let  $x$  = marks secured by  $y$  students

$$N = 5000$$

$$X \sim N(65, 40)$$

Let lowest mark of top 10% student =  $x_1$

$$P(X \geq x_1) = 10\% = 0.1$$

$$\text{Define } Z = \frac{X - \mu}{\sigma}$$

$$\text{When } X = x_1, Z = \frac{x_1 - 65}{\sqrt{40}} = z_1 \text{ (say)} \quad \dots \text{(i)}$$

$$P(X \geq x_1) = 0.1$$

$$\Rightarrow P(Z \geq z_1) = 0.1$$

$$\Rightarrow 0.5 - P(0 \leq Z \leq z_1) = 0.1$$

$$\Rightarrow 0.5 - 0.1 = P(0 \leq Z \leq z_1)$$

$$\Rightarrow 0.4 = P(0 \leq Z \leq z_1)$$

$$\Rightarrow z_1 = 1.28$$

Substitute  $z_1$  in (i),

$$\frac{x_1 - 65}{\sqrt{40}} = 1.28$$

$$\Rightarrow \frac{x_1 - 65}{6.324} = 1.28$$

$$\Rightarrow x_1 = 6.324 \times 1.28 + 65 \\ = 8.09 + 65 = 73.09$$

Hence lowest marks of top 10% student is 73.09.

Let highest marks by poorest 500 students =  $x_2$ .

$$P(X \leq x_2) = \frac{500}{5000}$$

$$\Rightarrow P(X \leq x_2) = 0.1$$

$$\text{When } X = x_2, Z = \frac{x_2 - 65}{\sqrt{40}} = -Z_2 \text{ (say)} \quad \dots \text{(ii)}$$

$$P(X \leq x_2) = 0.1$$



$$\Rightarrow P(Z \leq -Z_2) = 0.1$$

$$\Rightarrow 0.5 - P(-Z_2 \leq Z < 0) = 0.1$$

$$\Rightarrow 0.5 - 0.1 = P(0 \leq Z \leq Z_2)$$

$$\Rightarrow P(0 \leq Z \leq Z_2) = 0.4$$

$$\Rightarrow Z_2 = 1.28$$



Substitute  $Z_2$  in (ii)

$$\frac{x_2 - 65}{\sqrt{40}} = -1.28$$

$$\Rightarrow \frac{x_2 - 65}{6.324} = -1.28$$

$$\Rightarrow x_2 = 6.324 \times (-1.28) + 65 = 56.91$$

Hence, highest marks of poorest 500 students is 56.91.

Let limit of marks of middle 80% students be  $x_3$  and  $x_4$

$$P(x_3 \leq X \leq x_4) = 80\%$$

$$\Rightarrow P(X_3 \leq X \leq X_4) = 0.8$$

When  $X = X_3$ ;

$$Z = \frac{X_3 - 65}{6.324} = -Z_3 \text{ (Say)} \dots (2)$$

When  $X = X_4$ ;

$$Z = \frac{X_4 - 64}{6.324} = Z_4 \text{ (Say)} \dots (3)$$

$$P(X_3 \leq X \leq X_4) = 0.8$$

$$\Rightarrow P(-2/3) \leq Z \leq Z_4 = 0.8$$

$$\Rightarrow 2P(0 \leq Z \leq Z_4) = 0.8 \text{ (Since it is middle)}$$

$$\Rightarrow P(0 \leq Z \leq Z_4) = 0.4$$

$$\Rightarrow Z_4 = 1.28, Z_3 = 1.28$$

Substitute values

$$\frac{X_3 - 65}{6.324} = -1.28$$

$$\Rightarrow X_3 = 56.91$$

$$\frac{X_4 - 65}{6.324} = 1.28$$

$$\Rightarrow X_4 = 73.09$$

68. Let  $x$  = marks secured

$$P(X > 80) = 5\%$$

$$P(X < 30) = 10\%$$

$$Z = \frac{X - \mu}{\sigma}$$

$$\text{When } X = 80, Z = \frac{80 - \mu}{\sigma} = Z_1 \text{ (Say)} \dots (1)$$

$$\text{When } X = 30, Z = \frac{30 - \mu}{\sigma} = -Z_2 \text{ (say)} \dots (2)$$



$$P(X > 80) = 5\%$$

$$\text{Or } P(Z > Z_1) = 0.05$$

$$\text{Or } 0.5 - P(0 < Z < Z_1) = 0.05$$

$$\text{Or } P(0 < Z < Z_1) = 0.45$$

$$\text{Or } Z_1 = 1.64$$



**94 CHAPTER - 6 | A Complete Solutions of Statistics I for BSc. CSIT**


Or  $P(X < 30) = 10\%$

Or  $P(Z < -Z_2) = 0.1$

Or  $0.5 - P(-Z_2 < Z < 0) = 0.1$

Or  $0.4 = P(0 < Z < Z_2)$

Or  $Z_2 = 1.28$



Substitute  $Z_1$  in equation (1)  $\frac{80 - \mu}{\sigma} = 1.64 \dots \dots (3)$

Substitute  $Z_2$  in equation (2)  $\frac{30 - \mu}{\sigma} = -1.28 \dots \dots (4)$

Divide  $\frac{80 - \mu}{\sigma} = -1.3125$

Or  $\mu = 49.02$


Substitute  $\mu$  in (3);  $\sigma = 18.88$

$P(45 < X < 60)$

When  $X = 45$ ,  $Z = \frac{45 - 49.02}{18.88} = -0.21$

When  $X = 60$ ,  $Z = \frac{60 - 49.02}{18.88} = 0.58$

$$\begin{aligned} P(45 < X < 60) &= P(-0.21 < Z < 0.58) \\ &= P(-0.21 < Z < 0) + P(0 < Z < 0.58) \\ &= 0.0832 + 0.219 = 0.3022 = 30.22\% \end{aligned}$$



69. The mean IQ

Let  $X \sim N(\mu, \sigma^2)$   $\mu = 100, \sigma = 16$


$X$  = IQ score of children

Define  $Z = \frac{X - \mu}{\sigma} = \frac{X - 100}{16}$

i.  $P(X < 80)$

When  $X = 80$ ,  $Z = \frac{80 - 100}{16} = -1.25$

$$\begin{aligned} P(X < 80) &= P(Z < -1.25) \\ &= 0.5 - P(-1.25 < Z < 0) \\ &= 0.5 - P(0 < Z < 1.25) \\ &= 0.5 - 0.3944 = 0.1056 = 10.56\% \end{aligned}$$



Let limit for middle 40% students be  $x_1$  and  $x_2$

$P(x_1 < X < x_2) = 40\%$

When,  $X = x_1, Z = \frac{x_1 - 100}{16} = -Z_1 \dots \dots \dots (i)$

$X = x_2, Z = \frac{x_2 - 100}{16} = Z_1 \dots \dots \dots (ii)$

$$\begin{aligned} P(x_1 < x < x_2) &= 0.4 \\ \text{Or, } P(-z_1 < z < z_1) &= 0.4 \\ \text{Or, } 2P(0 < z < z_1) &= 0.4 \\ \text{Or, } P(0 < z < z_1) &= 0.2 \\ \therefore z_1 &= 0.52 \end{aligned}$$

Substitute  $z_1$  in (i) and (ii)

$$\frac{x_1 - 100}{10} = -0.52$$

$$\text{Or, } x_1 = -5.2 + 100 = 94.8 \approx 95$$

$$\frac{x_2 - 100}{10} = 0.52$$

$$\text{Or, } x_2 = 5.2 + 100 = 105.2 \approx 105$$

Hence limit of IQ for middle 40% is 95 to 105

$$\begin{aligned} \text{IQ range} &= \mu \pm 1.96 \times 16 \\ &= 100 \pm 31.36 \\ &= (68.64, 131.36) \end{aligned}$$

$$P(68.64 < X < 131.36)$$

$$\text{When } x = 68.64, Z = \frac{68.64 - 100}{16} = -1.96$$

$$x = 131.36, Z = \frac{131.36 - 100}{16} = 1.96$$

$$\begin{aligned} P(68.64 < X < 131.36) &= P(-1.96 < Z < 1.96) \\ &= P(-1.96 < Z < 0) + P(0 < Z < 1.96) \\ &= 2P(0 < Z < 1.96) \\ &= 2 \times 0.475 = 0.95 \end{aligned}$$

$\therefore$  95 children had IQ with range  $\mu \pm 1.96\sigma$

70. The life of

Let  $x$  = life of printer

$$\mu = 4, \sigma = 200$$

$$P(x < 400) = 10\%$$

$$\text{Or, } P(x < 400) = 0.1$$

$$\text{Define } Z = \frac{X - \mu}{\sigma}$$


$$\text{When } X = 400, Z = \frac{400 - \mu}{200} = -z_1 \text{ (say)} \quad \dots \dots \dots \text{ (i)}$$

$$P(X < 400) = 0.1$$

$$\Rightarrow P(Z < -z_1) = 0.1$$

$$\Rightarrow 0.5 - P(-z_1 < Z < 0) = 0.1$$

$$\Rightarrow 0.5 - 0.1 = P(-z_1 < Z < 0)$$



$$\Rightarrow P(0 < Z < Z_1) = 0.4$$

$$\therefore Z_1 = 1.28$$

Substitute  $Z_1$  in (i)


$$\frac{400 - \mu}{200} = -1.28$$

$$\text{Or, } 400 - \mu = -256$$

$$\text{Or, } 400 + 256 = \mu$$

$$\therefore \mu = 656$$

Hence mean life of bulb is 656 hrs.



71. Solution.

Let Daily sale of book =  $X$

$$X \sim N(\mu, \sigma^2)$$

$$\sigma = 5000; \mu = ?$$

$$P(X > 40,000) = 90\%$$

$$P(X > 40,000) = 0.9$$

$$\text{Let } Z = \frac{X - \mu}{\sigma}$$

$$\text{When } X = 40,000; Z = \frac{40000 - \mu}{5000} = -Z_1 \text{ (say)} \dots \dots (1)$$

$$P(X > 40,000) = 0.9$$

$$\Rightarrow P(Z > -Z_1) = 0.9$$

$$\Rightarrow 0.5 + P(-Z_1 < Z < 0) = 0.9$$

$$\Rightarrow P(0 < Z < Z_1) = 0.4$$

$$\Rightarrow Z_1 = 1.28$$


Substitute  $Z_1$  in equation (1)

$$\frac{40000 - \mu}{5000} = -1.28$$

$$\Rightarrow 40,000 - \mu = -6400$$

$$\Rightarrow 40,000 + 6400 = \mu$$

$$\mu = 46400$$



72. Sacks of grain

Let,  $x$  = weight of bag

$$\mu = 114, X \sim N(\mu, \sigma^2), \sigma = ?$$

$$\text{Here } P(X > 115) = 15\%$$

$$\text{Or, } P(X > 115) = 0.15$$

$$\text{Define } Z = \frac{X - \mu}{\sigma}$$

$$\text{When } X = 115, Z = \frac{115 - 114}{\sigma} = \frac{1}{\sigma} = z_1 \text{ (say)} \dots \dots \dots (i)$$

$$P(X > 115) = 0.15$$

$$\Rightarrow P(Z > z_1) = 0.15$$

$$\Rightarrow 0.5 - P(0 < Z < z_1) = 0.15$$



$$\begin{aligned} \Rightarrow 0.5 - 0.15 &= P(0 < Z < Z_1) \\ \Rightarrow P(0 < Z < Z_1) &= 0.35 \\ \Rightarrow Z_1 &= 1.04 \\ \therefore \text{Substitute } Z_1 \text{ in (i)} \\ \frac{1}{\sigma} &= 1.04 \end{aligned}$$

$$\text{Or, } \sigma = \frac{1}{1.04} = 0.96$$

Hence, S.D. = 0.96.

73. Solution.  
Let marks secured =  $X$

$$X \sim N(\mu, \sigma^2)$$

$$P(X > 60) = 10\%; \quad P(X < 40) = 30\%$$



$$\text{Define } Z = \frac{X - \mu}{\sigma}$$

$$\text{When } X = 60, Z = \frac{60 - \mu}{\sigma} = Z_1 \text{ (say)}$$

.... (1)

$$\text{When } X = 40, Z = \frac{40 - \mu}{\sigma} = -Z_2 \text{ (say)}$$

.... (2)



$$\begin{aligned} P(X > 60) &= 10\% \\ \Rightarrow P(Z > Z_1) &= 0.1 \\ \Rightarrow 0.5 - P(0 < Z < Z_1) &= 0.1 \\ \Rightarrow 0.5 - 0.1 &= P(0 < Z < Z_1) \\ \Rightarrow P(0 < Z < Z_1) &= 0.4 \\ \Rightarrow Z_1 &= 1.28 \\ \therefore \text{Substitute } Z_1 \text{ in equation (1)} \end{aligned}$$

$$\frac{60 - \mu}{\sigma} = 1.28 \quad \dots (3)$$

$$P(X < 40) = 30\%$$

$$\begin{aligned} \Rightarrow P(Z < -Z_2) &= 0.3 \\ \Rightarrow 0.5 - P(-Z_2 < Z < 0) &= 0.3 \\ \Rightarrow 0.5 - 0.3 &= P(-Z_2 < Z < 0) \\ \Rightarrow P(0 < Z < Z_2) &= 0.2 \\ \Rightarrow Z_2 &= 0.52 \end{aligned}$$

Substitute  $Z_2$  in equation (2)

$$\frac{40 - \mu}{\sigma} = -0.52 \quad \dots (4)$$

Divide (3) by (4)

$$\frac{\frac{60 - \mu}{\sigma}}{\frac{40 - \mu}{\sigma}} = \frac{1.28}{-0.52}$$

$$\begin{aligned} \Rightarrow \frac{60 - \mu}{40 - \mu} &= -2.461 \\ \Rightarrow 60 - \mu &= -98.46 + 2.461 \mu \\ \Rightarrow 60 + 98.46 &= 2.461 \mu + \mu \end{aligned}$$

$$\Rightarrow \mu = \frac{158.46}{3.461} = 45.78$$

Substitute  $\mu$  in equation (3)

$$\frac{60 - 45.78}{\sigma} = 1.28$$

$$\Rightarrow \sigma = \frac{14.22}{1.28} = 11.109$$

74. Of a large group of

Let,  $x \sim N(\mu, \sigma^2)$  where,  $x$  = height

$$P(x < 60) = 0.05$$

$$P(60 < x < 65) = 0.4$$

$$\text{Define } z = \frac{x - \mu}{\sigma}$$

$$\text{When } x = 60, z = \frac{60 - \mu}{\sigma} = -z_1 \text{ (say)} \quad \dots \dots \dots \text{(i)}$$

$$\text{When } x = 65, z = \frac{65 - \mu}{\sigma} = -z_2 \text{ (say)} \quad \dots \dots \dots \text{(ii)}$$

Now,

$$P(x < 60) = 0.05$$

$$\Rightarrow P(z < -z_1) = 0.05$$

$$\Rightarrow 0.5 - P(-z_1 < z < 0) = 0.05$$

$$\Rightarrow 0.5 - 0.05 = P(-z_1 < z < 0)$$

$$\Rightarrow P(0 < z < z_1) = 0.45$$

$$z_1 = 1.64$$

Substitute  $z_1$  in (i)

$$\frac{60 - \mu}{\sigma} = -1.64 \quad \dots \dots \dots \text{(iii)}$$

$$P(60 < x < 65) = 0.4$$

$$\Rightarrow P(-z_1 < z < -z_2) = 0.4$$

$$\Rightarrow P(-z_1 < z < 0) - P(-z_2 < z < 0) = 0.4$$

$$\Rightarrow P(0 < z < z_1) - P(0 < z < z_2) = 0.4$$

$$\Rightarrow 0.45 - 0.4 = P(0 < z < z_2)$$

$$\Rightarrow P(0 < z < z_2) = 0.05$$

$$\therefore z_2 = 0.13$$

Substitute  $z_2$  in (ii)

$$\frac{65 - \mu}{\sigma} = -0.13 \quad \dots \dots \dots \text{(iv)}$$



Divide (iii) by (iv)

$$\frac{\frac{60 - \mu}{\sigma}}{\frac{65 - \mu}{\sigma}} = \frac{-1.64}{-0.13}$$

$$0.13(60 - \mu) = 1.64(65 - \mu)$$

$$\text{Or, } 7.8 - 0.13\mu = 106.6 - 1.64\mu$$

$$\text{Or, } 1.64\mu - 0.13\mu = 106.6 - 7.8$$



$$\text{Or, } 1.51 \mu = 98.8$$

$$\therefore \mu = \frac{98.8}{1.51} = 65.43$$

Substitute  $\mu$  in (iv)

$$\frac{65 - 65.43}{\sigma} = -0.13$$

$$\text{Or, } \sigma = \frac{-0.43}{-0.13} = 3.30$$

Hence, mean = 65.43 and S.D. = 3.3

75. If the skulls

Let  $x$  = length breadth index

$$X \sim N(\mu, \sigma^2)$$

$$P(x < 75) = 58\%; P(75 < x < 80) = 38\%; P(x > 80) = 4\%$$

$$\text{Define } z = \frac{x - \mu}{\sigma}$$

$$\text{When, } X = 75, z = \frac{75 - \mu}{\sigma} = z_1 \text{ (say)} \quad \dots \quad (\text{i})$$

$$\text{When, } X = 80, z = \frac{80 - \mu}{\sigma} = z_2 \text{ (say)} \quad \dots \quad (\text{ii})$$

Now,

$$P(x < 75) = 0.58$$

$$\Rightarrow P(z < z_1) = 0.58$$

$$\Rightarrow 0.5 + P(0 < z < z_1) = 0.58$$

$$\Rightarrow P(0 < z < z_1) = 0.58 - 0.5$$

$$\Rightarrow P(0 < z < z_1) = 0.08$$

$$\therefore z_1 = 0.2$$

Substitute  $z_1$  in (i)

$$\frac{75 - \mu}{\sigma} = 0.2$$

$$\text{Or, } 75 = 0.2 \sigma + \mu$$

$$\therefore \mu + 0.2 \sigma = 75$$

$$P(x > 80) = 4\%$$

$$\Rightarrow P(z > z_2) = 0.04$$

$$\Rightarrow 0.5 - P(0 < z < z_2) = 0.04$$

$$\Rightarrow 0.5 - 0.04 = P(0 < z < z_2)$$

$$\Rightarrow P(0 < z < z_2) = 0.46$$

$$Z_2 = 1.75$$


Substitute  $z_2$  in (ii)

$$\frac{80 - \mu}{\sigma} = 1.75$$


$$\text{Or, } 80 = 1.75 \sigma + \mu$$

$$\therefore \mu + 1.75 \sigma = 80$$

$$\text{Solving (iii) and (iv) } \mu = 74.35; \sigma = 3.22$$



..... (iii)



..... (iv)

76. Let waiting time =  $x$  hrs;  $\theta = 5$

$$P(x \geq \frac{15}{60}) = ?$$

$$\begin{aligned}
 P(x \geq 0.25) &= \int_{15}^{\infty} \theta e^{-\theta x} dx \\
 &= \int_{0.25}^{\infty} 5 e^{-5x} dx \\
 &= 5 \left[ \frac{e^{-5x}}{-5} \right]_{0.25}^{\infty} = -[e^{-5x}]_{0.25}^{\infty} = -[e^{-\infty} - e^{-1.25}] = e^{-1.25} = 0.28
 \end{aligned}$$

$$E(x) = \frac{1}{\theta} = \frac{1}{5} \text{ hr} = 12 \text{ min}$$

77.  $y_0 = 2000 \Rightarrow \theta = \frac{1}{2000}$

Let  $x$  = daily consumption of electricity

$$\begin{aligned}
 P(x \geq 1500) &= \int_{1500}^{\infty} \theta e^{-\theta x} dx \\
 &= \int_{1500}^{\infty} \frac{1}{2000} e^{-\frac{1}{2000}x} dx \\
 &= \frac{1}{2000} \left[ \frac{e^{-\frac{x}{2000}}}{-\frac{1}{2000}} \right]_{1500}^{\infty} \\
 &= - \left[ e^{-\infty} - e^{-\frac{1500}{2000}} \right] \\
 &= e^{-0.75} \\
 &= 0.47
 \end{aligned}$$

$$\begin{aligned}
 P(x \leq 2500) &= \int_{\infty}^{2500} \theta e^{-\theta x} dx \\
 &= \int_{\infty}^{2500} \frac{1}{2000} e^{-\frac{1}{2000}x} dx \\
 &= \frac{1}{2000} \left[ \frac{e^{-\frac{1}{2000}x}}{-\frac{1}{2000}} \right]_{0}^{2500} \\
 &= - \left[ e^{-\frac{x}{2000}} \right]_{0}^{2500} \\
 &= - \left[ e^{-\frac{2500}{2000}} - e^0 \right] \\
 &= - e^{-1.25} + 1 \\
 &= 1 - e^{-1.25} \\
 &= 1 - 0.286 \\
 &= 0.713
 \end{aligned}$$

### 78. Solution.

Let  $x$  = No. of pages print by printer for min.

$$\lambda = 15$$

Here,  $\mu = 15, \sigma = 15$

$$P(x > 25.5) = ? \quad x = 25.5 \text{ due to continuity correction for } x > 25.5$$

$$P(x < 10) = ?$$

$$\text{Let } z = \frac{x - \mu}{\sigma} = \frac{x - 15}{15}$$

When  $x = 25.5$

$$z = \frac{25.5 - 15}{15} = \frac{10.5}{15} = 0.7$$

$$\begin{aligned} P(x > 25.5) &= P(z > 0.7) \\ &= 0.5 - P(0 < z < 0.7) \\ &= 0.5 - 0.258 \\ &= 0.242 \end{aligned}$$

$x = 9.5$  due to continuity correction for  $x < 10$

When  $x = 9.5$

$$z = \frac{9.5}{15} = \frac{-5.5}{15} = -0.36$$


$$\begin{aligned} P(x < 9.5) &= P(z < -0.36) \\ &= 0.5 - P(-0.36 < z < 0) \\ &= 0.5 - P(0 < z < 0.36) \\ &= 0.5 - 0.1406 \\ &= 0.3594 \end{aligned}$$

### 79. Solution.

$$\alpha = 2$$

$$\begin{aligned} P(x > 2) &= \int_2^{\infty} \frac{e^{-x} x^{\alpha-1}}{\sqrt{2}} dx \\ &= \int_2^{\infty} \frac{e^{-x} x^{\alpha-1}}{\sqrt{\alpha}} dx \\ &= \int_2^{\infty} \frac{e^{-x} x^{2-1}}{\sqrt{2}} dx \\ &= \int_2^{\infty} x e^{-x} dx \\ &= \left[ x \int e^{-x} dx - \int \left( \frac{dx}{dx} \int e^{-x} dx \right) dx \right]_2^{\infty} \\ &= \left[ -x e^{-x} - e^{-x} \right]_2^{\infty} \\ &= -0 - 0 + 2e^{-2} + e^{-2} \\ &= 3e^{-2} = 3 \times 0.135 = 0.405 \end{aligned}$$

$$\begin{aligned} P(3 < x < 5) &= \int_3^5 \frac{e^{-x} x^{\alpha-1}}{\sqrt{\alpha}} dx \\ &= \int_3^5 x e^{-x} dx \\ &= \left[ -x e^{-x} - e^{-x} \right]_3^5 \\ &= -5e^{-5} - e^{-5} + 3e^{-3} + e^{-3} \\ &= -6e^{-5} + 4e^{-3} \\ &= -0.040 + 0.199 \\ &= 0.159 \end{aligned}$$



## Practical Questions

1. Fit the binomial

x	0	1	2	3	4	5	6	Total
f	7	6	19	35	23	7	1	$\sum f = 98$
fx	0	6	38	105	92	35	6	$\sum fx = 282$

$$\bar{x} = \frac{\sum fx}{N} = \frac{282}{98} = 2.877$$

Here, n = 6

$$\bar{x} = np$$

$$\text{Or, } 2.877 = 6p$$

$$\text{Or, } p = \frac{2.877}{6} = 0.479, q = 0.52$$

x	$P(x) = C(n, x) p^x q^{n-x}$	Expected frequency = N P(x)
0	$C(6, 0) (0.479)^0 (0.52)^6 = 0.019$	1.86 ≈ 2
1	$C(6, 1) (0.479)^1 (0.52)^5 = 0.1092$	10.70 ≈ 11
2	$C(6, 2) (0.479)^2 (0.52)^4 = 0.2516$	24.65 ≈ 25
3	$C(6, 3) (0.479)^3 (0.52)^3 = 0.3090$	30.28 ≈ 30
4	$C(6, 4) (0.479)^4 (0.52)^2 = 0.2153$	21.1 ≈ 21
5	$C(6, 5) (0.479)^5 (0.52)^1 = 0.078$	7.64 ≈ 8
6	$C(6, 6) (0.479)^6 = 0.012$	1.176 ≈ 1

2. Five fair coin

No. of heads (x)	Frequency (f)	fx
0	2	0
1	10	10
2	24	48
3	38	114
4	18	72
5	8	40
	$N = \sum f = 100$	$\sum fx = 284$

$$\bar{x} = \frac{\sum fx}{N} = \frac{284}{100} = 2.84$$

n = 5

$$\bar{x} = n p$$

$$\text{Or, } 2.84 = 5p$$

$$\text{Or, } p = \frac{2.84}{5} = 0.568; q = 1 - p = 0.432$$

$$x \sim B(n, p)$$

x	$P(x) = C(n, x) p^x q^{n-x}$	Expected frequency = N P(x)
0	$C(5, 0) (0.568)^0 (0.432)^5 = 0.015$	1.5 ≈ 2
1	$C(5, 1) (0.568)^1 (0.432)^4 = 0.098$	9.8 ≈ 10
2	$C(5, 2) (0.568)^2 (0.432)^3 = 0.2601$	26.01 ≈ 26
3	$C(5, 3) (0.568)^3 (0.432)^2 = 0.341$	34.10 ≈ 34
4	$C(5, 4) (0.568)^4 (0.432)^1 = 0.224$	22.4 ≈ 22
5	$C(5, 5) (0.568)^5 (0.432)^0 = 0.059$	5.9 ≈ 6

## 3. Solution.

192 Families

No. of girl child (x)	No. of families (f)	$f_x$
0	77	0
1	90	90
2	20	40
3	5	15
	$N = \sum f = 192$	$\sum f_x = 145$

$$\bar{x} = \frac{\sum f x}{N}$$

$$= \frac{145}{192} = 0.755$$

$$n = 3$$

(i) When both sexes are equally probable

$$p = q = \frac{1}{2}$$

$$n = 3$$

No. of girl child (x)	No. of families (f)	$f_x$
0	77	0
1	90	90
2	20	40
3	5	15
	$N = \sum f = 192$	$\sum f_x = 145$

$$\bar{x} = \frac{\sum f x}{N} = \frac{145}{192} = 0.755; \quad n = 3$$

i. When both sexes are equally probable

$$p = q = \frac{1}{2} \quad n = 3$$

x	$P(x) = C(n, x) p^x q^{n-x}$	Expected frequency = $N P(x)$
0	$C(3, 0) \left(\frac{1}{2}\right)^0 \left(\frac{1}{2}\right)^3 = 0.125$	24
1	$C(3, 1) \left(\frac{1}{2}\right)^1 \left(\frac{1}{2}\right)^2 = 0.375$	72
2	$C(3, 2) \left(\frac{1}{2}\right)^2 \left(\frac{1}{2}\right)^1 = 0.375$	72
3	$C(3, 3) \left(\frac{1}{2}\right)^3 = 0.125$	24

ii. When probability vary

$$\bar{x} = np \quad \text{Or, } 0.755 = 3p \quad \text{Or, } p = \frac{0.755}{3} = 0.25,$$

$$q = 1 - p = 1 - 0.25 = 0.75$$

x	$P(x) = C(n, x) p^x q^{n-x}$	Expected frequency = $N P(x)$
0	$C(3, 0) (0.25)^0 (0.75)^3 = 0.421$	$80.83 \approx 81$
1	$C(3, 1) (0.25)^1 (0.75)^2 = 0.421$	$80.83 \approx 81$
2	$C(3, 2) (0.25)^2 (0.75)^3 = 0.1406$	$26.99 \approx 27$
3	$C(3, 3) (0.25)^3 = 0.0156$	3

## 4. Fit the Poisson

Mistaken per page (x)	No. of page (f)	$\sum fx$	$P(x) = \frac{e^{-\lambda} \lambda^x}{x!}$	Expected frequency = $N P(x)$
0	142	0	$\frac{e^{-0.95} (0.25)^0}{0!} = 0.386$	154.4 $\approx$ 154
1	156	156	$\frac{e^{-0.95} (0.25)^1}{1!} = 0.367$	146.8 $\approx$ 147
2	69	138	$\frac{e^{-0.95} (0.25)^2}{2!} = 0.174$	69.6 $\approx$ 70
3	27	81	$\frac{e^{-0.95} (0.25)^3}{3!} = 0.055$	22
4	5	20	$\frac{e^{-0.95} (0.25)^4}{4!} = 0.013$	5.2 $\approx$ 5
5	1	5	$\frac{e^{-0.95} (0.25)^5}{5!} = 0.006$	2.57 $\approx$ 3
	$N = \sum f = 400$	$\sum fx = 380$		


$$\lambda = \bar{x} = \frac{\sum fx}{N} = \frac{380}{400} = 0.95$$

## 5. Fit Poisson distribution

x	f	$\sum fx$	$P(x) = \frac{e^{-\lambda} \lambda^x}{x!}$	Expected frequency = $N P(x)$
0	71	0	$\frac{e^{-1.74} (1.74)^0}{0!} = 0.175$	69.82 $\approx$ 70
1	112	112	$\frac{e^{-1.74} (1.74)^1}{1!} = 0.305$	121.695 $\approx$ 122
2	117	234	$\frac{e^{-1.74} (1.74)^2}{2!} = 0.265$	105.73 $\approx$ 106
3	57	161	$\frac{e^{-1.74} (1.74)^3}{3!} = 0.154$	61.446 $\approx$ 61
4	27	108	$\frac{e^{-1.74} (1.74)^4}{4!} = 0.067$	26.733 $\approx$ 27
5	11	55	$\frac{e^{-1.74} (1.74)^5}{5!} = 0.023$	9.177 $\approx$ 9
6	3	18	$\frac{e^{-1.74} (1.74)^6}{6!} = 0.0067$	2.67 $\approx$ 3
7	1	7	$\frac{e^{-1.74} (1.74)^7}{7!} = 0.0016$	0.67 $\approx$ 1
	$N = \sum f = 399$	$\sum fx = 695$		

$$\lambda = \bar{x}$$

$$= \frac{\sum fx}{N} = \frac{695}{399} = 1.74$$



## 14. Solution.

Age of husband (x)	Age of wife (y)	Rank of x (R <sub>1</sub> )	Rank of y (R <sub>2</sub> )	d = R <sub>1</sub> - R <sub>2</sub>	d <sup>2</sup>
23	21	1	1	0	0
27	22	2	2	0	0
28	23	3	3	0	0
29	24	4	4	0	0
30	25	5	5	0	0
31	26	6	6	0	0
33	28	7	7	0	0
35	29	8	8	0	0
				$\sum d = 0$	$\sum d^2 = 0$

$$r_s = 1 - \frac{6 \sum d^2}{n(n^2 - 1)} = 1 - \frac{6 \times 0}{6(6^2 - 1)} = 1$$

## 15. Calculate Spearman's

x	y	Rank of x (R <sub>1</sub> )	Rank of y (R <sub>2</sub> )	d = R <sub>1</sub> - R <sub>2</sub>	d <sup>2</sup>
20	52	1	3	-2	4
25	50	3	1.5	1.5	2.25
60	55	7	4	3	9
45	50	5	1.5	3.5	12.25
80	60	10	5	5	25
25	70	3	7	-4	16
55	72	6	8	-2	4
65	78	8	9	-1	1
25	80	3	10	-7	49
75	63	9	6	3	9
				$\sum d = 0$	$\sum d^2 = 131.5$

Here, m<sub>1</sub> = 3, m<sub>2</sub> = 2

$$r_s = 1 - \frac{6 \left\{ \sum d^2 + \frac{m_1(m_1^2 - 1)}{12} + \frac{m_2(m_2^2 - 1)}{12} \right\}}{n(n^2 - 1)}$$

$$= 1 - \frac{6 \left[ 131.5 + \frac{3(3^2 - 1)}{12} + \frac{2(2^2 - 1)}{12} \right]}{10(10^2 - 1)} = 1 - \frac{6 \times 134}{990} = 0.187$$

## 16.

Applicant	Marks in Skill	Marks in Ability	R <sub>x</sub>	R <sub>y</sub>	d = R <sub>x</sub> - R <sub>y</sub>
A	38	48	2.5	4	-1.5
B	41	39	8	3	2
C	68	38	8	2	6
D	41	36	5	1	4
E	38	58	2.5	5	-2.5
F	55	61	7	6.5	0.5
G	85	72	10	8	2
H	81	83	9	10	-1
I	28	61	1	6.5	-5.5
J	41	82	5	9	-4
					$\sum d = 0$

$$R = 1 - \frac{6 \left[ \sum d^2 + \frac{m_1 (m_1^2 - 1)}{12} + \frac{m_2 (m_2^2 - 1)}{12} + \frac{m_3 (m_3^2 - 1)}{12} \right]}{n (n^2 - 1)}$$

$$= 1 - \frac{6 \left[ 116 + \frac{2 \times 3}{12} + \frac{3 \times 8}{12} + \frac{2 \times 3}{12} \right]}{10 (100 - 1)}$$

$$= 1 - 0.72 = 0.278$$

17. Ten competitors in Nepal

1 <sup>st</sup> Judge (R <sub>1</sub> )	2 <sup>nd</sup> Judge (R <sub>2</sub> )	3 <sup>rd</sup> Judge (R <sub>3</sub> )	d <sub>12</sub> = R <sub>1</sub> - R <sub>2</sub>	d <sub>12</sub> <sup>2</sup>	d <sub>13</sub> = R <sub>1</sub> - R <sub>3</sub>	d <sub>13</sub> <sup>2</sup>	d <sub>23</sub> = R <sub>2</sub> - R <sub>3</sub>	d <sub>23</sub> <sup>2</sup>
1	4	6	-3	9	-5	25	-2	4
5	8	7	-3	9	-2	4	1	1
4	7	8	-3	9	-4	16	-1	1
8	6	1	2	4	7	49	5	25
9	5	5	4	16	4	16	0	0
6	9	10	-3	9	-4	16	-1	1
10	10	9	0	0	1	1	1	1
7	3	2	4	16	5	25	1	1
3	2	3	1	1	0	0	-1	1
2	1	4	1	1	-2	4	-3	9
			$\sum d_{12} = 0$	$\sum d_{12}^2 = 74$	$\sum d_{13} = 0$	$\sum d_{13}^2 = 156$	$\sum d_{23} = 0$	$\sum d_{23}^2 = 44$

$$r_{s(12)} = 1 - \frac{6 \sum d_{12}^2}{n (n^2 - 1)} = 1 - \frac{6 \times 74}{10 (10^2 - 1)} = 0.55$$

$$r_{s(13)} = 1 - \frac{6 \sum d_{13}^2}{n (n^2 - 1)} = 1 - \frac{6 \times 156}{10 (10^2 - 1)} = 0.054$$

$$r_{s(23)} = 1 - \frac{6 \sum d_{23}^2}{n (n^2 - 1)} = 1 - \frac{6 \times 44}{10 (10^2 - 1)} = 0.73$$

18. Solution.

Dell (R <sub>1</sub> )	HP (R <sub>2</sub> )	d = R <sub>1</sub> - R <sub>2</sub>	d <sup>2</sup>
5	10	-5	25
2	5	-3	9
9	1	8	64
8	3	5	25
1	8	-7	49
10	6	4	16
3	2	1	1
4	7	-3	9
6	9	-3	9
7	4	3	9
		$\sum d = 0$	$\sum d^2 = 216$

$$R = 1 - \frac{6 \sum d^2}{n (n^2 - 1)} = 1 - \frac{6 \times 216}{10 \times 99} = -0.309$$

19. The coefficient of

$$n = 10, r_s = 0.8$$

Wrong d = 7

Correct d = 9

$$r_s = 1 - \frac{6 \sum d^2}{n(n^2 - 1)}$$

$$\text{or, } 0.8 = 1 - \frac{6 \sum d^2}{10(10^2 - 1)}$$

$$\text{Or, } \frac{6 \sum d^2}{990} = 1 - 0.8$$

$$\text{Or, } \sum d^2 = \frac{0.2 \times 990}{6} = 33$$

$$\text{Correct } \sum d^2 = 33 - 7 + 9 = 35$$

$$\text{Correct } r_s = 1 - \frac{6 \text{ Correct } \sum d^2}{n(n^2 - 1)} = 1 - \frac{6 \times 35}{10(10^2 - 1)} = 0.78$$

20. R = 0.143,  $\sum d^2 = 48$ , n = ?

$$R = 1 - \frac{6 \sum d^2}{n(n^2 - 1)}$$

$$\text{or } 0.143 = \frac{6 \times 48}{n(n^2 - 1)}$$

$$\text{or } \frac{288}{n(n^2 - 1)} = 1 - 0.143$$

$$\text{or } n(n^2 - 1) = 336.05$$

$$\text{or } \text{By trial method } n = 7$$

21. Solution.

$$x = 22, \sum x = 950, \sum y = 26953, \sum y^2 = 35528893, \sum x^2 = 49250, \sum xy = 1263940$$

For  $y = a + bx$

$$\sum y = na + b \sum x$$

$$26953 = 22a + 950b \quad \dots (1)$$

$$\sum xy = a \sum x + b \sum x^2$$

$$1263940 = 950a + 49250b \quad \dots (2)$$

Solving (1) and (2)

$$a = 699.957; \quad b = 12.162$$

$$y = a + bx = 699.957 + 12.162x$$

$$\text{TSS} = \sum y^2 - n \bar{y} = 35528893 - 22 \left( \frac{26953}{22} \right)^2 = 2507792.951$$

$$\text{SSE} = \sum y^2 - a \sum y - \sum xy$$

$$= 35528893 - 699.957 \times 26953 - 12.162 \times 1263940$$

$$= 1290913.699$$

$$\text{SSR} = \text{TSS} - \text{SSE}$$

$$= 2507792.951 - 1290913.699$$

$$= 1216878.892$$

$$R^2 = \frac{\text{SSR}}{\text{TSS}} = \frac{1216878.892}{2507792.951} = 0.4852$$

x	y	$u = x - 66$	$v = y - 65$	$uv$	$u^2$	$v^2$
67	68	1	3	3	1	9
63	66	-3	1	-3	9	1
66	65	0	0	0	0	0
71	70	5	5	25	25	25
69	67	3	2	6	9	4
65	67	-1	2	-2	1	4
62	64	-4	-1	4	16	1
70	71	4	6	24	16	36
61	62	-5	-3	15	25	9
72	63	6	-2	-12	36	4
		$\Sigma u = 6$	$\Sigma v = 13$	$\Sigma uv = 60$	$\Sigma u^2 = 138$	$\Sigma v^2 = 93$

First find regression equation of v on u

$$v = a + bu$$

$$\Sigma v = na + b \Sigma u$$

$$13 = 10a + 6b \quad \dots\dots (1)$$

$$\Sigma uv = a \Sigma u + b \Sigma u^2$$

$$60 = 6a + 138b$$

$$\text{or } a = \frac{60 - 138b}{6}$$

Substitute a in (1)

$$13 = 10 \frac{60 - 138b}{6} + 6b$$

$$\text{or } 13 = \frac{600 - 1380b + 36}{6}$$

$$\text{or } 78 = 600 - 1344b$$

$$\text{or } 1344b = 600 - 78$$

$$\text{or } b = \frac{522}{1344} = 0.388$$

Substitute b in (1)

$$a = \frac{60 - 138 \times 0.388}{6} = \frac{6.401}{6} = 1.066$$

$$\therefore v = a + bu$$

$$\text{or } v = 1.066 + 0.388u$$

$$\text{or } y - 65 = 1.066 + 0.388(x - 66)$$

$$\text{or } y = 1.066 + 65 + 0.388x - 25.608$$

$$\text{or } y = 40.45 + 0.388x$$

When x = 70,

$$y = 40.45 + 0.388 \times 70 = 67.61$$

$$r = \frac{n \Sigma uv - \Sigma u \Sigma v}{\sqrt{n \Sigma u^2 - (\Sigma u)^2} \sqrt{n \Sigma v^2 - (\Sigma v)^2}} = \frac{10 \times 60 - 6 \times 13}{\sqrt{10 \times 138 - 6^2} \sqrt{10 \times 93 - 13^2}} = 0.51$$

Coefficient of determination,  $R^2 = r^2 = (0.51)^2 = 0.266$

23. Solution.

Operator	x	y	xy	$x^2$
I	16	87	1372	256
II	12	88	1056	144
III	18	89	1602	324
IV	4	68	272	16
V	3	78	234	9
VI	10	80	800	100
VII	5	75	375	25
VIII	12	83	996	144
	$\Sigma x = 80$	$\Sigma y = 648$	$\Sigma xy = 6727$	$\Sigma x^2 = 1018$

Regression equation of y on x is

$$y = a + bx$$

$$\text{Now, } \Sigma y = na + b \Sigma x$$

$$\text{or } 648 = 8a + 80b \quad \dots (1)$$

$$\Sigma xy = a \Sigma x + b \Sigma x^2$$

$$\text{or } 6727 = 80a + 1018 b$$

$$\text{or } a = \frac{6727 - 1018b}{80} \quad \dots (2)$$

Substitute 'a' in equation (1)

$$648 = 8 \times \frac{6727 - 1018b}{80} + 80b$$

$$\text{or } 648 = \frac{6727 - 1018 + 800b}{10}$$

$$\text{or } 6480 = 6727 - 218b$$

$$\text{or } b = \frac{6727 - 6480}{218} = 1.133$$

Substitute b in equation (2)

$$a = \frac{6727 - 1018 \times 1.133}{80} = 69.67$$

$$\therefore y = a + bx = 69.67 + 1.133x$$

When x = 8,

$$y = 69.67 + 1.133 \times 8 = 78.73$$

24. Solution.

The city council

x (football game)	y (minor accidents)	xy	$x^2$	$y^2$
20	6	120	400	36
30	9	270	900	81
10	4	40	100	16
12	5	60	144	25
15	7	105	225	49
25	8	200	625	64
34	9	306	1156	81
$\Sigma x = 146$	$\Sigma y = 48$	$\Sigma xy = 1101$	$\Sigma x^2 = 3550$	$\Sigma y^2 = 352$

$$b_{yx} = \frac{n \sum xy - \sum x \sum y}{n \sum x^2 - (\sum x)^2} = \frac{7 \times 1101 - 146 \times 48}{7 \times 3550 - (146)^2} = \frac{699}{3534} = 0.197$$

$$\bar{x} = \frac{\sum x}{n} = \frac{146}{7} = 20.85$$

$$\bar{y} = \frac{\sum y}{n} = \frac{48}{7} = 6.85$$

Regression equation of  $y$  on  $x$

$$\text{Or, } y - \bar{y} = b_{yx} (x - \bar{x})$$

$$\text{Or, } y - 6.85 = 0.197 (x - 20.85)$$

$$\text{Or, } y = 0.197x - 4.107 + 6.85$$

$$\therefore y = 2.742 + 0.197x \quad \text{It is in the form } y = a + bx$$

$$y = 2.742 + 0.197 \times 30 = 8.65 \approx 9$$

Hence no. of minor accidents is 9.

$$\text{Now, TSS} = \sum y^2 - n \bar{y}^2 = 352 - 7 \times (6.85)^2 = 23.542$$

$$\begin{aligned} \text{SSE} &= \sum y^2 - a \sum y - b \sum xy \\ &= 352 - 2.752 \times 48 - 0.197 \times 1101 = 3.007 \end{aligned}$$

$$\text{SSR} = \text{TSS} - \text{SSE} = 23.542 - 3.007 = 20.535$$

$$\text{Coefficient of determinants (R}^2\text{)} = \frac{\text{SSR}}{\text{TSS}} = \frac{20.535}{23.542} = 0.87$$

$$\text{Standard error of estimate} = \sqrt{\frac{\text{SSE}}{n - k - 1}} = \sqrt{\frac{3.007}{7 - 1 - 1}} = 0.77$$

25. Solution.

$x$	$y$	$xy$	$x^2$	$y^2$
27	57	1539	729	3249
45	64	2880	2025	4096
41	80	3280	1681	6400
19	46	874	361	2116
35	62	2170	1225	3844
39	72	2868	1521	5184
19	52	988	361	2704
$\Sigma x = 225$	$\Sigma y = 433$	$\Sigma xy = 74539$	$\Sigma x^2 = 7903$	$\Sigma y^2 = 27593$

Regression equation of  $y$  on  $x$

$$y = a + bx$$

$$\text{Here, } \Sigma y = na + b \Sigma x$$

$$433 = 7a + 225 b \quad \dots (1)$$

$$\Sigma xy = a \Sigma x + b \Sigma x^2$$

$$14539 = 225 a + 7903 b$$

$$a = \frac{14539 - 7903b}{225} \quad \dots (2)$$

Substitute a in equation (1)

$$433 = 7 \times \left( \frac{14539 - 7903b}{225} \right) + 225 b$$

$$\text{or } 433 = \frac{101773 - 55321b + 550625b}{225}$$

$$\text{or } 97425 = 101773 - 4696b$$

$$\text{or } b = \frac{4348}{4696} = 0.926$$

Putting the value of b in equation (2)

$$a = \frac{14539 - 7903b}{225} = \frac{14539 - 7903 \times 0.926}{225} = 32.09$$

$$y = a + bx = 32.09 + 0.926x$$

where  $x = 35$

$$\therefore y = 64.5$$

$$R^2 = \frac{\text{SSR}}{\text{SST}}$$

### 26. Solution.

x	y	$x^2$	$y^2$	xy
0	12	0	144	0
5	15	25	225	75
10	17	100	289	170
15	22	225	484	330
20	24	400	576	480
25	30	625	900	750
$\Sigma x = 75$	$\Sigma y = 120$	$\Sigma x^2 = 1375$	$\Sigma y^2 = 248$	$\Sigma xy = 1805$

To fit

$$y = a + bx$$

$$\Sigma y = na + b \Sigma x$$

$$120 = 6a + 75b \quad \dots (1)$$

$$\Sigma xy = a \Sigma x + b \Sigma x^2$$

$$1805 = 75a + 1375b$$

$$a = \frac{1805 - 1375b}{75} \quad \dots (2)$$

Substitute a in (1)

$$120 = 6 \frac{1805 - 1375b}{75} + 75b$$

$$\text{or } 120 = \frac{10830 - 8250b + 5625b}{75}$$

$$\text{or } 9000 = 10830 - 2625b$$

$$\text{or } 2625b = 10830 - 9000$$

$$\text{or } b = 0.697$$

Putting the value of b in equation (2)

$$a = \frac{1805 - 1375 \times 0.697}{75} = \frac{846.625}{75} = 11.29$$

$$y = a + bx = 11.29 + 0.697x$$

Now, coefficient of determination

$$R^2 = \frac{SSR}{TSS}$$

$$TSS = \sum y^2 - n \bar{y}^2$$

$$= 2618 - 6 \times \left(\frac{120}{6}\right)^2$$

$$= 218$$

$$SSE = \sum y^2 - a \sum y - b \sum xy$$

$$= 2618 - 11.29 \times 10 - 0.697 \times 1809$$

$$= 5.115$$

$$SSR = TSS - SSE = 218 - 5.115 = 212.88$$

$$R^2 = \frac{212.88}{218} = 0.97$$

### 27. Solution.

y	x	xy	$x^2$	$y^2$
3	50	150	2500	9
1	200	200	40000	1
4	70	280	4900	16
1	100	100	10000	1
2	90	180	8100	4
3	40	120	1600	9
$\Sigma y = 14$		$\Sigma x = 550$	$\Sigma xy = 1030$	$\Sigma x^2 = 67100$
				$\Sigma y^2 = 40$

Now,

$$\Sigma y = na + b \sum x$$

$$14 = 6a + 550b \quad \dots (1)$$

$$\Sigma xy = a \sum x + b \sum x^2$$

$$\text{or } 1030 = 550a + 67100b \quad \dots (2)$$

Solving equation (1) and (2)

$$a = 3.725, \quad b = -0.01518$$

$$y = a + bx = 3.725 - 0.01518x$$

When  $x = 110$

$$y = 3.725 - 0.01518 \times 110 = 2.055$$

$$TSS = \sum y^2 - n \bar{y}^2 = 40 - 6 \times \left(\frac{14}{6}\right)^2 = 7.333$$

$$SSE = \sum y^2 - a \sum y - b \sum xy = 40 - 3.725 \times 14 - (-0.01518) \times 1030 = 3.487$$

$$SSR = TSS - SSE = 7.333 - 3.487 = 3.846$$

$$R^2 = \frac{SSR}{TSS} = \frac{3.846}{7.333} = 0.524$$

28. Solution.

Year	Adv. exp (x)	y	xy	$x^2$	$y^2$
1	10	20	200	100	400
2	12	30	360	144	900
3	14	37	518	196	1369
4	16	50	800	256	2500
5	18	56	1008	324	3136
6	20	78	1560	400	6084
7	22	85	1958	484	7921
8	24	100	2400	576	10000
9	26	120	3120	676	14400
10	28	110	3080	784	12100
	$\Sigma x = 190$	$\Sigma y = 690$	$\Sigma xy = 15004$	$\Sigma x^2 = 3940$	$\Sigma y^2 = 58810$

$$r = \frac{n \sum xy - \sum x \sum y}{\sqrt{n \sum x^2 - (\sum x)^2} \sqrt{n \sum y^2 - (\sum y)^2}}$$

$$= \frac{10 \times 15004 - 190 \times 690}{\sqrt{10 \times 3940 - (190)^2} \sqrt{10 \times 58810 - (690)^2}} = 0.985$$

For  $y = a + bx$ 

$$\Sigma y = na + b \Sigma x$$

$$690 = 10a + 190b \quad \dots (1)$$

$$\Sigma xy = a \Sigma x + b \Sigma x^2$$

$$15004 = 190a + 3940b \quad \dots (2)$$

Solving equation (1) and (2)

$$a = -40.048$$

$$b = 5.739$$

$$\therefore y = a + bx = -40.048 + 5.739x$$

When  $x = 27$ ,

$$y = -40.048 + 5.739 \times 27 = 114.915$$

29. Solution.

x	y	$x^2$	$y^2$	xy
250	83	62500	6889	20750
340	89	115600	7921	30260
320	88	102400	7744	28160
330	89	108900	7921	29370
346	92	119716	8464	31832
260	85	67600	7225	22100
280	84	78400	7056	23520
395	92	156025	8464	36340
380	93	144400	8649	35340
400	96	160000	9216	38400
$\Sigma x = 3301$	$\Sigma y = 891$	$\Sigma x^2 = 1115541$	$\Sigma y^2 = 79549$	$\Sigma xy = 296072$

$$\begin{aligned}
 r &= \frac{n \sum xy - \sum x \sum y}{\sqrt{n \sum x^2 - (\sum x)^2} \sqrt{n \sum y^2 - (\sum y)^2}} \\
 &= \frac{10 \times 296072 - 3301 \times 891}{\sqrt{10 \times 1115541 - (3301)^2} \sqrt{10 \times 79549 - (891)^2}} \\
 &= 0.957
 \end{aligned}$$

Equation,  $y = a + bx$

$$\sum y = na + b \sum x$$

$$\text{or } 891 = 10a + 3301 b \quad \dots (1)$$

$$\sum xy = a \sum x + b \sum x^2$$

$$\text{or } 296072 = 3301 a + 1115541 b \quad \dots (2)$$

Solving (1) and (2)

$$a = 64.1915; b = 0.075$$

$$\therefore y = a + bx = 64.1915 + 0.075x$$

### 30. Solution.

Data size (x)	Processed request (y)	$x^2$	$y^2$	$xy$
6	40	36	1600	240
7	55	49	3025	385
7	50	49	2500	350
8	41	64	1681	328
10	17	100	289	170
10	26	100	676	260
15	16	225	256	240
$\sum x = 63$		$\sum y = 245$	$\sum x^2 = 623$	$\sum y^2 = 10027$
				$\sum xy = 1973$

To find  $y = a + bx$

$$\sum y = na + b \sum x$$

$$245 = 7a + 63b \quad \dots (1)$$

$$\sum xy = a \sum x + b$$

$$1973 = 63a + 623b \quad \dots (2)$$

Solving (1) and (2)  $a = 72.278, b = -4.142$

$$y = a + bx = 72.278 - 4.142x$$

$$\text{TSS} = \sum y^2 - n \bar{y}^2 = 10027 - 7 \times \left( \frac{245}{7} \right)^2 =$$

$$\text{SEE} = \sum y^2 - a \sum y - b \sum xy = 10027 - 72.278 \times 245 - (-4.142) \times 1973 =$$

$$\text{SSR} = \text{TSS} - \text{SEE} =$$

$$R^2 = \frac{\text{SSR}}{\text{TSS}} = 0.662$$

$$\text{When } x = 12, y = 22.57$$