



JEE (Main)

PAPER-1 (B.E./B. TECH.)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

2021

COMPUTER BASED TEST (CBT) Memory Based Questions & Solutions

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Duration: 3 Hours | Max. Marks: 300

SUBJECT: PHYSICS

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/resonancewatch blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



| JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

PART : PHYSICS

1. A deuteron & α -particle both enters in a region of magnetic field perpendicular to it with same kinetic energy find the ratio of their radii ?

(1) 2

(2) $2\sqrt{2}$

(3) $\sqrt{2}$

(4) $\frac{1}{2}$

Ans. (3)


Sol. $r = \frac{mv}{qB} = \frac{\sqrt{2mK}}{qB}$

$$r \propto \frac{\sqrt{m}}{q}$$

$$m_a = 2m_d$$

$$\frac{r_d}{r_a} = \frac{\sqrt{m_d}}{q_d} \times \frac{2q_d}{\sqrt{2m_d}} = \sqrt{2}$$

2. In the given arrangement, spring of spring constant 100 N/m is compressed by 0.5m. The height of the arrangement is 2m. A basket is placed at distance d such that after projection, ball will fall in the basket. If the mass of the ball is 100 gm, find maximum value of d?



(A) 5 m (B) 10 m (C) 15 m (D) 20 m

Sol. By energy conservation

$$\frac{1}{2}kx^2 = \frac{1}{2}mv^2 \Rightarrow v = \sqrt{\frac{k}{m}} \quad v = 0.5 \times \sqrt{\frac{100}{0.1}} = 5\sqrt{10} \text{ m/s}$$

$$\text{Time of flight of ball } T = \sqrt{\frac{2H}{g}} = \sqrt{\frac{2 \times 2}{10}} = \frac{2}{\sqrt{10}} \text{ sec}$$

Range of ball $s = ut$

$$d = 5\sqrt{10} \times \left(\frac{2}{\sqrt{10}} \right) = 10 \text{ m}$$

Resonance Eduventures Ltd.


Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029
Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/resonancewatch blog.resonance.ac.in

Resonance® Educating for better tomorrow | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

3. When a disc slides on smooth inclined surface from rest, the time taken to move from A to B is t_1 . When disc performs pure rolling from rest then time taken to move from A to B is t_2 . If $\frac{t_2}{t_1} = \sqrt{\frac{3}{x}}$ find x.



B

Ans. 2

Sol. When disc slides $a_1 = g \sin \theta$ So $S = ut_1 + \frac{1}{2} a_1 t_1^2 = \frac{1}{2} g \sin \theta \cdot t_1^2 \quad \dots(1)$

$$\text{When disc do pure rolling } a_2 = \frac{gsin\theta}{1+k^2/R^2} = \frac{gsin\theta}{1+1/2} = \frac{2}{3} g \sin \theta$$

$$\text{So } S = ut_2 + \frac{1}{2} a_2 t_2^2 = \frac{1}{2} \cdot \frac{2}{3} g \sin \theta \cdot t_2^2 \quad \dots(2)$$

From (1) & (2)

$$\frac{t_2}{t_1} = \sqrt{\frac{3}{2}}$$

4. We have a charge of magnitude Q. If we divide charge in two parts, what should be their ratio so that there will be max repulsion force between them?

(1) 1 : 1 (2) 2 : 1 (3) 1 : 2 (4) 3 : 2

Ans. (1)


Sol.

+q

Q-q

$$\begin{aligned} dF &= 0 \\ \frac{d}{dq} &= 0 \\ Q - 2q &= 0 \\ q &= Q/2 \\ \text{ratio} &= 1 : 1 \end{aligned}$$


5. Four planks are arranged in a lift going upwards with an acceleration of 0.2 m/s^2 as shown in figure. Find the normal reaction applied by the lift on a 10 kg block : ($g = 9.8 \text{ m/s}^2$)



- Ans. (1) 500 (2) 700 (3) 672 (4) 800




Sol.



$$\begin{aligned} N - 70g &= 70 \times 0.2 \\ N &= 70(g + 0.2) \\ N &= 700 \end{aligned}$$

6. For given PV curve, Find net heat taken by gas system in cyclic process.



- Ans. (1) 25π (2) 50π (3) 75π (4) 100π


$$\begin{aligned} &= \left[\frac{40-20}{2} \times 10^3 \right] \times \left[\frac{40-20}{2} \times 10^{-3} \right] \\ &= 100\pi \text{ Joule} \end{aligned}$$

7. A butterfly is flying in North-East with $4\sqrt{2}$ m/s w.r.t. wind. Wind is blowing at 1 m/s southwards. Displacement of butterfly in 3s is

- (1) 10 meter (2) 15 meter (3) 20 meter (4) 5m

Ans. (2)

Sol.



$$\vec{D} = v_{F,G} \times T$$

$D = 15\text{m}$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005


Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029
 Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/Resonance blog.resonance.ac.in


This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 3

8. In the given circuit, find current passing through 6Ω resistor



Ans.
 Sol.



Let potential at junction point = x

By KCL $\sum i_{in} = 0$

$$\Rightarrow \frac{160-x}{4} + \frac{90-x}{5} + \frac{0-x}{6} = 0$$



$$\Rightarrow \frac{160 \times 15 - 15x + 90 \times 12 - 12x + 0 - 10x}{60} = 0$$

$$\Rightarrow 37x = 2400 + 1080 \\ x = 94.05$$

So current $i_3 = \frac{x}{6}$

$$= \frac{94.05}{6} = 15.67$$

9. In the given system, uniform magnetic field exists from $x = 0$ to $x = b$. A rod is first moved from $x = 0$ to $x = 2b$ uniformly and then moved reverse uniformly from $x = 2b$ to $x = 0$. Match the quantities with proper curves



Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029
 Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/Resonance blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 4

Column-I

(a) Flux (ϕ)

(b) EMF (e)

(c) Power (P)

- (1) (a) – (iii), (b) – (i), (c) – (ii)
 (3) (a) – (iii), (b) – (ii), (c) – (i)

Ans.

Sol.

$$\text{Flux} = \phi = B \cdot A$$

$$\Rightarrow B \propto A \cos \theta$$

$$\text{Where } A = vt$$


$$\phi = B \cdot vt$$

One rod go at $x > b$ then ϕ stop changing this constant flux = B/b .

When rod come back and when $x < b$ flux start decreasing so graph ϕ /st



Column-II




b → (ii)

$$e = -\frac{d\phi}{dt}$$

e = – slope of ϕ – t graph

In $0-t_1$ graph slope +ve and constant so e = negative and zero.

in t_2-t_3 slope of ϕ – t is negative and constant so e = positive and zero



Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005


Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more, visit [RESCO](#) at [E6677](#) | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

$$\text{Power} = \frac{e^2}{R}$$

Resistance is only of rod so R of the circuit is constant

$$P = \frac{B^2 l^2 V^2}{R} = \text{constant}$$



10. A uniform rod of young's modulus Y is stretched by two tension T_1 and T_2 such that rods get expanded to length L_1 and L_2 respectively. Find initial length of rod?

$$(1) \frac{L_1 T_1 - L_2 T_2}{T_1 - T_2} \quad (2) \frac{L_2 T_1 - L_1 T_2}{T_2 - T_1} \quad (3) \frac{L_1 T_2 - L_2 T_1}{T_2 - T_1} \quad (4) \frac{L_1 \times T_2}{T_1 \times L_2}$$

Ans. (3)

Sol. Let initial length of rod be L_0 and Area A.

$$\text{As } \frac{T}{A} = Y \frac{\Delta\ell}{\ell}$$

$$\text{So, } \frac{T_1}{A} = \frac{Y(L_1 - L_0)}{L_0}$$

$$\frac{T_2}{A} = \frac{Y(L_2 - L_0)}{L_0}$$

Dividing

$$\frac{T_1}{T_2} = \frac{L_1 - L_0}{L_2 - L_0}; \quad T_1 L_2 - T_1 L_0 = T_2 L_1 - T_2 L_0; \quad L_0 = \frac{L_1 T_2 - L_2 T_1}{T_2 - T_1}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$(1) \sqrt{A^2 + B^2 - \sqrt{2}AB}$$

$$(2) \sqrt{A^2 + B^2 + \sqrt{2}AB}$$

$$(3) A - B$$

$$(4) A + B$$

Ans. (1)

Sol. $\vec{A} \cdot \vec{B} = |\vec{A} \times \vec{B}|$

$$\Rightarrow AB \cos \theta = AB \sin \theta$$

$$\therefore \theta = 45^\circ$$

$$\therefore |\vec{A} - \vec{B}| = \sqrt{A^2 + B^2 - 2AB \cos 45^\circ}$$

$$= \sqrt{A^2 + B^2 - \sqrt{2}AB}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [facebook.com/ResonanceEdu](#) | [twitter.com/ResonanceEdu](#) | [www.youtube.com/Resonance](#) | [blog.resonance.ac.in](#)

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 6


Resonance® | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

12. In the L-C-R series A.C. circuit shown below, current leads source voltage by 45° . Find capacitance of the capacitor.

$$(1) 2.1 \text{ mF} \quad (2) 3.33 \text{ mF} \quad (3) 4.3 \text{ mF} \quad (4) 5.1 \text{ mF}$$

Ans.

Sol.



$$\tan 45^\circ = \frac{V_{0C} - V_{0L}}{V_{0R}}$$

$$\therefore C = \frac{1}{\omega C}$$

$$0.1\Omega = \frac{1}{300C} - 3 \times 10^{-3} \times 300$$

$$\therefore C = 3.33 \text{ mF}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- (1) 36.67 m/s (2) 30.12 m/s (3) 22.37 m/s (4) 20.25 m/s
Ans. (1)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005
 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222
 To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029
 Toll Free : 1800 258 5555     www.youtube.com/resonancewatch blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 8

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



| JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS


Sol. Frequency received by wall $f' = \left(\frac{v_s}{v_s - v} \right) f_0$

Reflected frequency received by man is $f'' = \left(\frac{v_s + v}{v_s} \right) f'$

$$\Rightarrow f'' = \left(\frac{v_s + v}{v_s} \right) \left(\frac{v_s}{v_s - v} \right) f_0 \Rightarrow f'' = \left(\frac{v_s + v}{v_s - v} \right) f_0 \Rightarrow 500 = \left(\frac{330 + v}{330 - v} \right) 400$$

$$\Rightarrow v = \frac{330}{9} = 36.67 \text{ m/s}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029
 material is 44×10^{-2} ohm \times m. Find electric field inside the rod?



(1) 0.567

(2) 0.367

(3) 0.667

(4) 0.767

Ans. (2)

Sol. $J = \sigma E$

$$\frac{I}{A_{\text{effective}}} = \frac{E}{\rho}$$

$$E = \frac{\rho I}{A} \cos 60^\circ = \frac{44 \times 10^{-2} \times 5}{3 \times 2}; E = 0.367$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029
 by it is zero.

(1) 700 R

(2) 600 R

(3) 500 R

(4) 100 R

Ans. (3)

Sol. $n = 4$

$\Delta T = 50K$

$$C_v = \frac{5R}{2}$$

As $W = 0$. It means isochoric process

$Q = \Delta U$

$$= nC_v \Delta T = 4 \times \frac{5R}{2} \times 50 = 500 R$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Toll Free : 1800 258 5555     www.youtube.com/resonancewatch blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal


PAGE # 9

19. A car is moving on a Banked rough road, the mass of car is 800 kg. The angle of Banking is 30° , car is moving with maximum speed given that $\mu_s = 0.2$. find the Normal Reaction (in Newton)?
 (1) 24000 (2) 5000 (3) 10000 (4) 9000

Ans. (3)

Sol.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



Perpendicular to inclined plane

$$N = mg \cos 30^\circ + \frac{mv^2}{R} \sin 30^\circ$$

$$N - mg \cos 30^\circ = \frac{mv^2}{R} \sin 30^\circ \quad \dots(1)$$

along inclined plane

$$mg \sin 30^\circ + \mu_s N = \frac{mv^2}{R} \cos 30^\circ \quad \dots(2)$$


Dividing (1) by (2)

$$N - mg \cos 30^\circ = \frac{mg \sin 30^\circ + \mu_s N}{\frac{mv^2}{R} \cos 30^\circ}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029
 Solving $N = 10000$ (Approx)

20. A particle of mass m moving with speed v collide elastically with the end of a uniform rod of mass M and length L perpendicularly as shown in figure. If the particle comes to rest after collision find the value of

$$\frac{m}{M}$$



- (1) 1/3 (2) 1/2 (3) 1/4 (4) 1/5

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-3916722

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029
Toll Free : 1800 258 5555


This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 10

Sol.

Rest

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029




Conservation of angular momentum about centre of mass of rod

$$\frac{1}{2} M L^2$$

$$\begin{aligned}
 mv \left(\frac{\omega}{2} \right) &= \frac{mv\omega}{12} \quad \dots(i) \\
 mv = Mv_1 &\dots(ii) \\
 1 &= \frac{v_1 + \omega L}{v} \quad \dots(iii) \\
 \text{Putting } v_1 \text{ from (ii) and } \omega L \text{ from (i) in (iii)} \\
 v &= \frac{m}{M} v + \frac{6mv}{2M} \\
 &= 4m \quad \dots(iv)
 \end{aligned}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

21. An object is moved from earth to moon. Choose the correct weight vs distance curve. Gravitational acceleration on earth surface is 10 m/s^2 and that on moon is 4 m/s^2 . Mass of the object is 1kg.



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



Ans. (2)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555

7340010333



facebook.com/ResonanceEdu



twitter.com/ResonanceEdu



www.youtube.com/Resonance

blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 11

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

 | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

Sol. \vec{g} (at any point) = $\vec{g}_{\text{Earth}} + \vec{g}_{\text{moon}}$. Since distance is large so $|\vec{g}| = |\vec{g}_E| = 10$.

As we move away from earth, It decrease to zero at a point where $\vec{g}_E + \vec{g}_M = 0$

Then it increase to $|\vec{g}| = |\vec{g}_M| = 4$ at moon surface.

22. For the spherical interface of radius of curvature $R = 30 \text{ cm}$ shown in figure. The two different media having refractive indices $n_1 = 1.4$ and $n_2 = 1.25$, an object is placed at 40 cm from the interface as shown in figure. Find position of image.



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

(1) 41.67

(2) 35.42

(3) 22.27

(4) 15.25

Ans. (1)

$$\frac{n_2}{v} - \frac{n_1}{u} = \frac{n_2 - n_1}{R}$$

$$\Rightarrow \frac{1.25}{v} - \frac{1.4}{-40} = \frac{1.25 - 1.4}{(-30)}$$


$$\Rightarrow \frac{1.25}{v} = 0.005 - 0.035$$

$$\Rightarrow v = -41.67 \text{ cm}$$

23. A ball of charge to mass ratio $8\mu\text{C/g}$ is placed at a distance of 10 cm from a wall. An electric field 100 N/m is switched on in the direction of wall. Find time period of its oscillations? Assume all collisions elastic.



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



(1) 1 sec

(2) 2 sec

(3) 3 sec

(4) 4 sec.

Ans. (1)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029
Toll Free : 1800 258 5555 7340010333

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 12

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



| JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

$$\text{Sol. } a = \frac{qE}{m} = \frac{8 \times 10^{-6}}{10^{-3}} \times 100 = 0.8 \text{ m/s}^2$$

As electric field is switched on, ball first strikes to wall and returns back.
one oscillation.

$$\text{Thus } s = ut + \frac{1}{2}at_1^2$$

$$0.1 = \frac{1}{2} \times 0.8t_1^2$$

1

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\text{Thus time period } T = 2 \times \frac{1}{2} = 1 \text{ sec.}$$

24. A body of mass m emits a photon of frequency ν , then loss in its internal energy ?

(1) $h\nu$


$$(2) h\nu \left(1 - \frac{h\nu}{2mc^2}\right)$$

$$(3) h\nu \left(1 + \frac{h\nu}{2mc^2}\right)$$

(4) zero

Ans. (3)

Sol.



$$mv = \frac{h}{\lambda} = \left(\frac{h\nu}{c}\right)$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$= \frac{1}{2} \frac{p^2}{m} + h\nu$$

$$= \frac{1}{2m} \left(\frac{h\nu}{c}\right)^2 + h\nu$$

$$= h\nu \left(1 + \frac{h\nu}{2mc^2}\right)$$

Resonance Eduventures Ltd.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029
Toll Free : 1800 258 5555     

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 13



| JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

25. Consider an equation $S = \alpha^2 \beta / n \left(\frac{n k R}{\beta^2} - 1 \right)$

Where S = Entropy

n = No. of moles

k = Boltzmann constant

R = Universal gas constant

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- (1) $[M^0 L^0 T^0]$, $[M^1 L^2 T^2 K^{-1}]$
 (2) $[M^1 L^2 T^{-2}]$, $[M^1 L^2 T^{-2} K^{-1}]$
 (3) $[M^1 L^2 T^{-2} K^{-1}]$, $[M^0 L^0 T^0]$
 (4) None of these

Ans. (1)

Sol. $S = \frac{Q}{\Delta T}$

$$[S] = \frac{ML^2 T^{-2}}{K}$$

$$K = \frac{\text{Energy}}{T}$$

$$[K] = [S] = \frac{ML^2 T^{-2}}{K}$$

$$[R] = \left[\frac{\text{Energy}}{nT} \right] = \frac{ML^2 T^{-2}}{mol K}$$

$$[J] = M^0 L^0 T^0$$

$$\text{Now, } [nKR] = [J\beta^2]$$

$$(mol) \times \frac{ML^2 T^{-2}}{K} \times \frac{ML^2 T^{-2}}{mol K} = [\beta^2]$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$[\alpha^2] = \left[\frac{\alpha}{\beta} \right] = \frac{M^0 L^0 T^0}{K \times M^1 L^2 T^{-2} K^{-1}} ; \quad \alpha = M^0 L^0 T^0$$

26. The shape of travelling wave at $t = 0$, is given by $y = \frac{1}{1+x^2}$. If after 3 sec shape of the wave pulse is

represented by $y = \frac{1}{1+(1-x)^2}$, then speed of wave is :

- (1) $\frac{1}{2} \text{ m/s}$ (2) $\frac{4}{3} \text{ m/s}$ (3) $\frac{1}{3} \text{ m/s}$ (4) $\frac{5}{6} \text{ m/s}$

Ans. (3)

Sol. $x \rightarrow (x - vt)$

$$y = \frac{1}{1+(x-vt)^2}$$

$$\text{At } t = 0 ; \quad y = \frac{1}{1+x^2}$$

$$\text{at } t = 3 ; \quad y = \frac{1}{1+(x-3v)^2}$$

By comparing

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029
Toll Free : 1800 258 5555     

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 14

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

incident on a gold plate from which electron is emitted which will make a radius of 7 mm in a uniform magnetic field of intensity 5×10^{-4} T find the work function of gold plate?

- (1) 3.4 eV (2) 5.12 eV (3) 1.031 eV (4) 11.01 eV

Ans. (4)

$$\begin{aligned} \text{Sol. } E_p &= 13.6 \left[\frac{1}{R_1^2} - \frac{1}{R_2^2} \right] \text{eV} \\ &= 13.6 \left[\frac{1}{1} - \frac{1}{9} \right] \\ E_p &= 12.08 \text{ eV} \\ \text{For Gold plate} \\ \phi &= E_p - K_{\text{Emax}} \\ v &= \frac{RqB}{m} \end{aligned}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\begin{aligned} K.E. &= \frac{1}{2} m V^2 \\ K.E. &= \frac{1}{2} \times \frac{9.1 \times 10^{-31} \times (6.15 \times 10^5)^2}{1.6 \times 10^{-19}} \text{ eV} = 1.075 \text{ eV} \\ \phi &= 12.05 - 1.075 \\ \phi &= 11.01 \text{ eV} \end{aligned}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555  facebook.com/ResonanceEdu  twitter.com/ResonanceEdu  www.youtube.com/resonancewatch  blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 15

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



RESULT: JEE (Advanced),
JEE (Main), NEET

HIGHEST No. of Classroom Selections

in JEE (Advanced) 2020 from any Institute of Kota

5 AIRs in TOP-50 in JEE (Adv.) 2020 from Classroom

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



Total Selections in JEE (Advanced) 2020

1505 | **14755** | **2646**

Eligible for JEE (Advanced) Through JEE (Main) 2020

NEET 2020

ADMISSION OPEN for Session 2021-22

ONLINE + OFFLINE PROGRAMS

CLASS 11, 12 & 12+

Target: JEE (Main+Adv.) | JEE (Main) | NEET

Scholarship Upto 90%*

Toll Free: 1800 258 5555 | Visit us: www.resonance.ac.in

Terms & Condition

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



JEE (Main)

PAPER-1 (B.E./B. TECH.)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

2021

COMPUTER BASED TEST (CBT) Memory Based Questions & Solutions

Date: 20 July, 2021 (SHIFT-1) | TIME : (9.00 a.m. to 12.00 p.m)

Duration: 3 Hours | Max. Marks: 300

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | facebook.com/ResonanceEdu | twitter.com/ResonanceEdu | www.youtube.com/resonancewatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal



Resonance®

RCSIIIT JEE (Advanced),

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

≡ HIGHEST No. of Classroom Selections ≡

in JEE (Advanced) 2020 from any Institute of Kota

5 AIRs in TOP-50 in JEE (Adv.) 2020 from Classroom



AIR-2
(GEN-EWS)
AIR-15



Zonal Topper
IIT-Kharagpur
AIR-25



2nd Rank in
IIT-Kharagpur Zone
AIR-29



AIR-30



AIR-41

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



With us Since Class 9th



With us Since Class 11th



With us Since Class 11th



With us Since Class 9th



With us Since Class 10th

Total Selections in JEE (Advanced) 2020

4505

Classroom: 3441 | Distance: 1064

Eligible for JEE (Advanced) Through JEE (Main) 2020

14755

Classroom: 11047 | Distance: 3708

NEET 2020

2646

Classroom: 1823 | Distance: 811

ADMISSION OPEN for Session 2021-22

ONLINE + OFFLINE PROGRAMS


CLASS 11, 12 & 12+

Target: JEE (Main+Adv.) | JEE (Main) | NEET

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Scholarship Upto 90%*

Toll Free: 1800 258 5555 | Visit us: www.resonance.ac.in



Print & Download



| JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | CHEMISTRY

PART : CHEMISTRY

1. Arrange the following complex in increasing order of intensity of colour.

$\text{Co}(\text{CN})_6^{3-}$, $\text{Co}(\text{Cl})_6^{2-}$, $\text{Co}(\text{H}_2\text{O})_6^{2+}$

$\text{Co}(\text{H}_2\text{O})_6^{2+}$, $\text{Co}(\text{CN})_6^{3-}$, $\text{Co}(\text{Cl})_6^{2-}$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

(3) $[\text{Co}(\text{CN})_6]^{3-}$, $[\text{Co}(\text{Cl})_6]^{2-}$, $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$ (4) $\text{Co}(\text{H}_2\text{O})_6^{2+}$, $[\text{Co}(\text{CN})_6]^{3-}$, $[\text{Co}(\text{Cl})_6]^{2-}$

Sol. (1)

	Complex	Colour
1.	$[\text{Co}(\text{CN})_6]^{3-}$,	Yellow
2.	$[\text{Co}(\text{H}_2\text{O})_6]^{2+}$	Pink
3.	$[\text{Co}(\text{Cl})_6]^{2-}$	Blue

2. Which of the following does not disproportionate

(1) BrO^-

(2) BrO_2^-

(3) BrO_3^-

(4) BrO_4^{\ominus}

Ans. (4)

Sol. In BrO_4^- , Br is in maximum oxidation state. So it can only reduce

3. A metal M on reaction with excess oxygen give MO_2 type oxide (as main product) then possible metal is.

(1) Li

(2) Na

(3) K

(4) Mg

Ans. (3)

Potassium on reaction with excess oxygen give superoxide

4. Identify the correct increasing order of 1st ionisation energy of following

Al, Mg, Si, S, P

(1) Mg, Al, Si, P, S (2) Al, Mg, Si, S, P (3) Mg, Al, Si, S, P (4) Al, Mg, Si, P, S

Ans. (2)

Sol. Mg 3s²

Al 3s²3p¹

Si 3s²3p²

P 3s²3p³

S 3s²3p⁴

So correct order : - Al < Mg < Si < S < P

Resonance Eduventures I ltd



JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | CHEMISTRY

5. Four moles of a diatomic gas is heated from 0°C to 50°C, find the heat supplied to the gas if work done by it is zero.


(1) 780 R (2) 500 R (3) 100 R (4) 650 R

Ans. (2)

Sol. w = 0

$$\Delta E = q_v = nC_v\Delta T$$

6. HNO₃ on reaction with P₄O₁₀ gives an oxide 'A'



Nature of oxide A is

- (1) acidic (2) Basic (3) Neutral (4) Amphoteric

Sol. P₄O₁₀ + 4HNO₃ → 2N₂O₅ + 4HPO₃

'A'

Nature of oxide 'A' is "acidic".

7. An equimolar mixture of benzene (P⁰_{Benzene} = 70 torr and methyl benzene (P⁰_{Methyl Benzene} = 20 torr) is prepared, then find mole fraction of benzene in vapour phase.

Ans. 0.7

$$P_{\text{Total}} = P_{\text{Benzene}}^0 \times X_{\text{Benzene}} + P_{\text{Toluene}}^0 \times X_{\text{Toluene}}$$

$$= (70) \frac{1}{2} + (20) \frac{1}{2}$$

$$= 35 + 10$$

$$= 45$$

$$1 \quad Y_{\text{Benzene}} \quad Y_{\text{Toluene}}$$

$$\frac{1}{45} = \frac{Y_{\text{Benzene}}}{70} + \frac{1 - Y_{\text{Benzene}}}{20}$$

$$\frac{1}{45} = \frac{2Y_{\text{Benzene}} + 7(1 - Y_{\text{Benzene}})}{140}$$

$$\frac{1}{45} = \frac{2Y_{\text{Benzene}} + 7 - 7Y_{\text{Benzene}}}{140}$$

$$\frac{1}{45} = \frac{7 - 5Y_{\text{Benzene}}}{140}$$

$$\frac{1}{9} = \frac{7 - 5Y_{\text{Benzene}}}{28}$$

13. 4.5 gram mass of a substance [molar mass = 90 g/mol] is dissolved in 250 ml solution, then molarity of solution is -

Ans. 0.2

Sol. Molarity (M) = $\frac{W_{\text{solute}} \times 1000}{GMM_{\text{solute}} \times V_{\text{sol}}}$

$$M = \frac{4.5 \times 1000}{90 \times 250} = \frac{4.5 \times 4}{90} = 0.2 \text{ M.}$$

14. What is the magnetic moment (Spin only) of complex $[\text{Co}(\text{CN})_6]^{4-}$?
[Report your answer to nearest integer]

So number of unpaired electrons = 1.

$$\mu = \sqrt{n(n+2)} = \sqrt{3}$$

$$\mu = 1.73 \text{ BM} \approx 2 \text{ BM.}$$

15. 10000 KJ energy is needed per day, if heat of combustion of glucose is 2700 KJ/Mole. Then how many gram of glucose is needed per day for this : [Report your answer to nearest integer].

Ans. 667



No. of mole of glucose require for production of 10,000 KJ heat is = $\frac{10,000}{2700}$ mole.

Total mass of glucose = $\frac{10,000}{2700} \times 180 = 666.67 \text{ gram.}$

16. The value of ℓ (azimuthal quantum number) for valence shell electron of Ga^+ ion is _____.
[Report your answer to nearest integer]

Ans. 0

Azimuthal Quantum number (ℓ) for valence shell electron is 0.

17. What is the difference in energy between 2nd and 3rd orbit of He^+ ion (in eV) is -
[Report your answer to nearest integer]

Ans. 8

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 7340010333 [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) [youtube.com/resonancewatch](https://www.youtube.com/resonancewatch) blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 4

Sol. $(E_{\text{He}^+})_{n=2} = -13.6 \times \frac{(2)^2}{(2)^2} = -13.6 \text{ ev}$

$$(E_{\text{He}^+})_{n=3} = -13.6 \times \frac{(2)^2}{(3)^2} = -13.6 \times \frac{4}{9}$$

$$(E_{\text{He}^+})_{n=3} - (E_{\text{He}^+})_{n=2} = 13.6 \left[1 - \frac{4}{9} \right]$$

$$= 13.6 \left[\frac{5}{9} \right] = 7.55 \text{ eV}$$

18. Anion of a compound 'x' gives brown ring test and cation gives deep blue coloration with NH_4OH and also gives precipitate with $\text{HCl} & \text{H}_2\text{S}$, then compound 'x' is

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol. Nitrates give brown ring test.



Deep Blue



Black

19. What is the value of second excitation energy of Li^{2+}

- (1) 108.8 eV (2) 81.6 eV (3) 13.6 eV (4) 95.2 eV

Ans. (1)



$$|\Delta E_{\text{u}^{2+}}|_{1 \rightarrow 3} = 13.6 \times 2^2 \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$= 13.6 \times 9 \left[\frac{8}{9} \right]$$

$$= 13.6 \times 8 = 108.8 \text{ eV}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555

7340010333

[facebook.com/ResonanceEdu](#)

[twitter.com/ResonanceEdu](#)

[www.youtube.com/resonance](#)


[blog.resonance.ac.in](#)

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 5

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

 | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | CHEMISTRY



Identify compound P and Q respectively





- (1) Primary amine, secondary amine (2) Primary amine, Primary amine
(3) Primary amine, Isonitrile (4) Secondary amine, Isonitrile

Ans. (3)

Sol. Only 1° amines give carbylaminies reaction



21. In which of the following carbocation are resonance stabilised



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029


- (1) A, B

- (2) A, C

- (3) A, D


- (4) A, B, C

Ans. (3)




are resonance stabilised carbocations.

Ans. (2)



Sol.



-C -CH₂CH₃

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029





Toll Free : 1800 258 5555 **7340010333** facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/resonancewatch blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal


PAGE # 6

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

23. Ninhydrin + $\text{NH}_2\text{-CH-}^{\text{R}}\text{-COOH}$ $\xrightarrow{\alpha\text{-amino acid}}$ Product is ?



Ans. (1)



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Ninhydrin

Deep Blue
Rube mann's purple

Ninhydrin is useful for identification of α -amino acid which react with ninhydrin and give deep blue colour.


24.

$\text{Cyclohexene} \xrightarrow[\text{H}_2\text{SO}_4 \text{ } \Delta]{\text{KMnO}_4} \text{A}$
 $\text{Cyclohexene} \xrightarrow[\text{H}_2\text{O}_2 / 273\text{ K}]{\text{KMnO}_4} \text{B}$


A, B are respectively

- (1) both diol
 - (2) both dicarboxylic acid
 - (3) A is dicarboxylic acid and B is diol
 - (4) A is diol and B is dicarboxylic acid

Ans (3)



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005
 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222
 To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029
 Toll Free : 1800 258 5555 7340010333 www.youtube.com/resonancewatch blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 7




| JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | CHEMISTRY

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

(1) Polyamide (2) Polyester (3) Polyacrylonitrile (4) Polycarbonate

Ans. (3)

Sol. Orlon is a polymer of acrylonitrile also known as PAN



Acrylonitrile

PAN or orlon

26. Which of the following is better for green chemistry in day to day life (Domestic Purpose)

(1) $\text{Cl}_2\text{C} = \text{CCl}_2$ as dry cleaning agent liquid (2) Liquid CO_2 for cloth cleaning
 (2) Cl_2 gas a bleaching agent of paper (4) CCl_4 as dry cleaning agent

Ans. (2)

Sol. $\text{CCl}_2 = \text{CCl}_2$ was earlier used as solvent for dry cleaning agent but it is carcinogen. So liquid CO_2 is used. Replacement of halogenated solvent by liquid CO_2 will result in less harm to ground water.


27. Which of the following incorrect :

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Ans. (1)

28. How many mole of CH_3MgBr are required to convert ethylethanoate to 2-methylpropan-2-ol :

Ans. (2)



Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005
 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029
 Toll Free : 1800 258 5555 7340010333 www.youtube.com/resonancewatch blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 8



JEE (Main)

PAPER-1 (B.E./B. TECH.)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

2021

COMPUTER BASED TEST (CBT) Memory Based Questions & Solutions

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Duration: 3 Hours | Max. Marks: 300

SUBJECT: MATHEMATICS

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 7340010333 [facebook.com/ResonanceEdu](#) [twitter.com/ResonanceEdu](#) [www.youtube.com/resonancewatch](#) [blog.resonance.ac.in](#)

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



RESULT: JEE (Advanced),
JEE (Main), NEET

HIGHEST No. of Classroom Selections

in JEE (Advanced) 2020 from any Institute of Kota

5 AIRs in TOP-50 in JEE (Adv.) 2020 from Classroom

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



Total Selections in JEE (Advanced) 2020

4505

Eligible for JEE (Advanced) Through JEE (Main) 2020

14755

NEET 2020

2646

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

ADMISSION OPEN for Session 2021-22

ONLINE + OFFLINE PROGRAMS

CLASS 11, 12 & 12+

Target: JEE (Main+Adv.) | JEE (Main) | NEET

Scholarship Upto 90%*

*T&C's Apply

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



| JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS

PART : MATHEMATICS

- 1 All possible words with or without meaning were formed using all the letters of the word 'EXAMINATION'.
The probability that 'M' appears at fourth position is :

$$(1) \frac{2}{11} \quad (2) \frac{1}{11} \quad (3) \frac{4}{11} \quad (4) \frac{8}{11}$$

Ans. (2)

Sol. EXAMINATION

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

X → 1

$$n(E) = \frac{10!}{2! 2! 2!}$$

A → 2

$$P(E) = \frac{n(E)}{n(s)} = \frac{1}{11}$$

M → 1

O → 1

T → 1

N → 2

I → 2

- 2 If a cricket team consist of 15 players have 6 batsmen , 7 Ballers and 2 wicket keepers then the number

Sol. **Case-I :** Team consist 5 Batsman , 5 Bowlers

and 1 wicket keeper then number of ways.

$$= {}^6C_5 \times {}^7C_5 \times {}^2C_1 = 6 \times 21 \times 2 = 252$$

Case - II 4 Batsmen, 6 bowlers and 1 wicket keeper

$$= {}^6C_4 \times {}^7C_6 \times {}^2C_1 = 15 \times 7 \times 2 = 210$$

Case-III 4 Batsmen, 5 bowler and 2 wicket keepers

$$= {}^6C_4 \times {}^7C_5 \times {}^2C_2 = 15 \times 21 \times 1 = 315$$


Total $252 + 210 + 315 = 777$

 **JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS**

- 3 Tangent drawn at a point $P(2,2)$ to parabola $y^2 = 2x$ cuts x-axis at point Q and normal drawn at point $P(2,2)$ to parabola cut parabola again at point R then area of ΔPQR is

- (1) 25 (2) $\frac{25}{2}$ (3) $\frac{15}{2}$ (4) 50

Sol.



Equation of tangent at $P(2,2)$ is $T = 0$

$$2y = x + 2$$

$$y^2 = 4 So, Q(-2,0)$$

$$2at_1 = 1 \Rightarrow t_1 = 2$$

$$\therefore + 2 \sim 2 \sim$$

$$\therefore R\left(\frac{1}{2}(-3)^2, 2\left(\frac{1}{2}\right)(-3)\right) = \left(\frac{9}{2}, -3\right)$$

$$\text{Area of } \Delta PQR = \frac{1}{2} \begin{vmatrix} 2 & 2 & 1 \\ -2 & 0 & 1 \\ \frac{9}{2} & -3 & 1 \end{vmatrix}$$

$$= \frac{1}{2} [2(0+3) - 2(-2-9/2) + 1(6-0)] = \frac{1}{2} [6 + 4 + 9 + 6] = \frac{25}{2} \text{ sq. unit.}$$

4. Coefficient of x^{256} in the expansion of $(1-x)^{101}(x^2+x+1)^{100}$ is

- (1) ${}^{100}C_{86}$ (2) ${}^{100}C_{85}$ (3) ${}^{100}C_{84}$ (4) ${}^{100}C_{83}$

Ans. (2)

Sol. $\Rightarrow (1-x)^{101}(x^2+x+1)^{100}$

$$\Rightarrow (1-x)^{100}(x^2+x+1)^{100}(1-x)$$

$$\Rightarrow (1-x^3)^{100}(1-x)$$

$$\Rightarrow 100C_{85}x^{256}$$

so, the coefficient of x^{256} is $100C_{85}$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555



This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 2

5. The value of $\lim_{x \rightarrow 0} (2 - \cos x\sqrt{\cos 2x})^{\frac{x^2+2}{x}}$ is

Ans. 1

$$\begin{aligned} \text{Sol. } & \lim_{x \rightarrow 0} (2 - \cos x\sqrt{\cos 2x})^{\frac{x^2+2}{x}} \quad (1^\infty \text{ form}) \\ & = e^{\lim_{x \rightarrow 0} \frac{(1 - \cos x\sqrt{\cos 2x})}{x}(x^2 + 2)} \\ & = e^{\lim_{x \rightarrow 0} \frac{1 - \cos^2 x(\cos 2x)}{x}(x^2 + 2)} \\ & = e^{\lim_{x \rightarrow 0} \frac{1 - \cos^2 x(\cos 2x)}{x} \left(\frac{x^2 + 2}{1 + \cos x\sqrt{\cos 2x}} \right)} \\ & = e^{\lim_{x \rightarrow 0} \frac{1 - \cos^2(2\cos^2 x - 1)}{x} \left(\frac{x^2 + 2}{1 + \cos x\sqrt{\cos 2x}} \right)} \\ & = e^{\lim_{x \rightarrow 0} (1 - 2\cos^4 x + \cos^2 x) \cdot \frac{x^2 + 2}{x^2 + 2}} \end{aligned}$$

$$\begin{aligned} & = e^{\lim_{x \rightarrow 0} \frac{-(2\cos^4 x - \cos^2 x - 1)}{x} \cdot \frac{x^2 + 2}{1 + \cos x\sqrt{\cos 2x}}} \\ & = e^{\lim_{x \rightarrow 0} \frac{(2\cos^2 x + 1)(\cos^2 x - 1)}{x} \cdot \frac{x^2 + 2}{1 + \cos x\sqrt{\cos 2x}}} \\ & = e^{\lim_{x \rightarrow 0} \frac{+(2\cos^2 x + 1)\sin^2 x}{x} \cdot \frac{x^2 + 2}{1 + \cos x\sqrt{\cos 2x}}} \\ & = e^{\lim_{x \rightarrow 0} (2\cos^2 x + 1) \frac{\sin x}{x} \cdot \frac{x^2 + 2}{1 + \cos x\sqrt{\cos 2x}}} \\ & = e^0 = 1 \end{aligned}$$

6. If the focal chord $y = mx + c$ of parabola $y^2 = -64x$ is also the tangent to the circle $(x+10)^2 + y^2 = 4$ then absolute value of $4\sqrt{2}(m+c)$ is

Ans. 34

Now slope form of tangent to the circle

$(x+10)^2 + y^2 = 4$ is given by

$$y = m(x+10) \pm 2\sqrt{1+m^2}$$

$$\text{So, } c = 10m \pm 2\sqrt{1+m^2} \dots\dots\dots (ii)$$

By (i) and (ii)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555



This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 3

$$16m = 10m \pm 2\sqrt{1+m^2}$$

$$\Rightarrow 9m^2 = 1 + m^2 \Rightarrow m = \pm \frac{1}{2\sqrt{2}}$$

$$\Rightarrow c = 16m = \pm \frac{8}{\sqrt{2}}$$

$$\therefore 4\sqrt{2}(m+c) = \pm 34$$

7. The mean of 6 numbers is 6.5 and its variance is 10.25 if 4 numbers are 2, 4, 5 and 7, then find the other two :

(1) 10 11

(2) 11 12

(3) 9 12

(4) 9 11

Sol. Let two number x and y according to question

$$18 + x + y = 39$$

$$x + y = 21 \quad \dots\dots\dots(1)$$

$$10.25 = \frac{\sum x_i^2}{n} - (\bar{x})^2$$

$$10.25 = \frac{x^2 + y^2 + 4 + 16 + 25 + 49}{6} - (6.5)^2$$

$$10.25 = \frac{x^2 + y^2 + 94}{6} - (6.5)^2$$

$$\Rightarrow x^2 + y^2 = 221 \quad \dots\dots\dots(2)$$

solving (1) and (2)

So, $x = 10$ or $y = 11$

(1) point of local maxima


(2) point of local minima

(3) point of Inflection

(4) None of these

Ans. (1)

Sol. Roughly graph of $f(x)$ can be drawn as



Thus $x = \frac{3}{2}$ is a point of local maxima

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more, visit [RESCO](http://www.resonance.ac.in) at [Eduventures](http://www.resonance.ac.in) | www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

9. If the roots of the quadratic equation $x^2 + 3^{\frac{1}{4}}x + 3^{\frac{1}{2}} = 0$ are α and β then the value of $\alpha^{96}(\alpha^{12}-1) + \beta^{96}(\beta^{12}-1)$

(1) 50.3^{24}

(2) 51.3^{24}

(3) 52.3^{24}

(4) 104.3^{24}

Ans. (3)

$$\text{Sol. } x^2 + \sqrt{3}x + 3^{\frac{1}{2}} = 0$$

$$\begin{aligned}
&\Rightarrow x^8 + 3x^4 + 9 = 3x^8 \\
&\Rightarrow x^8 + 3x^4 + 9 = 0 \\
&\Rightarrow \alpha^8 = -9 - 3\alpha^4 \\
&\Rightarrow \alpha^{12} = -9\alpha^4 - 3\alpha^8 = -9\alpha^4 - 3(-9 - 3\alpha^4) = 27 \\
&\text{Similarly } \beta^{12} = 27 \\
&\Rightarrow \alpha^{96}(\alpha^{12} - 1) + \beta^{96}(\beta^{12} - 1) = (27)^8 \cdot 26 + (27)^8 \cdot 26 = 52 \cdot (27)^8 = 52 \cdot 3^{24}
\end{aligned}$$

10. In a $\triangle ABC$, If $AB = 5$, $\angle B = \cos^{-1}(3/5)$ and radius of circumcircle of triangle is 5 then the area of $\triangle ABC$ is

$$(1) 6+8\sqrt{3} \quad (2) 3+4\sqrt{3} \quad (3) 3+8\sqrt{3} \quad (4) 6+4\sqrt{3}$$

Ans. (1)

Resonance 3 4

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow \frac{b}{2R} = \frac{4}{5} \Rightarrow b = 8, c = 5$$

$$\cos B = \frac{a^2 + c^2 - b^2}{2ac} = \frac{3}{5} \Rightarrow \frac{a^2 + 25 - 64}{2a(5)} = \frac{3}{5}$$

$$a^2 - 39 = 6a \Rightarrow a^2 - 6a - 39 = 0$$

$$\Rightarrow a = \frac{6+8\sqrt{3}}{2} \Rightarrow a = 3+4\sqrt{3}$$

$$\Delta = \frac{abc}{4R} = \frac{(3+4\sqrt{3})(8)(5)}{4(5)} = 6+8\sqrt{3}$$

11. The number of integral terms is the expansion of $\left(\frac{1}{4^{\frac{1}{2}}} + \frac{1}{5^{\frac{1}{6}}}\right)^{120}$ is :

$$(1) 11 \quad (2) 21 \quad (3) 20 \quad (4) 30$$

Ans. (2)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 5

Resonance® Educating for better tomorrow | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS

Sol. General term of $\left(\frac{1}{2^{\frac{1}{2}}} + \frac{1}{5^{\frac{1}{6}}}\right)^{120}$ is

$$\text{given by } T_{r+1} = {}^{120}C_r \left[2^2 \right]^{120-r} \left[5^6 \right]^r$$

For integral term, r should be a multiple of 6

$$\text{i.e. } r \in \{0, 6, 12, 18, \dots, 120\}$$

$$\therefore 21 \text{ integral terms are there in the expansion } \left(\frac{1}{2^{\frac{1}{2}}} + \frac{1}{5^{\frac{1}{6}}}\right)^{120}$$

12. If the shortest distance between the lines $\vec{r}_1 = \hat{\alpha i} + 2\hat{j} + 2\hat{k} + \lambda(\hat{i} - 2\hat{j} + 2\hat{k}), \lambda \in \mathbb{R}, \alpha > 0$ and

$$\vec{r}_2 = -4\hat{i} - \hat{k} + \mu(3\hat{i} - 2\hat{j} - 2\hat{k}), \mu \in \mathbb{R}$$

$$(1) 2 \quad (2) 4 \quad (3) 6 \quad (4) \sqrt{6}$$

Ans. (3)

$$\text{Sol. Shortest distance} = \frac{|(\mathbf{a}_2 - \mathbf{a}_1) \cdot (\mathbf{b}_1 \times \mathbf{b}_2)|}{|\mathbf{b}_1 \times \mathbf{b}_2|}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow 9 = \frac{|((\alpha + 4)\hat{i} + 2\hat{j} + 3\hat{k}) \cdot (8\hat{i} + 8\hat{j} + 4\hat{k})|}{\sqrt{64 + 64 + 16}}$$

$$\Rightarrow \left| \frac{8(\alpha + 4) + 16 + 12}{12} \right| = 9$$

$$\therefore \alpha = 6$$

13. If $\vec{a}, \vec{b}, \vec{c}$ are mutually \perp unit vectors equally inclined to $\vec{a} + \vec{b} + \vec{c}$ at an angle θ , find $36\cos^2 2\theta$.

Ans. 4

$$\text{Sol. } |\vec{a} + \vec{b} + \vec{c}|^2 = (\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c}) = |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 = 3$$

$$\Rightarrow |\vec{a} + \vec{b} + \vec{c}| = \sqrt{3}$$

$$\text{Now } \vec{a}(\vec{a} + \vec{b} + \vec{c}) = |\vec{a}| |\vec{a} + \vec{b} + \vec{c}| \cos\theta$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow \cos 2\theta = -\frac{1}{3} \Rightarrow \cos^2 2\theta = \frac{1}{9} \Rightarrow 36\cos^2 2\theta = 4$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 7340010333

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 6

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

14. If z and ω are complex number such that $|z\omega| = 1$, $\arg(z) - \arg(\omega) = \frac{3\pi}{2}$. Find the $\arg\left(\frac{1-2\bar{z}\omega}{1+3\bar{z}\omega}\right)$.

(1) $\frac{\pi}{4}$

(2) $-\frac{\pi}{4}$

(3) $\frac{3\pi}{4}$

(4) $-\frac{3\pi}{4}$

Ans. (4)

$$\text{Sol. Let } z = re^{i\theta} \text{ & } \omega = \frac{1}{r} e^{i\left(\theta - \frac{3\pi}{2}\right)}$$

$$\text{then } \frac{1-2\bar{z}\omega}{1+3\bar{z}\omega} = \frac{1-2re^{-i\theta} \cdot \frac{1}{r} e^{i\left(\theta - \frac{3\pi}{2}\right)}}{1+3re^{-i\theta} \cdot \frac{1}{r} e^{i\left(\theta - \frac{3\pi}{2}\right)}}$$

$$1-2e^{-\frac{3\pi}{2}} \quad 1-2i$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$= -\frac{1}{2} - \frac{1}{2}i$$

$$\text{The } \arg\left(-\frac{1}{2} - \frac{1}{2}i\right) = -\frac{3\pi}{4}$$

15. If $f(x) = \begin{cases} \sin x - e^x & ; x \leq 0 \\ a + [-x] & ; 0 < x < 1 \\ 2x - b & ; x \geq 1 \end{cases}$ is continuous and differentiable function then find the value of $a + b$.

(where $[x]$ is GIF)

Ans. (03.00)

Sol. Since $f(x)$ is continuous at $x = 0$

$$\text{So } \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$$

$$-1 = a - 1 = -1 \Rightarrow a = 0$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = f(1)$$

$$a - 1 = 2 - b = 2 - b$$

$$\Rightarrow a = 0, \text{ so } 0 - 1 = 2 - b$$

$$\Rightarrow -3 = -b$$

$\Rightarrow b = 3$

So the value of $a + b = 3$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/resonancewatch blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 7

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



| JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS

16. If $A = [a_{ij}]_{3 \times 3}$ where $a_{ij} = \begin{cases} 1 & i=j \\ -x & |i-j|=1 \\ 2x+1 & \text{otherwise} \end{cases}$ and $f(x) = \det(A)$, then the sum of local maximum and

local minimum value of $f(x)$ is:

$$(1) \frac{20}{27}$$

$$(2) \frac{-20}{27}$$

$$(3) \frac{88}{27}$$

$$(4) \frac{-88}{27}$$

Ans. (4)

Sol. $|\mathbf{A}| = \begin{vmatrix} 1 & -x & 2x+1 \\ -x & 1 & -x \\ 2x+1 & -x & 1 \end{vmatrix} = 1+x^2(2x+1)+x^2(2x+1)-(2x+1)^2-x^2-x^2$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\rightarrow f(x) = 4x^3 - 4x^2 - 4x$$

$$\Rightarrow f'(x) = 12x^2 - 8x - 4$$

$$= 4(3x^2 - 2x - 1)$$

$$= 4(x-1)(3x+1)$$

$$\begin{array}{c|ccc} + & & - & + \\ \hline -1 & & & 1 \\ \hline 3 & & & \end{array}$$

$\Rightarrow f(x)$ is maximum at $x = -\frac{1}{3}$ and minimum at $x = 1$

$$\therefore \text{maximum value} = \frac{20}{27} \text{ and minimum value} = -4$$

$$\therefore \text{sum} = \frac{20}{27} - 4 = -\frac{88}{27}$$

17. The coefficient of $a^3 b^4 c^5$ in $(ab + bc + ac)^6$ is:

$$(1) 60$$

$$(2) 45$$

$$(3) 40$$

$$(4) 90$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol. $(ab + bc + ac)^6 = \sum_{p+q+r=6} \frac{6!}{p!q!r!} (ab)^p (bc)^q (ca)^r$

$$= \sum_{p+q+r=6} \frac{6!}{p!q!r!} a^{p+r} b^{p+q} c^{q+r}$$

For $a^3 b^4 c^5$, we need

$$p + r = 3$$

$$p + q = 4$$

$$q + r = 5$$

Solving we get, $p = 1, q = 3, r = 2$

$$\therefore \text{coefficient of } a^3 b^4 c^5 \text{ in } (ab + bc + ac)^6 \text{ is } \frac{6!}{1!2!3!} = 60$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/resonancewatch blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 8

18. If an invertible function $f(x)$ is defined as $f(x) = 3x - 2$, $g(x)$ is also an invertible function such that

$f^{-1}(g^{-1}(x)) = x - 2$ then $g(x)$ is

- (1) $\frac{x-8}{3}$ (2) $\frac{x+8}{3}$ (3) $\frac{x-3}{8}$ (4) $\frac{x+3}{8}$

Ans. (2)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$3(x-2) - 2 = g^{-1}(x)$$

$$3x - 8 = g^{-1}(x)$$

$$g^{-1}(x) = 3x - 8$$

or $x = 3g(x) - 8$

$$g(x) = \frac{x+8}{3}$$

19. $\int_{-1}^1 \ln(\sqrt{1-x} + \sqrt{1+x}) dx = ?$

- (1) $\pi + \ln 2$ (2) $2/\ln 2$ (3) $\frac{\pi}{2} - 1 + \ln 2$ (4) $\ln 2 - \frac{\pi}{2} - 1$

Ans. (4)

Sol. $f(x) = \ln(\sqrt{1-x} + \sqrt{1+x})$ $x \in [-1, 1]$ is an even function

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\text{Put } x = \cos 2\theta \Rightarrow dx = -2\sin 2\theta d\theta$$

$$\therefore \cos 2\theta = 2\cos^2 \theta - 1 = 1 - 2\sin^2 \theta$$

$$\Rightarrow I = -4 \int_{\frac{\pi}{4}}^0 [\ln(\sin \theta + \cos \theta)\sqrt{2}] \sin 2\theta d\theta$$

$$= 4 \int_0^{\frac{\pi}{4}} [\ln(\sin \theta + \cos \theta)\sqrt{2}] \sin 2\theta d\theta$$

$$= 4 \int_0^{\frac{\pi}{4}} [\ln(\sin \theta + \cos \theta) \sin 2\theta d\theta + 4 \int_0^{\frac{\pi}{4}} \ln \sqrt{2} \sin 2\theta d\theta]$$

$$= 4 \left[\left| -\ln(\sin \theta + \cos \theta) \frac{\cos 2\theta}{2} \right|_0^{\frac{\pi}{4}} + \left| \frac{\cos \theta - \sin \theta}{\sin \theta + \cos \theta} \cdot \frac{\cos 2\theta}{2} \right|_0^{\frac{\pi}{4}} \right] + 4 \ln \sqrt{2} \left(-\frac{\cos 2\theta}{2} \right)_0^{\frac{\pi}{4}}$$

$$= \left[1 \right] + \left[\ln \sqrt{2} \right]$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/ResonanceWatch blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 9

$$= -4 \left[0 - \frac{1}{4} \int_0^{\frac{\pi}{4}} (1 - \sin 2\theta) d\theta \right] + 2 \ln \sqrt{2}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$= -4 \left[\theta + \frac{\cos 2\theta}{2} \right]_0^{\frac{\pi}{4}} + 2 \ln \sqrt{2}$$

$$= -2 \left[\frac{\pi}{4} - \frac{1}{2} \right] + 2 \ln \sqrt{2}$$

$$= -\frac{\pi}{2} - 1 + 2 \ln \sqrt{2}$$

20. The probability of selecting integers $a \in [-5, 30]$, such that $x^2 + 2(a+4)x - 5a + 64 > 0$ for all $x \in \mathbb{R}$ is :

(1) $\frac{7}{9}$

(2) $\frac{4}{9}$

(3) $\frac{5}{9}$

(4) $\frac{1}{3}$

Ans. (3)

Sol. $x^2 + 2(a+4)x - (5a - 64) > 0$

$D < 0$

$\therefore 4(a+4)^2 + 4(5a - 64) < 0$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$\Rightarrow a^2 + 13a - 48 < 0$

$a = \frac{-13 \pm \sqrt{169 + 192}}{2}$

So, $a \in [-16, 3]$

Total integers are 20 \therefore Probability = $\frac{20}{36} = \frac{5}{9}$

21. If $\int_0^a e^{-[x]} dx = 10e - 9$, then the value of 'a' is (where $[x]$ is GIF)

(1) $9 + \ln 2$

(2) $10 + \ln 2$

(3) 10

(4) 9

Ans. (2)

Sol. Let $a = 10 + K$, $0 \leq K < 1$

$\int_0^a e^{-[x]} dx = 10e - 9$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$e^K - 1 = 1$

$K = \ln 2$

so, $a = 10 + \ln 2$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555

7340010333



facebook.com/ResonanceEdu



twitter.com/ResonanceEdu



www.youtube.com/Resonance



blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 10

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

22. If $\vec{A} \cdot \vec{B} = |\vec{A} \times \vec{B}|$, then $|\vec{A} - \vec{B}|$ is

(1) $\sqrt{A^2 + B^2 + \sqrt{2}AB}$ (2) $\sqrt{A^2 + B^2 - \sqrt{2}AB}$ (3) $\sqrt{A^2 + B^2 + \sqrt{2}AB}$ (4) $\sqrt{A^2 + B^2 - \sqrt{2}AB}$

Ans. (4)

Sol. $\vec{A} \cdot \vec{B} = |\vec{A} \times \vec{B}| \Rightarrow \cos\theta = \sin\theta \Rightarrow \tan\theta = 1$

$\theta = \frac{\pi}{4}$

$|\vec{A} - \vec{B}|^2 = A^2 + B^2 - 2\vec{A} \cdot \vec{B}$

$= A^2 + B^2 - 2AB\cos\left(\frac{\pi}{4}\right)$

$= A^2 + B^2 - \sqrt{2}AB$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/resonancewatch blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 11