

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac in | CIN - U80302RJ/2007PLC024029

2021

COMPUTER BASED TEST (CBT) Memory Based Questions & Solutions

Date: 20 July, 2021 (SHIFT-2) | TIME : (3.00 p.m. to 6.00 p.m)

Duration: 3 Hours | Max. Marks: 300

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

Resonance® | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

PART: PHYSICS

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Ans. (2

Sol. $K.E. \Rightarrow K = \frac{P^2}{2m}$

P∝√K

P₂ K₂ P₂ 4K

$$\overline{P_1} = \sqrt{K_1} \rightarrow \overline{P_1} = \sqrt{K}$$

$$\Rightarrow \frac{P_2}{P_4} = 2$$

$$\Rightarrow \frac{P_2 - P_1}{P_1} \% = \left(\frac{P_2}{P_1} - 1\right) \times 100 = (2 - 1) \times 100 = 100$$

$$\Rightarrow \frac{\Delta P}{P_1} \% = 100\%$$

- A RLC circuit is in its resonance condition. Its circuit components have value

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

- (2) 10kW
- (3) 12kW
- (4) 12.5kW

As circuit is in resonance. Thus

$$\therefore$$
 Z = R so i_{ms} = V/Z = V/R

$$P = \frac{V^2}{R} = \frac{250 \times 250}{5} = 12500 \text{ J/s} = 12.5 \text{ kW}$$

- A wheel rotating with an angular speed of 600 rpm is given an constant angular acceleration of 1800 rpm² for 10 sec. Number of revolutions revolved by wheel is:
 - (1) 125
- (2) 100
- (4)50

Ans. (1)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555 | 7340010333 | entertok contribution of the contributi

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#1

Resonance* | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

 $\omega_0 = 600 \text{ rpm}$

$$\alpha = 1800 \text{ rpm}^2$$

$$\theta = \omega_0 t + \frac{1}{2} \alpha t^2$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

= 100 + 25 = 125 revolution.

- $|\vec{P}| = |\vec{Q}|, |\vec{P} + \vec{Q}| = |\vec{P} \vec{Q}|$. Find angle between $\vec{P} & \vec{Q}$
 - (1) 45°
- (2) 90°
- (3) 135°
- (4) 150°

Ans. (2)

Sol.
$$|\vec{P} + \vec{Q}| = |\vec{P} - \vec{Q}|$$

$$|\vec{P}|^2 + |\vec{Q}|^2 + 2|\vec{P}||\vec{Q}|\cos\theta = |\vec{P}|^2 + |\vec{Q}|^2 - 2|\vec{P}||\vec{Q}|\cos\theta$$

$$|\vec{P}||\vec{Q}|\cos\theta = 0^{\circ}$$

Thus,
$$\theta = 90^{\circ}$$

- A body is moved from rest along straight line by a machine delivering a constant power. Time taken by body to travel a distance "S" is proportional to
 - (1) S1/3
- (2) S2/3
- (3) S1/2
- (4) S1/4

Ans.

Energy supply = Pt

$$t^{3/2} = \frac{3S}{2C}$$

$$t = S^{2/3} \left(\frac{3}{2C}\right)^{2/3}$$

T ∞ S^{2/3}

Decompose Eduvanturas I td

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac in | CIN - U80302RJ2007PLC024029

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222 To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555 🕲 7340010333 🌠 sockook ours ResonanceEdu 💟 twitter.com/ResonanceEdu 🛗 www.youtube.com/tresonation 🛅 biog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

Resonance® | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

- A uniform rod of young's modulus Y is stretched by two tension T₁ and T₂ such that rods get expanded to length L₁ and L₂ respectively. Find initial length of rod ?
 - (1) $\frac{L_1T_1-L_2T_2}{L_1T_1}$
- $(2) \ \frac{L_2T_1 L_1T_2}{T_2 T_1}$
- (3) $\frac{L_1 T_2 L_2 T_1}{T_2 T_1}$ (4) $\frac{L_1}{T_1} \times \frac{T_2}{L_2}$

Ans. (3)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

As
$$\frac{1}{A} = Y \frac{\Delta \ell}{\ell}$$

So, $\frac{T_1}{A} = \frac{Y(L_1 - L_0)}{L_0}$
 $\frac{T_2}{A} = \frac{Y(L_2 - L_0)}{L_0}$

$$\frac{T_1}{T_2} = \frac{L_1 - L_0}{L_2 - L_0} \quad ; \quad T_1 L_2 - T_1 L_0 = T_2 L_1 - T_2 L_0 \quad ; \quad L_0 = \frac{L_1 T_2 - L_2 T_1}{T_2 - T_1}$$

- Time (T), velocity (C) and angular momentum (h) are choosen as fundamental quantities instead of mass, length and time. In term of these, dimension of mass would be
 - (1) $[M] = [T^{-1}C^{-2}h]$

(2) $[M] = [T^{-1}C^2h]$

(3) $[M] = [T^{-1}C^{-2}h^{-1}]$

(4) $[M] = [T^{-1}C^{-2}h]$

Ans. (1)

Sol. M or TxCyhz

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

On comparing powers
$$z = 1$$
 ...(1)

$$x - y - z = 0$$
 ...(2)

$$y + 2z = 0$$
 ...(3)
 $y + 2 \times 1 = 0$

$$x - (-2)-1 = 0$$

T 10 2 L

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-7777777 2777700 | FAX No.: +91-022-39167722

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac_in | CIN - U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

Resonance | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

- 8. Find relation between γ (adiabatic constant) and degree of freedom (t)
 - (1) $f = \frac{2}{x-1}$
- (2) $f = \frac{\gamma}{\gamma 1}$
- (3) $f = \frac{y 1}{2}$
- (4) $f = \frac{\gamma 1}{\gamma}$

PAGE#3

Ans. (1

Sol. $C_V = \frac{fR}{2}$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302R.J2007PLC024029

$$\Rightarrow \qquad \gamma = \frac{6\beta}{C_V} = 1 + \frac{2}{f}$$

$$\gamma = \frac{1 + \frac{2}{f}}{f}$$

$$f = \frac{2}{\gamma - 1}$$

a (f .)

- Two identical drops of Hg coalesce to form a bigger drop. Find ratio of surface energy of bigger drop to smaller drop.
 - (1) 23/2
- (2) 32/5
- (3) 22/3
- (4) 52/3

Ans. (3


Sol.
$$2 \times \frac{4}{3} \pi r^3 = \frac{4}{3} \pi R^3$$

$$\frac{R}{2} = 2^{1/3}$$

...(1)

Now
$$\frac{U_{\text{bigger}}}{U_{\text{smaller}}} = \frac{S \times 4\pi R^2}{S \times 4\pi r^2} = \left(\frac{R}{r}\right)^2 = 2^{2/3}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744-2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



Ans. (3)

Sol. PV = nRT

 \Rightarrow PV = CT

Therefore, PV v/s T graph is straight line.

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 27777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555 | 7340010333 | socioux oun-fleorencetta | www.resonance.ac.in | www.youtube.com/fleorencetta | www.resonance.ac.in | www.reson

Resonance | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

- For a body in pure rolling, its rotational kinetic energy is 1/2 times of its translation kinetic energy. They body should be ?
 - (1) solid cylinder
- (2) Ring
- (3) solid sphere
- (4) Hollow sphere

Ans. (1)

Sol. Given

$$\frac{1}{2}$$
I $\omega^2 = \frac{1}{2} \times \frac{1}{2}$ my ²

as v = R @ (pure rolling

1. 0 1 _00

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302R.J2007PLC024029

$$I = \frac{1}{2} mR^2$$

Thus, solid cylinder.

- Magnetic susceptibility of material is 499 & $\mu_0 = 4\pi \times 10^{-7}$. SI unit then find μ_r
 - (1) 500
- (2)400
- (3) 300
- (4)200

Ans. (1)

Sol. $\mu_r = 1 + \chi$

= 1 + 499 = 500

- 13. A plane electromagnetic wave travels in free space. Electric field is $\ddot{E}=E_0i$ and magnetic field is represented by $\vec{B} = B_0 k$. What is the unit vector along the direction of propagation of electromagnetic wave?
 - (1) j
- (2) k
- (3) j
- (4) k

Ans. (3)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Unit vector in direction
$$\vec{E} \times \vec{B} \Rightarrow \frac{\vec{E} \times \vec{B}}{|\vec{E} \times \vec{B}|}$$

$$\Rightarrow \frac{\hat{E_0 i \times B_0 k}}{\hat{E_0 B_0 s in 90}}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555
7340010333
to extend confliction confliction contact with the contact with the confliction contact with the co

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#5

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

RESONANCE | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

- Two satellites of mass MA and MB are revolving around a planet of mass M in radius RA and RB respectively. Then ?
 - (1) TA > TB if
- RA > RB
- (2) TA > TB

- (3) TA = TB
- if MA > MB
- RA < RB

Ans. (1)

Sol. T ∞ r^{3/2}

$$\frac{T_A}{T_B} \propto \left(\frac{R_A}{R_B}\right)^{3/2}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$N_0 \xrightarrow{t_{1/2}} \frac{N_0}{2} \xrightarrow{t_{1/2}} \frac{N}{4} \xrightarrow{t_{1/2}} \frac{N}{8} \xrightarrow{t_{1/2}} \frac{N}{16}$$

$$4 \times t_{1/2} = 80 \text{ days}$$

16. A satellite is revolving around a planet in an orbit of radius R. Suddenly radius of orbit becomes 1.02 R then what will be percentage change in its time period of revolution?

Ans.

Sol. As
$$T \propto R^{3/2}$$

$$T_1 = kR^{3/2}$$

$$\frac{\Delta T}{T} = \frac{3}{2} \times \frac{\Delta R}{R} = 3\%$$

17. A person walks up a stationary escalator in the time t₁. If he remains stationary on the escalator, then it can take him up in time t2. Determine the time it would take to walk up on the moving escalator?

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555 | 7340010333 | evertex confliction models | www.youtube.com/resoverced.

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

Suppose length of escalator = L

Speed of man wrt escalator = $\frac{L}{t}$.

Speed of escalator = L

Speed of man wrt ground when escalator is moving = $\frac{L}{t_1} + \frac{L}{t_2}$

Time taken by the man to walk on the moving escalator $= \frac{L}{\frac{L}{t_1} + \frac{L}{t_2}} = \frac{t_1 t_2}{t_1 + t_2}$

For given graph between decay rate & time. Find half life (where R = decay rate)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



$$(1) \frac{10}{3} \ln 2$$

$$(2) \frac{20}{3} \ln 2$$

$$(3) \frac{3}{20} \ln 2$$

(3)
$$\frac{3}{20} \ln 2$$
 (4) $\frac{20}{3} \ln 2$

Sol. $R = R_0e^{-\lambda t}$

$$lnR = lnR_0 - \lambda t$$

slope =
$$-\lambda = \frac{-6}{40}$$

$$\lambda = \frac{3}{20}$$

$$t_{1/2} = \frac{\ln 2}{\lambda} = \frac{\ln 2}{3} \times 20 = \frac{20}{3} \ln 2$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

time period of oscillation?

(1)
$$2\pi \sqrt{\frac{x_2^2 + x_1^2}{v_1^2 - v_2^2}}$$

(2)
$$2\pi \sqrt{\frac{x_2^2 - x_1^2}{v_1^2 + v_2^2}}$$

$$(1) \ \ \, \frac{2\pi}{v_1^2-v_2^2} \sqrt{\frac{x_2^2+x_1^2}{v_1^2-v_2^2}} \qquad \qquad (2) \ \ \, \frac{2\pi}{v_1^2+v_2^2} \sqrt{\frac{x_2^2-x_1^2}{v_1^2+v_2^2}} \qquad \qquad (3) \ \ \, 2\pi \sqrt{\frac{x_2^2-x_1^2}{v_1^2-v_2^2}} \ \ \, (4) \ \ \, 2\pi \sqrt{\frac{x_2^2+x_1^2}{v_1^2+v_2^2}}$$

(3) Ans.

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
besteek own fleeconceds: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
besteek own fleeconceds: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$V = \omega \sqrt{A^2 - x^2}$$

$$v_1 = \omega \sqrt{A^2 - x_1^2}$$

$$v_2 = \omega \sqrt{A^2 - x_2^2}$$

$$\left(\frac{\mathbf{v}_1}{\omega}\right)^2 - \left(\frac{\mathbf{v}_2}{\omega}\right)^2 = \mathbf{x}_2^2 - \mathbf{x}_1^2$$

$$\omega^2 = \frac{v_1^2 - v_2^2}{x_2^2 - x_1^2}$$

$$\omega = \sqrt{\frac{v_1^2 - v_2^2}{x_2^2 - x_1^2}}$$

$$T = 2\pi \sqrt{\frac{x_2^2 - x_1^2}{v_1^2 - v_2^2}}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Ans. (3)

$$eV = \frac{hc}{\lambda} - W$$

For first case

$$e(3V_0) = \frac{hc}{\lambda_0} - W \qquad \dots (i)$$

For second case

$$eV_0 = \frac{hc}{2\lambda_0} - W$$
 ...(ii)

From equation (i) and (ii)

$$W = \frac{hc}{4\lambda_0}$$



Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555 S 7340010333 cottok om/lesonancela: void more in the same particles of the same particle

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#8

RESONANCE | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

(1)
$$\tan^{-1} \left(\frac{1}{\sqrt{6}} \right)$$

(2)
$$\tan^{-1} \left(\frac{1}{\sqrt{2}} \right)$$

(3)
$$\tan^{-1} \left(\frac{1}{\sqrt{4}} \right)$$

(4)
$$\tan^{-1} \left(\frac{1}{\sqrt{3}} \right)$$

Ans. (1)

Let vertical and horizontal component of earth's magnetic field at meridian will be V and H. Sol.

Angle of dip,
$$\tan\theta = \frac{V}{H}$$
 ...(i)

at angle of 45° from magnetic meridian, angle of dip = 30°

$$\tan 30^{\circ} = \frac{V}{H\cos 45^{\circ}}$$
 \Rightarrow $\frac{1}{\sqrt{3}} = \frac{V}{H\cos 45^{\circ}}$

$$\frac{V}{H} = \frac{1}{\sqrt{6}}$$

$$\tan\theta = \frac{V}{H} \Rightarrow \frac{1}{\sqrt{6}}$$

$$\theta = \tan^{-1} \left(\frac{1}{\sqrt{6}} \right)$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

A sodium lamp in space was emitting waves of wavelength 2880Å. When observed from a planet, its wavelength was recorded 2886 Å. Find the speed of planet ?

$$(2) 6.25 \times 10^{5} \text{ m/s}$$

Ans.

Sol.
$$\frac{V_{rel}}{C} = \frac{\Delta \lambda}{\lambda}$$

$$V_{rel} = \frac{6}{2880} \times 3 \times 10^8$$

$$= 6.25 \times 10^5 \text{ m/s}$$

- An electron having debroglie wavelength is falls on an X-ray tube. The cut off wave length of emitted X-Ray is
 - (1) 2m cλ²
- (2) 2h
- $(3) \frac{h}{mc}$
- $(4) \ \frac{2}{3} \frac{\text{mc}\lambda^2}{h}$

Ans. (1)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
Tooltok contifesconcedu
www.resonancedu
www

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

De-broglie wavelength

1 _ h

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow P = \frac{\pi}{\lambda_B}$$

∴ Kinetic energy of electron \Rightarrow E = $\frac{P^2}{2m_e} = \frac{h^2}{2m_e \lambda_B^2}$

For cut-off wavelength of emitted X-Ray

$$E = \frac{hc}{\lambda}$$

$$\Rightarrow \frac{h^2}{2m_e\lambda_B^2} = \frac{hc}{\lambda}$$

$$\Rightarrow \lambda = \frac{2m_e c \lambda_B^2}{h} = \frac{2mc\lambda^2}{h} \text{ where } \lambda_B = \lambda \& m_e = m.$$

A gas is undergoing change in state by an isothermal process AB as follows. Work done by gas in process 24.



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



(1) 100 In2 Joule

(2) - 100 ln2 Joule (3) 200 ln2 Joule

(4) - 200 In2 Joule

Ans.

Wisothermal = $P_1V_1 \ln \frac{V_2}{V_1}$ Sol.

 $V_1 = 100m^3$

 $V_2 = 200 m^3$

 $P_1 = 2 N/m^2$

 $W = 2 \times 100 \ln \frac{200}{100}$

= 200 ln2 Joule

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555 | 7340010333 | socioik.com/Resonance.ac.in | www.youtube.com/Resonance.ac.in | www.youtube.com/Resonance.ac.in | cin: uses the contact of t

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 10

RESONANCE | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

A block is projected upto a rough plane of inclination 30°. If time of ascending is half the time for descending and the coefficient of friction is $\mu = \frac{3}{5\sqrt{n}}$. Then n =

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$S = \frac{1}{2} a_A t_A^2$$
(1)

$$S = \frac{1}{2} a_D t_D^2$$
(2)

From Equation (1) & (2)

$$\frac{t_A^2}{t_D^2} = \frac{a_D}{a_A}$$

$$\Rightarrow \frac{t_A^2}{t_A^2} = \frac{g \sin \theta - \mu g \cos \theta}{1 - \mu g \cos \theta}$$

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow \frac{t_A}{t_D} = \sqrt{\frac{g\sin\theta + \mu g\cos\theta}{g\sin\theta + \mu g\cos\theta}}$$

$$\Rightarrow \qquad \frac{1}{2} = \sqrt{\frac{1 - \sqrt{3}\mu}{1 + \sqrt{3}\mu}}$$

$$\Rightarrow 1 + \sqrt{3}\mu = 4 - 4\sqrt{3}\mu$$

$$\Rightarrow$$
 $5\sqrt{3}\mu = 3$

$$\Rightarrow \qquad \mu = \frac{3}{5\sqrt{3}}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302R12007PLC024029

Toll Free: 1800 258 5555
7340010333
Tooltexicon/Resonanceds


www.resonanceds

www.resonanceds

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

Resonance | Jee Main-2021 | Date : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

26. I - V characteristic curve of a diode in forward bias is given in fig. find out dynamic resistance -



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- (1) 212.3 Ω
- (2) 205.3Ω
- (3) 245.3Ω (4) 233.3Ω

Ans.

Sol. Dynamic resistance =
$$\frac{\Delta V}{\Delta I}$$

$$=\frac{0.7}{3\text{mA}}=233.3\Omega$$

- An electron is accelerated through a voltage of 40 kV. What will be its wavelength?
 - (1) 0.061Å
- (2) 0.011Å
- (3) 0.021Å
- (4) 0.161Â


Ans. (1)

Sol.
$$\lambda_B = \frac{h}{P}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$= \frac{12.27}{\sqrt{40 \times 10^3}} \dot{A} = 0.061 \dot{A}$$

28. Find value of R_P in given ckt ? ($V_Z = 8V$)



Resonance Eduventures Ltd.

(3) 3Ω

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302R.J2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal PAGE # 12

(4) 5Ω

RESONANCE | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

Ans. (2

Sol. Applying KVL

$$20 - 8 - 2RP = 0$$


$$R_P = 6\Omega$$

29. Two stars of masses m₁ and m₂ are in mutual interaction and revolving in orbits of radii r₁ and r₂ respectively. Time period of revolution for this system will be ?

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Ans. (2)

Sol.



Let angular velocity will be ω

For mass mi

$$Gm_1m_2 - m_1 r_1 \omega^2 - m_1 \times \frac{m_2(r_1 + r_2)}{\omega^2} \omega^2$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\omega = \frac{\sqrt{G(m_1 + m_2)}}{(r_1 + r_2)^{3/2}}$$

$$T = \frac{2\pi}{\omega}$$

$$= 2\pi \sqrt{\frac{(r_1 + r_2)^3}{G(m_1 + m_2)}}$$

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029










© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



COMPUTER BASED TEST (CBT) Memory Based Questions & Solutions

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



5 AIRs in TOP-50 in JEE (Adv.) 2020 from Classroom

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302R.J2007PLC024029

ADMISSION OPEN for Session 2021-22

ONLINE + OFFLINE PROGRAMS

CLASS 11, 12 & 12+

Target: JEE (Main+Adv.) | JEE (Main) | NEET

Scholarship Upto 90%*

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Resonance* | Jee Main-2021 | Date : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

PART : MATHEMATICS

1 If element of matrix A is defined as A = $[a_{ij}]_{3\times3}$ where A = $\begin{cases} (-1)^{j+i} & i < j \\ 2 & i = j \\ (-1)^{j+j} & i > j \end{cases}$, then the value of $|3Adj(2A^{-1})|$

(1) 72 (2) 36 (3) 108 (4) 48

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$A = \begin{bmatrix} -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$$

So, $|A| = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$
 $= 2(4-1) + 1(-2+1) + 1(1-2)$
 $= 2(3) + 1(-1) + 1(-1)$
 $= 4$
 $|3Ac|(2A^{-1})| = 3^3 |Ac|(2A^{-1})| = 3^3 \times |2A^{-1}|^2$
 $= 3^3 \times 2^6 \times |A^{-1}|^2 = 3^3 \times 2^6 \times \frac{1}{|A|^2} = 108$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$|\overrightarrow{AB}| = 7$$
, $|\overrightarrow{BC}| = 5$, $|\overrightarrow{CA}| = 3$,



Projection of BC on CA is = |BC| cos∠BCA

$$5\left(\frac{3^2+5^2-7^2}{2.3.5}\right) = 5\left|\frac{-15}{30}\right| = \frac{5}{2}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302R.J2007PLC024029

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

RESONANCE* | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

- The value of $tan(2tan^{-1}(3/5) + sin^{-1}(5/13))$ is:
- (2) $\frac{110}{21}$ (3) $\frac{55}{21}$

Δne

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$\tan^{-1} \frac{5}{1 - \frac{9}{25}} + \tan^{-1} \frac{5}{12}$$

$$\tan\left(\tan^{-1}\left(\frac{15}{8}\right) + \tan^{-1}\left(\frac{5}{12}\right)\right) = \frac{\frac{15}{8} + \frac{5}{12}}{1 - \frac{15}{8} \cdot \frac{5}{12}} = \frac{220}{21}$$

(2) 1

Mean of 6 observations is 10 and their variance is $\frac{20}{3}$. If observations are 15, 11, 10, 7, a, b then

|a - b| is equal to:

(3) 3

(1)2Ans

Mean = 10

$$\frac{7+10+11+15+a+b}{6}=10$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$\frac{49+100+121+225+a^2+b^2}{6}-100=\frac{2}{3}$$

$$a^2 + b^2 = 145$$

 $(a + b)^2 = 289$

ab = 72

$$(a - b)^2 = (a + b)^2 - 4ab$$

$$(a - b)^2 = 289 - 288 = 1$$

a -b | = 1

- If f(x) = x + 1, then find $\lim_{n \to \infty} \frac{1}{n} \left[f(0) + f\left(\frac{5}{n}\right) + f\left(\frac{10}{n}\right) + \dots + f\left(\frac{5(n-1)}{n}\right) \right]$

Ans.

Sol. =
$$\lim_{r \to \infty} \frac{1}{r} \int_{r}^{r-1} f(\frac{5r}{r}) = \int_{r}^{1} f(\frac{5r}{r}) dr = \int_{r}^{1} f(\frac{5r}{r})$$

$$=\left(\frac{5x^2}{2} + x\right)_0^1 = \frac{5}{2} + 1 = \frac{7}{2}$$
 Ans

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555
7840010333
Toolkok contresonance ac.in | www.resonance ac.in | www.youtube.com/resonance ac.in | CIN: U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

M Posnoance*

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777; 2777700 | contact@resonance ac in | CIN - U80302RJ2007PLC024029

Ans. (3)

Sol. 2 log₉x +3 log₉x +4 log₉x21 terms

=
$$(2 + 3 + 4 + 5 + \dots + 22) \log_9 x = \frac{21}{2} (2 + 22) \log_9 x$$

= 21 x 12 logsx

= 252 log9x

Given sum = 252 ⇒ logex = 1

$$\Rightarrow$$
 x = 9

 $([x] - [\sin x]) dx = ?$ (Where [-] represents G.I.F.)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$I = \int_{-\pi}^{\frac{\pi}{2}} ([x] - [\sin x]) dx$$

using property
$$\int_{-a}^{a} f(x)dx = \int_{0}^{a} f(x)dx + \int_{0}^{a} f(-x)dx$$

$$I = \int_{0}^{\frac{\pi}{2}} ([x] + [-x]) dx - \int_{0}^{\frac{\pi}{2}} ([\sin x] + [-\sin x]) dx = 0$$

8. If
$$\lim_{x\to 0} \frac{\alpha x e^x - \beta \ell n(1+x) + \gamma x^2 e^{-x}}{x^3} = 10$$
, then the value of $\alpha + \beta + \gamma$ is :

Ans.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow \alpha - \beta = 0, \Rightarrow \alpha = \beta$$

$$\Rightarrow \alpha + \frac{\beta}{2} + \gamma = 0 \Rightarrow \gamma = -\frac{3\beta}{2}$$

$$\Rightarrow \frac{\alpha}{2} - \frac{\beta}{3} - \gamma = 10$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555
7340010333
acceptable better conflection of the contact of the This solution was download from Resonance JEE (MAIN) 2021 Solution portal

RESONANCE* | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

$$\Rightarrow \frac{\beta}{2} - \frac{\beta}{3} + \frac{3\beta}{2} = 10 \Rightarrow \frac{3\beta - 2\beta + 9\beta}{6} = 10$$

:
$$\beta = 6$$
, $\alpha = 6$, $\gamma = -9$

So, the value of
$$\alpha + \beta + \gamma = 3$$

The value of x satisfying the equation $log_{(x+1)}(2x^2 + 7x + 5) + log_{(2x+5)}(x+1)^2 = 4$ is:

$$(3) -4$$

Ans. (2)

 $\log_{(x+1)}((2x+5)(x+1)) + \log_{(2x+5)}(x+1)^2 = 4$ Sol.

$$1 + \log_{(x+1)} (2x+5) + 2\log_{(2x+5)} (x+1) = 4$$

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$t^2 + t + 2 = 4t \Rightarrow t^2 - 3t + 2 = 0$$

For
$$t = 2$$

 $2x+5 = (x+1)^2$

$$\Rightarrow$$
 x = -4 (rejected)

$$x = 2$$
, $x = -2$ (rejected)

If (α, β) is the point on $y^2 = 6x$, that is closest to $\left(3, \frac{3}{2}\right)$ then find $2(\alpha + \beta)$

(1)6

(2)9

(3)7

(4)5

Ans. (2)

Sol.

(3,3/2)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



$$y^2 = 6x$$

$$2yy' = 6$$

$$-\frac{\beta}{3} = \frac{\beta - 3/2}{\alpha - 3}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : core DECO at EGG77 | Website : unit

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution pontal

RESONANCE* | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

$$-\frac{\beta}{3} = \frac{2\beta - 3}{2\alpha - 6}$$

$$-\beta(2\alpha-6)=6\beta-9$$

$$6\beta - 2\alpha\beta = 6\beta - 9$$

$$\alpha\beta = \frac{9}{2} \Rightarrow \beta = \frac{9}{2\alpha}$$

$$\alpha^3 = \frac{27}{8} \quad \alpha = \frac{3}{2} \ , \beta^2 = 9 \ \Rightarrow \beta = \pm 3$$

$$\alpha = \frac{3}{2}, \beta = 3$$

$$2(\alpha + \beta) = 9$$

11. Two circles pass through (-1,4) and their centres lie on $x^2 + y^2 + 2x + 4y = 4$ r. If r_1 and r_2 are maximum 4 minimum radii and $\frac{r_i}{r} = a + b\sqrt{2}$ then the value of a + b is

Ans. 3

Sol. Given circle

$$(x + 1)^2 + (y + 2)^2 = (3)^2$$

any point on this circle is $(3\cos\theta - 1, 3\sin\theta 2)$ equation of circle having centre $(3\cos\theta - 1, 3\sin\theta - 2)$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$r = \sqrt{(3\cos\theta - 1 + 1)^2 + (3\sin\theta - 2 - 4)^2}$$

$$= \sqrt{9\cos^2\theta + 9\sin^2\theta + 36 - 36\sin\theta}$$

 \Rightarrow r_{max} = 9 = r₁ and r_{min} = 3 = r₂

$$\Rightarrow \frac{\mathbf{r}_1}{\mathbf{r}_2} = \frac{9}{3} = 3 \ 3 + 0\sqrt{2}$$

$$\Rightarrow$$
 a + b = 3

12. If \triangle ABC is right angled triangle with sides $\frac{a}{b}$ & c and smallest angle $\frac{a}{b}$. If $\frac{1}{a}$, $\frac{1}{b}$ and $\frac{1}{c}$ are also the sides of right angled triangle then find $\sin \theta$

(1)
$$\sqrt{\frac{3-\sqrt{5}}{2}}$$

(2)
$$\frac{3-\sqrt{5}}{2}$$

(1)
$$\sqrt{\frac{3-\sqrt{5}}{2}}$$
 (2) $\frac{3-\sqrt{5}}{2}$ (3) $\sqrt{\frac{3+\sqrt{5}}{2}}$ (4) $\frac{3+\sqrt{5}}{2}$

(4)
$$\frac{3+\sqrt{5}}{2}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

nesonance Euuventures Ltu.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555 S 7340010333 ecolosic configuration of the contact of the conta

This solution was download from Resonance JEE (MAIN) 2021 Solution portal



Resonance* | Jee Main-2021 | Date : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

Sol. Let a > b > c

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029



$$in\theta = \frac{c}{a}$$

$$\frac{1}{a} < \frac{1}{b} < \frac{1}{c}$$

$$\frac{1}{c^2} = \frac{1}{a^2} + \frac{1}{b^2}$$

$$1 = \frac{c^2}{a^2} + \frac{c^2}{b^2}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$1 = \sin^2\theta + \frac{1}{\frac{a^2}{c^2} - 1} = \sin^2\theta + \frac{1}{\cos^2\theta - 1}$$

$$1 = \frac{1 - \sin^2 \theta + 1}{\cos \theta c^2 \theta - 1} \Rightarrow \sin^2 q + \csc^2 \theta = 3$$

13. If Re $[(1+\cos\theta + 2i\sin\theta)^{-1}] = 4$ then value of θ is :

$$(1) \frac{\pi}{2}$$

(2)
$$\frac{\pi}{3}$$

(3)
$$-\frac{\pi}{2}$$

Ans.

Sol.
$$\frac{1}{1+\cos^2\theta + 2\sin\theta} \times \frac{1+\cos\theta - 2\sin\theta}{1+\cos\theta - 2\sin\theta}$$
$$= \frac{1+\cos\theta - 2\sin\theta}{(1+\cos\theta)^2 + 4\sin^2\theta}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$1 + \cos^2 \theta + 2\cos \theta + 4\sin^2 \theta$$

$$\Rightarrow \frac{1 + \cos \theta}{1 + \cos^2 \theta + 2\cos \theta + 4 - 4\cos^2 \theta} = 4$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
besteek own fleeconceds: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
weekek own fleeconceds: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow \frac{1 + \cos \theta}{5 + 2\cos \theta - 3\cos^2 \theta} = 4$$

$$\Rightarrow 1 + \cos\theta = 20 + 8\cos\theta - 12\cos^2\theta$$

$$\Rightarrow 12\cos^2\theta - 7\cos\theta - 19 = 0$$

$$\Rightarrow 12\cos^2\theta - 19\cos\theta + 12\cos\theta - 19 = 0$$

$$\Rightarrow$$
 cos θ (12cos θ - 19) + 1 (12cos θ -19) = 0

$$\Rightarrow$$
 cosθ = -1 or cosθ = $\frac{19}{12}$ (rejected)

$$\Rightarrow \theta = \pi$$

If x = ay -1 = z -2, and x = 3y -2 = bz - 2 lie in same plane then the value of a, b, is

$$(1) a = 2 b = 3$$

(1)
$$a = 2$$
, $b = 3$ (2) $a = 1$, $b = 1$ (3) $b = 1$, $a \in R - \{0\}$

Ans. (3)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

a
$$(\vec{a}_1 - \vec{a}_2)(\vec{b}_1 \times \vec{b}_2) = 0$$
 $0 \quad \frac{1}{a} - \frac{2}{3} \quad 2 - \frac{2}{b}$ $0 \quad \frac{1}{a} = 0$

$$\Rightarrow \frac{1}{1} - \frac{1}{1} = 0$$

$$b = 1, a \in R - \{0\}$$

15. If
$$P(\overline{A} \cap B) + P(A \cap \overline{B}) = 1 - K$$

$$P(\overline{A} \cap C) + P(A \cap \overline{C}) = 1 - 2K$$

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 27777777, 2777770 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$P(A \cap B \cap C) = K^2, K \in (0,1)$$

Then the value of P(at least one of A,B,C) is:

$$(1) > \frac{1}{-}$$

$$(2)$$
 $\left[\frac{1}{2}, \frac{1}{4}\right]$

$$(4)^{-1}$$

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555 S 7340010333 toeteek.com/leconancedu my hetter.com/leconancedu www.youtube.com/retowate

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#7

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Resonance* | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

Sol. $P(A) + P(B) - 2P(A \cap B) = 1 - K$

 $P(A) + P(C) - 2P(A \cap C) = 1 - 2K$

 $P(B) + P(C) - 2P(B \cap C) = 1 - K$

 $P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$

$$= \frac{3-4k}{2} + k^2 = \frac{2k^2 - 4k + 3}{2}$$

.. The value of 2k2 -4k +3 is greater than 1

$$\therefore P(A \cup B \cup C) > \frac{1}{2}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

(1) -5

(2) 5

(3) 6

(4)-6

Ans. (1)

Sol. $f(f(x)) = \frac{5f(x) + 3}{6f(x) + a} = x \Rightarrow 5f(x) + 3 = 6x f(x) + ax$

 $\Rightarrow \frac{25x+15}{6x+a} + 3 = 6x \left(\frac{5x+3}{6x+a}\right) + ax$

 \Rightarrow 25x+ 15 + 18x +3a = 30x² + 18x + 6ax² + a²x

 \Rightarrow (30+6a)x² + (a² - 25) x - (3a+15) = 0

 \Rightarrow 6(a+5)x² + (a-5) (a+5)x-3 (a+5) = 0, $\forall x$

 \Rightarrow a+5= 0 \Rightarrow a = -5


17. If $g(t) = \begin{cases} max(t^3 - 6t^2 + 9t - 3.0), t \in [0.3] \\ 4 - t, t \in (3,4] \end{cases}$ then the number of points at which g(t) is non differentiable is :

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$y = t^3 - 6t^2 + 9t - 3$$

$$y' = 3t^2 - 12t + 9$$

$$=3(t^2-4t+3)$$



Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

DAGE#8

Resonance | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

A: if 2 + 4 = 7, then 3+4 = 8

B: if 3 + 5 = 8, then earth is flat

C: if A and B are true, then 5+4=11

(1) A is true, B and C are false

(2) B is true, A and C are false

(3) C is true, A and B are false

(4) B is false, A and C are true

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

р	q	$p \rightarrow q$
Т	T	T
Т	F	F
F	T	T
F	F	Т

A is true, B is false, C is true.

19. If
$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$
, $B \sum_{t=1}^{2021} A^t$ then value of $|B|$ is

(1)2021

(2) (2021)2

(3) - 2021

(4) 0

Ans. (2)

Sol. A=I, B = I + I +..... 2021 times

B - 202 1 0

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

|B| = (2021)*

Resonance Eduventures Ltd.


Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
acetesk com/Resonancedul www.resonancedul www

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#9



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



COMPUTER BASED TEST (CBT) Memory Based Questions & Solutions

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



5 AIRs in TOP-50 in JEE (Adv.) 2020 from Classroom

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302R.J2007PLC024029

ADMISSION OPEN for Session 2021-22

ONLINE + OFFLINE PROGRAMS

CLASS 11, 12 & 12+

Target: JEE (Main+Adv.) | JEE (Main) | NEET

Scholarship Upto 90%*

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Resonance* | Jee Main-2021 | Date : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

PART : MATHEMATICS

1 If element of matrix A is defined as A = $[a_{ij}]_{3\times3}$ where A = $\begin{cases} (-1)^{j+i} & i < j \\ 2 & i = j \\ (-1)^{j+j} & i > j \end{cases}$, then the value of $|3Adj(2A^{-1})|$

(1) 72 (2) 36 (3) 108 (4) 48

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$A = \begin{bmatrix} -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$$

So, $|A| = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$
 $= 2(4-1) + 1(-2+1) + 1(1-2)$
 $= 2(3) + 1(-1) + 1(-1)$
 $= 4$
 $|3Ac|(2A^{-1})| = 3^3 |Ac|(2A^{-1})| = 3^3 \times |2A^{-1}|^2$
 $= 3^3 \times 2^6 \times |A^{-1}|^2 = 3^3 \times 2^6 \times \frac{1}{|A|^2} = 108$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$|\overrightarrow{AB}| = 7$$
, $|\overrightarrow{BC}| = 5$, $|\overrightarrow{CA}| = 3$,



Projection of BC on CA is = |BC| cos∠BCA

$$5\left(\frac{3^2+5^2-7^2}{2.3.5}\right) = 5\left|\frac{-15}{30}\right| = \frac{5}{2}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302R.J2007PLC024029

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

RESONANCE* | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

- The value of $tan(2tan^{-1}(3/5) + sin^{-1}(5/13))$ is:
- (2) $\frac{110}{21}$ (3) $\frac{55}{21}$

Δne

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$\tan^{-1} \frac{5}{1 - \frac{9}{25}} + \tan^{-1} \frac{5}{12}$$

$$\tan\left(\tan^{-1}\left(\frac{15}{8}\right) + \tan^{-1}\left(\frac{5}{12}\right)\right) = \frac{\frac{15}{8} + \frac{5}{12}}{1 - \frac{15}{8} \cdot \frac{5}{12}} = \frac{220}{21}$$

(2) 1

Mean of 6 observations is 10 and their variance is $\frac{20}{3}$. If observations are 15, 11, 10, 7, a, b then

|a - b| is equal to:

(3) 3

(1)2Ans

Mean = 10

$$\frac{7+10+11+15+a+b}{6}=10$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$\frac{49+100+121+225+a^2+b^2}{6}-100=\frac{2}{3}$$

$$a^2 + b^2 = 145$$

 $(a + b)^2 = 289$

ab = 72

$$(a - b)^2 = (a + b)^2 - 4ab$$

$$(a - b)^2 = 289 - 288 = 1$$

a -b | = 1

- If f(x) = x + 1, then find $\lim_{n \to \infty} \frac{1}{n} \left[f(0) + f\left(\frac{5}{n}\right) + f\left(\frac{10}{n}\right) + \dots + f\left(\frac{5(n-1)}{n}\right) \right]$

Ans.

Sol. =
$$\lim_{r \to \infty} \frac{1}{r} \int_{r}^{r-1} f(\frac{5r}{r}) = \int_{r}^{1} f(\frac{5r}{r}) dr = \int_{r}^{1} f(\frac{5r}{r})$$

$$=\left(\frac{5x^2}{2} + x\right)_0^1 = \frac{5}{2} + 1 = \frac{7}{2}$$
 Ans

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555
7840010333
Toolkok contresonance ac.in | www.resonance ac.in | www.youtube.com/resonance ac.in | CIN: U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

M Posnoance*

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777; 2777700 | contact@resonance ac in | CIN - U80302RJ2007PLC024029

Ans. (3)

Sol. 2 log₉x +3 log₉x +4 log₉x21 terms

=
$$(2 + 3 + 4 + 5 + \dots + 22) \log_9 x = \frac{21}{2} (2 + 22) \log_9 x$$

= 21 x 12 logsx

= 252 log9x

Given sum = 252 ⇒ logex = 1

$$\Rightarrow$$
 x = 9

 $([x] - [\sin x]) dx = ?$ (Where [-] represents G.I.F.)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$I = \int_{-\pi}^{\frac{\pi}{2}} ([x] - [\sin x]) dx$$

using property
$$\int_{-a}^{a} f(x)dx = \int_{0}^{a} f(x)dx + \int_{0}^{a} f(-x)dx$$

$$I = \int_{0}^{\frac{\pi}{2}} ([x] + [-x]) dx - \int_{0}^{\frac{\pi}{2}} ([\sin x] + [-\sin x]) dx = 0$$

8. If
$$\lim_{x\to 0} \frac{\alpha x e^x - \beta \ell n(1+x) + \gamma x^2 e^{-x}}{x^3} = 10$$
, then the value of $\alpha + \beta + \gamma$ is :

Ans.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow \alpha - \beta = 0, \Rightarrow \alpha = \beta$$

$$\Rightarrow \alpha + \frac{\beta}{2} + \gamma = 0 \Rightarrow \gamma = -\frac{3\beta}{2}$$

$$\Rightarrow \frac{\alpha}{2} - \frac{\beta}{3} - \gamma = 10$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555
7340010333
acceptable better conflection of the contact of the

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

RESONANCE* | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

$$\Rightarrow \frac{\beta}{2} - \frac{\beta}{3} + \frac{3\beta}{2} = 10 \Rightarrow \frac{3\beta - 2\beta + 9\beta}{6} = 10$$

:
$$\beta = 6$$
, $\alpha = 6$, $\gamma = -9$

So, the value of
$$\alpha + \beta + \gamma = 3$$

The value of x satisfying the equation $log_{(x+1)}(2x^2 + 7x + 5) + log_{(2x+5)}(x+1)^2 = 4$ is:

$$(3) -4$$

Ans. (2)

 $\log_{(x+1)}((2x+5)(x+1)) + \log_{(2x+5)}(x+1)^2 = 4$ Sol.

$$1 + \log_{(x+1)} (2x+5) + 2\log_{(2x+5)} (x+1) = 4$$

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$t^2 + t + 2 = 4t \Rightarrow t^2 - 3t + 2 = 0$$

For
$$t = 2$$

 $2x+5 = (x+1)^2$

$$\Rightarrow$$
 x = -4 (rejected)

$$x = 2$$
, $x = -2$ (rejected)

If (α, β) is the point on $y^2 = 6x$, that is closest to $\left(3, \frac{3}{2}\right)$ then find $2(\alpha + \beta)$

(1)6

(2)9

(3)7

(4)5

Ans. (2)

Sol.

(3,3/2)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



$$y^2 = 6x$$

$$2yy' = 6$$

$$-\frac{\beta}{3} = \frac{\beta - 3/2}{\alpha - 3}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : core DECO at EGG77 | Website : unit

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution pontal

RESONANCE* | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

$$-\frac{\beta}{3} = \frac{2\beta - 3}{2\alpha - 6}$$

$$-\beta(2\alpha-6)=6\beta-9$$

$$6\beta - 2\alpha\beta = 6\beta - 9$$

$$\alpha\beta = \frac{9}{2} \Rightarrow \beta = \frac{9}{2\alpha}$$

$$\alpha^3 = \frac{27}{8} \quad \alpha = \frac{3}{2} \ , \beta^2 = 9 \ \Rightarrow \beta = \pm 3$$

$$\alpha = \frac{3}{2}, \beta = 3$$

$$2(\alpha + \beta) = 9$$

11. Two circles pass through (-1,4) and their centres lie on $x^2 + y^2 + 2x + 4y = 4$ r. If r_1 and r_2 are maximum 4 minimum radii and $\frac{r_i}{r} = a + b\sqrt{2}$ then the value of a + b is

Ans. 3

Sol. Given circle

$$(x + 1)^2 + (y + 2)^2 = (3)^2$$

any point on this circle is $(3\cos\theta - 1, 3\sin\theta 2)$ equation of circle having centre $(3\cos\theta - 1, 3\sin\theta - 2)$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$r = \sqrt{(3\cos\theta - 1 + 1)^2 + (3\sin\theta - 2 - 4)^2}$$

$$= \sqrt{9\cos^2\theta + 9\sin^2\theta + 36 - 36\sin\theta}$$

 \Rightarrow r_{max} = 9 = r₁ and r_{min} = 3 = r₂

$$\Rightarrow \frac{\mathbf{r}_1}{\mathbf{r}_2} = \frac{9}{3} = 3 \ 3 + 0\sqrt{2}$$

$$\Rightarrow$$
 a + b = 3

12. If \triangle ABC is right angled triangle with sides $\frac{a}{b}$ & c and smallest angle $\frac{a}{b}$. If $\frac{1}{a}$, $\frac{1}{b}$ and $\frac{1}{c}$ are also the sides of right angled triangle then find $\sin \theta$

(1)
$$\sqrt{\frac{3-\sqrt{5}}{2}}$$

(2)
$$\frac{3-\sqrt{5}}{2}$$

(1)
$$\sqrt{\frac{3-\sqrt{5}}{2}}$$
 (2) $\frac{3-\sqrt{5}}{2}$ (3) $\sqrt{\frac{3+\sqrt{5}}{2}}$ (4) $\frac{3+\sqrt{5}}{2}$

(4)
$$\frac{3+\sqrt{5}}{2}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

nesonance Euuventures Ltu.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555 S 7340010333 ecolosic configuration of the contact of the conta

This solution was download from Resonance JEE (MAIN) 2021 Solution portal



Resonance* | Jee Main-2021 | Date : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

Sol. Let a > b > c

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029



$$in\theta = \frac{c}{a}$$

$$\frac{1}{a} < \frac{1}{b} < \frac{1}{c}$$

$$\frac{1}{c^2} = \frac{1}{a^2} + \frac{1}{b^2}$$

$$1 = \frac{c^2}{a^2} + \frac{c^2}{b^2}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$1 = \sin^2\theta + \frac{1}{\frac{a^2}{c^2} - 1} = \sin^2\theta + \frac{1}{\cos^2\theta - 1}$$

$$1 = \frac{1 - \sin^2 \theta + 1}{\cos \theta c^2 \theta - 1} \Rightarrow \sin^2 q + \csc^2 \theta = 3$$

13. If Re $[(1+\cos\theta + 2i\sin\theta)^{-1}] = 4$ then value of θ is :

$$(1) \frac{\pi}{2}$$

(2)
$$\frac{\pi}{3}$$

(3)
$$-\frac{\pi}{2}$$

Ans.

Sol.
$$\frac{1}{1+\cos^2\theta + 2\sin\theta} \times \frac{1+\cos\theta - 2\sin\theta}{1+\cos\theta - 2\sin\theta}$$
$$= \frac{1+\cos\theta - 2\sin\theta}{(1+\cos\theta)^2 + 4\sin^2\theta}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$1 + \cos^2 \theta + 2\cos \theta + 4\sin^2 \theta$$

$$\Rightarrow \frac{1 + \cos \theta}{1 + \cos^2 \theta + 2\cos \theta + 4 - 4\cos^2 \theta} = 4$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
besteek own fleeconceds: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
weekek own fleeconceds: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow \frac{1 + \cos \theta}{5 + 2\cos \theta - 3\cos^2 \theta} = 4$$

$$\Rightarrow 1 + \cos\theta = 20 + 8\cos\theta - 12\cos^2\theta$$

$$\Rightarrow 12\cos^2\theta - 7\cos\theta - 19 = 0$$

$$\Rightarrow 12\cos^2\theta - 19\cos\theta + 12\cos\theta - 19 = 0$$

$$\Rightarrow$$
 cos θ (12cos θ - 19) + 1 (12cos θ -19) = 0

$$\Rightarrow$$
 cosθ = -1 or cosθ = $\frac{19}{12}$ (rejected)

$$\Rightarrow \theta = \pi$$

If x = ay -1 = z -2, and x = 3y -2 = bz - 2 lie in same plane then the value of a, b, is

$$(1) a = 2 b = 3$$

(1)
$$a = 2$$
, $b = 3$ (2) $a = 1$, $b = 1$ (3) $b = 1$, $a \in R - \{0\}$

Ans. (3)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$\begin{array}{c}
\overline{a} \\
(\vec{a}_1 - \vec{a}_2)(\vec{b}_1 \times \vec{b}_2) = 0 \\
0 \quad \frac{1}{a} - \frac{2}{3} \quad 2 - \frac{2}{b} \\
1 \quad \frac{1}{a} \quad 1 \\
1 \quad \frac{1}{3} \quad \frac{1}{b}
\end{array} = 0$$

$$\Rightarrow \frac{1}{ab} - \frac{1}{a} = 0$$

$$b = 1, a \in R - \{0\}$$

15. If
$$P(\overline{A} \cap B) + P(A \cap \overline{B}) = 1 - K$$

$$P(\overline{A} \cap C) + P(A \cap \overline{C}) = 1 - 2K$$

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 27777777, 2777770 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$P(A \cap B \cap C) = K^2, K \in (0,1)$$

Then the value of P(at least one of A,B,C) is:

$$(1) > \frac{1}{-}$$

$$(2)$$
 $\left[\frac{1}{2}, \frac{1}{4}\right]$

$$(4)^{-1}$$

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555 S 7340010333 toeteek.com/leconancedu my hetter.com/leconancedu www.youtube.com/retowate

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#7

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Resonance* | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

Sol. $P(A) + P(B) - 2P(A \cap B) = 1 - K$

 $P(A) + P(C) - 2P(A \cap C) = 1 - 2K$

 $P(B) + P(C) - 2P(B \cap C) = 1 - K$

 $P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$

$$= \frac{3-4k}{2} + k^2 = \frac{2k^2 - 4k + 3}{2}$$

.. The value of 2k2 -4k +3 is greater than 1

$$\therefore P(A \cup B \cup C) > \frac{1}{2}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

(1) -5

(2) 5

(3) 6

(4)-6

Ans. (1)

Sol. $f(f(x)) = \frac{5f(x) + 3}{6f(x) + a} = x \Rightarrow 5f(x) + 3 = 6x f(x) + ax$

 $\Rightarrow \frac{25x+15}{6x+a}+3=6x\left(\frac{5x+3}{6x+a}\right)+ax$

 \Rightarrow 25x+ 15 + 18x +3a = 30x² + 18x + 6ax² + a²x

 \Rightarrow (30+6a)x² + (a² - 25) x - (3a+15) = 0

 \Rightarrow 6(a+5)x² + (a-5) (a+5)x-3 (a+5) = 0, $\forall x$

 \Rightarrow a+5= 0 \Rightarrow a = -5


17. If $g(t) = \begin{cases} max(t^3 - 6t^2 + 9t - 3.0), t \in [0,3] \\ 4 - t, t \in (3,4] \end{cases}$ then the number of points at which g(t) is non differentiable is :

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$y = t^3 - 6t^2 + 9t - 3$$

$$y' = 3t^2 - 12t + 9$$

$$=3(t^2-4t+3)$$



Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#8

Resonance | JEE MAIN-2021 | DATE : 20-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

A: if 2 + 4 = 7, then 3+4 = 8

B: if 3 + 5 = 8, then earth is flat

C: if A and B are true, then 5+4=11

(1) A is true, B and C are false

(2) B is true, A and C are false

(3) C is true, A and B are false

(4) B is false, A and C are true

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

р	q	$p \rightarrow q$
Т	T	T
Т	F	F
F	T	T
F	F	Т

A is true, B is false, C is true.

19. If
$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$
, $B \sum_{t=1}^{2021} A^t$ then value of $|B|$ is

(1)2021

(2) (2021)2

(3) - 2021

(4) 0

Ans. (2)

Sol. A=I, B = I + I +..... 2021 times

B - 202 1 0

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

|B| = (2021)*

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
acetesk com/Resonancedul www.resonancedul www

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#9