



COMPUTER BASED TEST (CBT) Memory Based Questions & Solutions

Date: 27 July, 2021 (SHIFT-1) | TIME: (9.00 a.m. to 12.00 p.m)

 $@ \ Resonance \ Eduventures \ Limited \ | \ Toll Free \ 1800 \cdot 258 \cdot 5555 \ | \ (0)744 \ 27777777, 27777700 \ | \ contact @ resonance.ac. in \ | \ CIN - U80302RJ2007PLC024029 \ | \ contact = 1800 \cdot 258 \cdot 258$

Resonance Eduventures Ltd. Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222 To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029 Toll Free: 1800 258 5555 | 7340010333 | tectoric contributions | tector

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

PART: PHYSICS

In the given arrangement, block A of mass m moving with speed 9m/s, strikes with block B of mass 2m elastically. Then block B strikes with block C of mass 2m perfect in-elastically. Find the final speed of block C?

Α		В	С
m	9 m/s	2m	2m

(1) 3 m/s (2) 4 m/s (3) 5 m/s (4) 6 m/s

Ans. (1)

Sol. $9m = mv_A + 2mv_B$

$$9 = V_A + 2V_B$$
(1)

$$9 = v_B - v_A \qquad \dots (2)$$

Thus v_B = 6 m/s

Callinian batuman D. C.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

vc = 3 m/s

A particle is thrown upward at t = 0. It attains maximum height of h. It is found at height h/3 at t = t₁ & t = t₂ Find t₁/t₂.

(1)
$$\frac{\sqrt{3}-1}{\sqrt{3}+1}$$


 $(2)\frac{1}{3}$

(3)
$$\frac{\sqrt{6}-\sqrt{5}}{\sqrt{6}+\sqrt{5}}$$

 $(4) - \frac{1}{2}$

Ans. (3)

Sol.



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$V = \sqrt{2gh}$$

$$\frac{h}{3} = \sqrt{2ght} - \frac{1}{2}gt^2$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39157222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302812007PLC024029

Tall Free: 1800 258 5555 | 340010333 | socioik own/lesonancetal | witter.com/lesonancetal | witter.com/lesonancetal | witter.com/lesonancetal | witter.com/lesonancetal | description |

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#1

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Resonance* | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

$$gt^2 - 2\sqrt{2ght} + \frac{2h}{3} = 0$$

$$t = \frac{2\sqrt{2gh} \pm \sqrt{8gh - 4g \times \frac{2h}{3}}}{2g} = \frac{2\sqrt{2gh} \pm \sqrt{\frac{16gh}{3}}}{2g}$$

$$=\frac{2\sqrt{2gh}\pm4\sqrt{\frac{gh}{3}}}{2g}$$

$$\frac{t_1}{t_2} = \frac{2\sqrt{2gh} - 4\sqrt{\frac{gh}{3}}}{2\sqrt{2gh} + 4\sqrt{\frac{gh}{3}}}$$

$$2\sqrt{2} - \frac{4}{\sqrt{3}}$$
 $\sqrt{3} - \sqrt{2}$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- In YDSE, if the light-used is shifted from orange to blue then which of the following is correct.
 - (1) Fringe width first increases then decreases (2) Fringe width increase
 - (3) Fringe width decreases
- (4) Fringe width first decreases than increases

Ans. (3)

c-l α λD

4. In the circuit shown in the figure, S₁ remains closed for a long time and S₂ remains open. Now S₂ is closed and S₁ is opened. Find out the di/dt just after that moment.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



$$(2) - \frac{6\epsilon}{1}$$

$$(3) - \frac{2\epsilon}{1}$$

$$(4) - \frac{\varepsilon}{1}$$

Ans. (2)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222


This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#2

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

RESONANCE® | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

Sol. Before S_2 is closed and S_1 is opened current in the left part of the circuit = $\frac{\epsilon}{R}$. Now when S_2 closed S_1 opened, current through the inductor cannot change suddenly, current $\frac{\epsilon}{R}$ will continue to move in the



Applying KVL in loop 1.

$$L \frac{di}{dt} + \frac{\varepsilon}{R}(2R) + 4\varepsilon = 0$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

5. Two identical tennis balls of mass m and charge q are hinged by a common support with the help of a string of length '\ell'. If the system is in equilibrium, then find the distance between the balls? Ignore gravitational interactions between balls. (6 is very small)


$$(1) \left(\frac{kq^2 \ell}{mg} \right)^{1/2}$$

$$(2) \left(\frac{3kq^2\ell}{mq} \right)^1$$

$$(3) \left(\frac{2kq^2\ell}{mg}\right)^{1/3}$$

$$(4) \left(\frac{3kq^2 \ell}{2mg} \right)^{1/3}$$

Ans. (3)



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$tan\theta = \frac{kq^2}{mg} \approx sin\theta$$
$$kq^2 = d$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555
7340010333
Toottok contresonanced: www.resonanced. w

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#3

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance* | Jee Main-2021 | Date : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

For the given semicircle with centre O. choose the correct relation? If A, B, C & D are points on the semicircle such that $|\overrightarrow{AB}| = |\overrightarrow{BC}| = |\overrightarrow{CD}|$.



Ans. (1)

Cal For the aliver comisisale

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

AD = Diameter of circle

7. A monoatomic gas is kept in a 1 litre container at pressure 1 atm. If average energy per molecule is 2×10^{-9} J, find no. of molecules in the container:

 $(1) 0.75 \times 10^{-6}$

- $(2) 0.75 \times 10^9$
- $(3) 0.5 \times 10^{11}$
- $(4) \ 0.75 \times 10^{11}$

Ans. (4)

Sol. Total energy = $\frac{f}{2}$ nRT = $\frac{3}{2}$ PV

According to question $\frac{3}{2}PV = N \times 2 \times 10^{-9} J$

$$\frac{3}{2} \times 10^5 \times 1000 \times 10^{-6} = N \times 2 \times 10^{-9}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U90302RJ2007PLC024029

$$N = \frac{3}{4} \times \frac{10^{-9}}{10^{-9}} = \frac{3}{4} \times 10^{11} = 0.75 \times 10^{1}$$

8. The relative permittivity of distilled water is 81. The velocity of light in it will be $(\mu_r = 1)$

 $(1) 3.3 \times 10$

- (2) 5.3 × 10
- $(3) 4.3 \times 10^{7}$
- (4) None

Ans. (1)

Sol.
$$V = \frac{C}{\sqrt{\mu_r \epsilon_r}}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-45 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PL C024029

Tall Free: 1800 258 5555 | 7340010333 | sectors our Personancel | www.youtde.com/resonance.ac.in | Cin: U80302RJ2007PL C024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

RESONANCE | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

Match the following column if all the rods have same density and same radius.




12



(b)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

(d)



- (1) (i) b (ii) d (iii) a (iv) c
- (2) (i) c (ii) d (iii) a (iv) d
- (3) (i) a (ii) c (iii) b (iv) d
- (3) (i) d (ii) a (iii) c (iv) b

- Ans.
- Sol.

(a)

(b)

- © Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN U80302RJ2007PLC024029
 - A particle executing SHM having maximum kinetic energy 'E' and amplitude 'A'. Find displacement from mean position when its kinetic energy is $\frac{3E}{4}$:
 - $(1)\frac{A}{2}$

Ans. (1)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

RESONANCE | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

Sol.
$$V = A\omega$$
, $KE_{max} = \frac{1}{2}m\omega^2 A^2$
 $V = \omega \sqrt{A^2 - x^2}$ $KE_X = \frac{1}{2}m\omega^2 (A^2 - x^2)$


$$\frac{3}{4}KE_{max} = KE_{3}$$

$$\frac{3}{4}\frac{1}{2}m\omega^{2}A^{2} = \frac{1}{2}m\omega^{2}(A^{2} - x^{2})$$

$$x^2 = \frac{A^2}{4}$$

$$x = \pm \frac{A}{2}$$


For given PV diagram AB is isothermal at T1 temperature, CD is also isothermal at T2 temperature. given that T1 > T2. Then which one of the following is correct.



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Ans.

12. A container of area 'A' filled with liquid, a small hole of area 'a' is made at the bottom of curved surface as shown. Find the coefficient of friction between container and ground in order to prevent it from sliding.



Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#6

| JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

Ans.

Force exerted by leaving water on the container, Sol.

$$F = \begin{pmatrix} dm \\ ot \end{pmatrix} v = (\rho a v) v$$

By conservation of energy v = 2gh

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\mu\rho(Ah)g=\rho a2gh$$

$$\mu = \frac{2a}{\Delta}$$

Two solid spheres of radius 4R and R and having same mass density ρ kept at a distance of 8 R. Find Moment of Inertia of the system about the axis passes through centre of line joining their centres.

(1)
$$\frac{9660}{5} \pi \rho R^5$$

(3) $\frac{9664}{4} \pi \rho R^5$ (4) $\frac{9664}{3} \pi \rho R^5$

Ans.

Sol.



/! 40 !

Let mass density = ρ

$$m_2 = \frac{4}{3} \pi R^3 \rho$$

$$m_1 = \frac{4}{3}\pi (4R)^3 \rho$$

$$m_1 = 64 m_2$$

$$I_{final} = [I_2 + m_2 (4R)^2] + [I_1 + m_1(4R)^2]$$

$$= \frac{2}{5} m_2 R^2 + 16 m_2 R^2 + \frac{2}{5} m_1 (4R)^2 + m_1 (4R)^2$$

$$=\frac{2}{5}m_{2}R^{2}+16m_{2}R^{2}+\frac{2}{5}64m_{2}(4R)^{2}+64m_{2}(4R)^{2}=\frac{5800}{3}\pi\rho R^{5}$$

Decemens Eduventures Ltd

@ Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029


Toll Free: 1800 258 5555
7340010333
toll sectors confidenced: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
toll sectors confidenced: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

RESONANCE | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

For the given circuit, switch is closed at t = 0 find time after which voltage across capacitor becomes 50V



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- (3) 50 μsec (1) 69.3 µsec (2) 60 µsec
 - (4) 40 µsec

Ans. (1)


Sol.
$$V = V_0 e^{\frac{t}{RC}}$$

$$50 = 100e^{\frac{t}{RC}}$$

$$t = RC \ln 2$$

= 100 × 10⁻⁶ $\ln 2 = 10^{-4} \times 0.693$ sec = 69.3 µsec

Figure shows two capacitors in steady state. Now the cell is removed and a dielectric of dielectric constant k is inserted between the plates of the capacitor of capacitance C. Find new potential difference across any of the capacitors:



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777770 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$(1) \frac{V}{2+k}$$

$$(2) \frac{2V}{2}$$

(4)
$$\frac{V}{1+2k}$$

Ans. Sol.

Just after removing cell



After inserting dielectric



Conservation of charge

$$V_{ab} = \frac{3CV}{2C + CK}$$

Mesonance Eduventures Etc.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029 Toll Free : 1800 258 5555 🚳 7340010333 💥 tectoek own/liescharcelds 💆 witter con/liescharcelds 🖺 www.youtsie.com/recovert. 🛅 bits resmance.ac. in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#8

RESONANCE | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

16. A particle of mass 9.1 × 10⁻³¹ kg is moving with velocity 6 m/s. Momentum of photon is 2 × 10⁻²⁷ kg m/s. If de-Broglie wavelength of the particle is K times of the wavelength of photon. Find value of K:

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol. De-Broglie wavelength of particle

$$\lambda = \frac{h}{p} = \frac{h}{9.1 \times 10^{-31} \times 6}$$

Wavelength of photon

$$\lambda_{\text{photon}} = \frac{h}{p} = \frac{h}{2 \times 10^{-27}}$$

From question

$$\Rightarrow K = \frac{2 \times 10^{-27}}{9.1 \times 10^{-31} \times 6} = 366$$

17. Two prism of same angle of refraction are arranged as shown. If the light incident, on the system goes undeviated. Then find wave length of incident light

@ Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\eta_2 = 1.45 + \frac{1.8 \times 10^{-14}}{\lambda^2}$$



- (1) 600 nm
- (2) 675 nm
- (3) 575 nm
- (4) 475 nm

Ans.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
tooltox cont Resonanceds: with twitter contribution of the co

This solution was download from Resonance JEE (MAIN) 2021 Solution portal



$$\delta_{net} = A(n_1 - 1) - A(n_2 - 1) = 0$$

$$\Rightarrow 1.2 + \frac{10.8 \times 10^{-14}}{\lambda^2} = 1.45 + \frac{1.8 \times 10^{-14}}{\lambda^2}$$

$$\lambda^2 = \frac{(10.8 - 1.8) \times 10^{-14}}{(1.45 - 1.2)}$$

 $\lambda = 600 \text{ nm}.$

For the given circuit find current 'I' and the phase difference between V and I:



(1) 8.8A, tan-1 1.83

(2) 6.8A, tan⁻¹ 2.83 (3) 5.8A, tan⁻¹ 0.83 (4) 7.8A, tan⁻¹ 2.53

Ans. (1)

Sol. $\omega = 2\pi f = 100\pi$

Thus
$$x_L = \omega_L = 100 \times \frac{22}{7} \times 0.07 = 22\Omega$$

Thus, $z = \sqrt{R^2 + x_L^2}$

 $z = \sqrt{(12)^2 + (22)^2}$

Z ≈ 25.00

220

 $X_L = 22$ Z = 25R=12

Impedance triangle

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555 | 7340010333 | sectors confidence of the confidence of the


This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 10

RESCHARCE | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

@ Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

the velocity of projection such that particle reaches to another planet 'B', which is at a distance of 8R from planet 'A'. Planet 'B' has mass 9M and radius 2R.



Ans. (2)

Sol. Let at a distance 'X' from planet 'A', the net gravitational field becomes zero

$$\frac{GM}{X^2} = \frac{G \times 9M}{(8R - X)^2}$$

© Resonance Eduventures Limited | Toll-Free 1800-250-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$X = 2R$$

Now, Particle should be projected such that it covers a minimum distance of '2R'.

Thus

$$\begin{split} \frac{1}{2} \, m v^2 - \frac{GMm}{R} - \frac{G(9M)m}{7R} &= -\frac{GMm}{2R} - \frac{G(9M)m}{6R} \\ &\qquad \qquad \frac{1}{2} \, v^2 = \frac{2}{7} \, \frac{GM}{R} \implies v = \sqrt{\frac{4}{7} \, \frac{GM}{R}} \end{split}$$

20. Circular scale divisions of a screw gauge is 50. Five full rotations advances circular scale by 5mm.

Statement-1: Least count of screw gauge is 0.001 cm

- (1) Statement-1 & 2 both are true
- (2) Statement-1 & 2 both are true statement-2 is correct explant of statement-1
- (2) Statement 1 is false Statement 2 is true

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- Ans. (3)
- **Sol.** LC = 1 mm/50 = 0.02 mm

Statement -1 is wrong. Statement-2 is right

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222


To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555 Toll 740010333 Toeleek.com/Reconancelds toll www.youtube.com/recovator ac-

This solution was download from Resonance JEE (MAIN) 2021 Solution portal PAGE #

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302R.J2007PLC024029

21. Figure shows a conductor of tapered cone shape. As one goes from left to right on conductor, choose correct option



- (1) current decreases
- (2) drift velocity of electron increases
- (3) electric field inside conductor decreases
- (4) All of the above

Ans. (2)

- Sol. i = nAevd
 - .. v_d increases as area A decrease

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- 22. A body cools down from 61°C to 49°C in 4 min, and the surrounding temperature is 30°C. Find the time taken by this body to cool down from 49°C to 37°C.
- (1) 1.62 min
- (2) 1.25 min
- (3) 7.69 min
- (4) 0.92 min.

Ans. (3)

Sol.
$$\frac{61-49}{4} = K \left[\frac{61+49}{2} - 30 \right]$$
 ...(1

$$\frac{49-37}{t} = k \left[\frac{49+37}{2} - 30 \right] \qquad \dots (2)$$

Dividing (1) and (2)

$$\frac{t}{4} = \frac{25}{13}$$

$$t = \frac{4 \times 25}{13} = 7.69$$

A bar-magnet of magnetic moment 9.85 A-m2 and moment of inertia I = 10-5 kg-m2 makes

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

(Take $\pi^2 = 9.85$)

- (1) 20 µT

- (2) $25 \mu T$ (3) $16 \mu T$ (4) $10 \mu T$

(3) Ans.

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222 To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302R12007PLC024029

Toll Free: 1800 258 5555 | 7340010333 | sected cont-Reconaced: visiter.con/Reconaced: www.youtube.com/resonance.ac.in | Only 1909 | Sected cont-Reconaced: visiter.con/Reconaced: visiter.com/Reconaced: visiter.com/Reco

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

RESONANCE* | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

Sol.
$$T = 2\pi \sqrt{\frac{1}{MR}}$$

$$\frac{1}{2} = 2\pi \sqrt{\frac{10^{-5}}{9.85 \times B}} = 2\pi \sqrt{\frac{10^{-6}}{\pi^2 \times B}}$$

$$B = 16 \times 10^{-6} \text{ T}$$

- 24. If two discs are of same mass density but having different radii R and r.
 - I1 = Moment of inertia of disc of radius R, about an axis perpendicular to the plane of disc and passing through its centre.
 - I₂ = Moment of inertia of disc of radius r. about one of its diameter. Then choose correct option

$$(1) \frac{l_1}{l} = \frac{2r^2}{r^2}$$

(2)
$$\frac{l_1}{l_2} = \frac{2R^4}{r^4}$$
 (3) $\frac{l_1}{l_2} = \frac{2r^2}{R^2}$ (4) $\frac{l_1}{l_2} = \frac{2r}{R}$

(3)
$$\frac{l_1}{l_2} = \frac{2r^2}{R^2}$$

(4)
$$\frac{I_1}{I_2} = \frac{2r}{R}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$M_1 = \pi R^2 \sigma$$
, $M_2 = \pi r^2 \sigma$

$$I_1 = \frac{1}{2}M_1R^2 = \frac{\pi\sigma R^4}{2}$$

$$I_2 = \frac{1}{4}M_2r^2 = \frac{\pi\sigma r^4}{4}$$

$$\frac{I_1}{I_2} = \frac{2R^4}{r^4}$$

25. Find the equivalent capacitance of shown arrangement if all the plates are identical having surface



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029



(1) $\frac{24}{15} \frac{A \epsilon_0 d}{K}$ (2) $\frac{17}{15} \frac{A \epsilon_0 d}{K}$ (3) $\frac{15}{34} \frac{A \epsilon_0 d}{K}$ (4) $\frac{34}{15} \frac{A \epsilon_0 d}{4K}$

Ans. (3)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
Tootlerk.com/Resonancedu w outline.com/Resonancedu w outline.com/Resonance.ac.in | CIN: U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

| JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

Sol.

$$\begin{split} C_1 &= \frac{A\epsilon_0 K}{d} \ C_2 = \frac{3KA\epsilon_0}{2d} \ C_3 = \frac{5KA\epsilon_0}{3d} \\ 1 &= \frac{1}{C_{eq}} + \frac{1}{C_2} + \frac{1}{C_3} \\ &= \frac{d}{A\epsilon_0 K} + \frac{2d}{3A\epsilon_0 K} + \frac{3d}{5A\epsilon_0 K} \\ &= \frac{d}{A\epsilon_0 K} \left(1 + \frac{2}{3} + \frac{3}{5}\right) = \frac{34}{15} \frac{d}{A\epsilon_0 K} \end{split}$$

Thus 15 Ac d

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- An electron is revolving in a circular orbit of radius 0.5 m, with a speed of 2.2 x 10-6 m/s. Find equivalent 26.

- (1) 1.12×10^{-25} A (2) 2.15×10^{-25} A (3) 1.12×10^{-15} A (4) 2.15×10^{-13} A

Ans. (1)

Sol.
$$i = \frac{q}{t} = \frac{q}{2\pi r/v}$$

$$= \frac{1.6 \times 10^{-19} \times 2.2 \times 10^{-6}}{2\pi \times 0.5} = 1.12 \times 10^{-25} \,\text{A}$$

- A capacitor (C = 100 µF) discharging against a resistor R, at same time a radioactive substance decays with mean life 30 ms, if the ratio of charge on capacitor and activity of substance remain same for all the time then, find the resistance:
- (1) 300 Ω
- (2) 432Ω
- (3) 450Ω
- (4) 250 0

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$A = A_0 e^{-\lambda t}$$
 ...(2)

Divide (1) & (2)

$$\Rightarrow \frac{q}{A} = \frac{q_0}{A_0} \frac{\frac{t}{e^{RC}}}{e^{-\lambda t}} \Rightarrow -\lambda t = -\frac{-t}{RC}$$

 $RC = 1/\lambda = 30 \times 10^{-3} sec$

R = 300

Resonance Eduventures Ltd.


Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
across contact of the contact of the

PAGE # 14

This solution was download from Resonance JEE (MAIN) 2021 Solution portal



- 28. A wire of length 0.1 m and area of cross section 0.04 x 10⁻⁴ m² is stretched by 0.001 m, its young's modulus is 0.5 x 10⁹ pa. The energy stored in this wire is transferred completely as kinetic energy to a particle of mass 20 grams. Find speed of this particle.
 - (1) 0.5 m/
- (2) 1 m/s
- (3) 1.5 m/s
- (4) 10 m/s

Ans. (2)

Sol.
$$U = \frac{1}{2}Kx^2$$

$$= \frac{1}{2} \times \frac{AY}{L} \times (0.001)^2$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac in | CIN - U80302RJ2007PLC024029

$$\Rightarrow v^2 = \frac{0.04 \times 10^{-4} \times 0.5 \times 10^{3}}{0.1 \times 20 \times 10^{-3}} = 1 \text{ m/s}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.


Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222


This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 15

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029







2021

COMPUTER BASED TEST (CBT) Memory Based Questions & Solutions

Date: 27 July, 2021 (SHIFT-1) | TIME: (9.00 a.m. to 12.00 p.m)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



ADMISSION OPEN for Session 2021-22 © Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

CLASS 11, 12 & 12+

Target: JEE (Main+Adv.) | JEE (Main) | NEET

Scholarship Upto 90%*

Toll Free: 1800 258 5555 | Visit us: www.resonance.ac.in

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

| JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | CHEMISTRY

PART: CHEMISTRY

A weak acid HA of concentration 0.001 mole/litre have conductance 2 × 10-5 Scm⁻¹ and molar conductivity at infinite dilution is 190 Scm2 mole-1 then value of Ka of weak acid is [x] × 10-6, then value of x in nearest

Ans. 11.00

Sol.
$$\lambda_{M}^{C} = \frac{K \times 1000}{M} = \frac{2 \times 10^{-5} \times 10^{3}}{10^{-3}} = 20 \text{ Scm}^{3} \text{ mole}^{-1}$$

For weak acid (α) = $\frac{\lambda_M^C}{\lambda_M^\infty} = \frac{20}{190} = \frac{2}{19}$

 $Ka = \frac{C\alpha^2}{1-\alpha} \cong C\alpha^2 = 10^{-3} \times \left(\frac{2}{19}\right)^2 = 0.011 \times 10^{-3} = 11 \times 10^{-6}$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

List -I

List-II

NaOH

Acid

Be(OH)₂

Ca(OH)₂

(ii) Base

Al(OH)3 (d)

(iii) Amphoteric

Correct Matching of List-I and List -II is : (1) (a) - (ii), (b) - (iii), (c) - (ii), (d) - (iii), (e) - (i) (2) (a) - (ii), (b) - (ii), (c) - (iii), (d) - (iii), (e) - (iii) (3) (a) - (ii), (b) - (i), (c) - (iii), (d) - (ii), (e) - (i) (4) (a) - (ii), (b) - (iii), (c) - (ii), (d) - (ii), (e) - (ii) Ans. (1) Sol. Species Nature NACH Raco © Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029 (iii) Be(OH)2 Amphoteric Al(OH)3 Amphoteric (iv) B(OH)3 Acidic (v) Resonance Eduventures Ltd. Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222 To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029 Toll Free : 1800 258 5555 📓 7340010333 🚮 bootook own Resonancedul 💟 twitter som Resonancedul 🚵 www.youtube.com/resonance This solution was download from Resonance JEE (MAIN) 2021 Solution portal © Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029 Resonance | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | CHEMISTRY Lattice Parameter for a crystal Lattice is a ≠ b ≠ c and α = 90°, γ = 90 and β = 120° this represent which type of Bravais Lattice. (4) Orthorhombic (1) Monoclinic (2) Triclinic (3) Hexagonal Ans. Sol. Unit Cell Lattice Parameter Monoclinic $a \neq b \neq c \& \alpha = 90, \gamma = 90 \& \beta = 120^{\circ}$ (i) Seven Crystal System S.No. Crystal System Edge length Angles 1 Cubic a = b = c $\alpha = \beta = \gamma = 90^{\circ}$ 2 Tetragonal $a = b \neq c$ $\alpha = \beta = \gamma = 90^{\circ}$ Orthorhombic a ≠ b ≠ c $\alpha = \beta = \gamma = 90^{\circ}$ © Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029 $\neq 90^{\circ}, \neq 60^{\circ}$ $\alpha = \beta = 90^{\circ}$ **He**xagonal a=b≠c $\gamma \neq 120^{\circ}$ Rhombohedral 6 a = b = c $\alpha = \beta = \gamma \neq 90^{\circ}$ or Trigonal Triclinic a ≠ b ≠ c $\alpha \neq \beta \neq \gamma \neq 90^{\circ}$ Number of Geometrical isomers of complex's [Ni(CO)4], [PtCl₂(NH₃)₂], [RuCl₃(NH₃)₃], are respectively : (2) 2, 2, 2 (3) 0, 1, 2 (4) 0, 0, 2 Ans. (1) Number of G.I. Sol. Complex [Ni(CO)4] 0 (1) [PtCl₂(NH₃)₂] © Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac in | CIN - U80302RJ2007PLC024029 NH, NH, CI

Geometrical isomers (cis and trans) of Pt(NH₃)₂Cl₂

(e) B(UH)3

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
3 7340010333
Tooltok confesonancedu: www.resonancedu: www.resonancedu: www.youtule.com/resonancedu: www.youtule.com/resonan

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#2

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance* | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | CHEMISTRY

(3) [RuCl₃(NH₃)₃]

7

Cl Ru NH₃ H₃N Cl Ru Cl NH₃ Cl NH

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- 5. Identify the correct hybridisation and Magnetic nature of complex [MnCl_e]³-
 - (1) sp³d², Diamagnetic
- (2) sp³d², Paramagnetic
- (3) d²sp³, Paramagnetic
- (4) d²sp³, Diamagnetic

- Ans. (2)
- Sol. [MnCl₆]³-

 \Rightarrow Mn³⁺ \Rightarrow 3d⁴ \Rightarrow $t_{2\alpha}^{1,1,1}, e_{\alpha}^{10} \Rightarrow$ sp³d² Hybridisation \Rightarrow Paramagnetic

- 6. Difference between bond order of CO and NO+ is $\left(\frac{x}{2}\right)$, then value of 'x' to the nearest integer
- Ans. 0

Sol. Species

Bond order

(i) CO

CO

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

So value of x = 0

7. The value of (ΔH – ΔU) for vaporisation of water at 100°C is 'x' × 10°2 J/mole, assume water vapour to be an ideal gas [Take R = 8.31 J/mole.K]

[report your answer to nearest integer]

Ans. 31.00

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#3

Resonance* | JEE MAIN-2021 | DATE: 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | CHEMISTRY

Sol.
$$H_2O(\ell) \longrightarrow H_2O(g)$$

$$\Delta H^{\circ} = \Delta U^{\circ} + \Delta n_{\circ} RT$$

$$\Delta H^{\circ} - \Delta U^{\circ} = \Delta n_{g}RT$$

Daneity of agrees a solution of NaOLI is 1.2 along then find its molelity

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302R.J2007PLC024029

Ans.

Sol. Let volume of solution is x one

So mass of solution = 1.2 x

& mass of water (solvent) = x gram

So mass of solute = 0.2 x gram

$$Molality = \frac{W_{solute} \times 1000}{GMM_{solute} \times W_{solvert}}$$

$$= \frac{0.2x \times 1000}{40 \times x} = \frac{200}{40} = 5m$$

Ans = 5m

- Identify the wrong statement from following about Ellingham diagram
 - (1) It gives rate of reaction
- (2) It tells about the stability of oxide
- (3) It gives idea about reduction of metal oxide (4) It gives idea about free energy of reduction.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- The main product of electrolysis of conc. H₂SO₄ is 10.

(2) HO3SO - OSO3H

(3) HO₂SO - OSO₂H

(4) O₂

Ans.

Main product of electrolysis of conc. H₂SO₄ is H₂S₂O₈ that is HO₃SO-OSO₃H

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

RESCHARGE | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | CHEMISTRY

Oxidation number of P in H₄P₂O₇, H₄P₂O₅ and H₄P₂O₈ are respectively:

(1) 5, 3, 4

(2) 3, 4, 5

(3) 5, 4, 3

Ans.

Sol. Compound Oxidation number of P

(i) H₄P₂O₇

5

(ii) H₄P₂O₅

(iii) H₄P₂O₆

3

Statement-II: Bohr's model is not in accordance with heisenberg's uncertainty principle (1) Both statements are true (2) Both statement are false (3) Statement-I is true and Statement-II is false (4) Statement-I is false and Statement-II is true Ans. (1) Sol. Theory Based Statement-I: Generally halides of Li are covalent 13. Statement-II: Lithium has high polarising power (1) Both Statement-I & Statement-II are correct (2) Statement-I is correct and Statement-II is incorrect (3) Statement-I is incorrect and Statement-II is correct (4) Both Statement-I and Statement-II are incorrect

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- 14. Unit of rate constant of nth order reaction is :
 - (1) mole 1-n Lit.n-1 Sec-1
- (2) molex Lit.-n Sec-1
- (3) mole-n Lit.n Sec-1
- (4) molen-1 Lit.1-n Sec-1

PAGE#5

Ans. (1)

Sol. Rate = k[A]

$$\frac{\text{mole}}{\text{Lit} \times \text{Sec}} = k \left[\frac{\text{mole}}{\text{Lit}} \right]^{n}$$

⇒ Unit of k = (mole)1-n Lit.n-1 Sec-1

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

© Resonance Eduventures Limited | Toti-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Toll Free : 1800 258 5555 📓 7340010333 🔝 lecetox com Head-arcetds 💟 twitter.com/Head-arcetdd 🛍 www.youtube.com/Head-arcetdd

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

15. Equilibrium constant Kc for dissociation of PCIs according to PCIs(g) PCIs(g) + CIz(g) is 1.844.

Initially 3 mole of PCIs is present in a flask of 1 Lit., then number of moles of PCIs after equilibrium is set Up. Is:

Ans. 1.60

Sol.

PCI₅(g) PCI₃(g) + CI₂(g) K_C = 1.844

Initially

0

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$K_C = \frac{x^2}{3-x} = 1.844$$

$$x^2 = 1.844 (3 - x)$$

$$x^2 = 5.532 - 1.844x$$

$$x^2 + 1.844x - 5.532 = 0$$

$$x = \frac{-1.844 \pm \sqrt{(1.844)^2 - 4(1)(-5.532)}}{2}$$

On solving

$$x = 1.60$$

So at equilibrium number moles of PCI₅ = 1.60

16. According to Freundlich adsorption isotherm $\left(\frac{x}{m}\right) = k(P)^{\frac{1}{n}}$ when pressure increased 2 times, then extent

Sol.
$$\left(\frac{x}{m}\right) = k(P)^{\frac{1}{n}}$$
 (i)

$$64\left(\frac{x}{m}\right) = k(2P)^{\frac{1}{n}}$$
 (ii)

From equation (i) to (ii)

So
$$\frac{1}{n} = 6$$

Decompose Eduventures Ltd

@ Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
tollers continued by butter confidence of the continued by butter continued by butt

This solution was download from Resonance JEE (MAIN) 2021 Solution portal PA

Resonance* | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | CHEMISTRY

17.
$$\begin{array}{c} OH \xrightarrow{H_3PO_4} X \xrightarrow{(BH_3)_2} Y & Product(Y) \text{ is :} \\ CH_3 & OH & CH_3 & OH & CH_3 \\ \end{array}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Ans. (2)

What shall be added so the reaction proceeds in one direction-

- (1) Con.HIO₃
- (2) HOCI
- (3) NH₃
- (4) HNO₂

Ans. (1)

Reaction is reversible due to reducing nature of HI. Oxidising agent like HNO 3, HIO3 should be added to make it irreversible (one direction).

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

- (1) Iodine test
- (2) Seliwanoff test
- (3) Barfoed test
- (4) Tollen's test

Ans. (3

Carbohydrates are polyhydroxy aldehydes and ketones. Carbohydrates may have an aldehyde group (aldoses) or ketose group. Barfoed's test distinguishes monosaccharides from disaccharides. In this test, copper acetate in dilute acid is reduced in 30 seconds by monosaccharides whereas disaccharides take several minutes.

- 20. In DNA complementary base Thymine is-
 - (1) Uracil
- (2) Adenine
- (3) Cytosine
- (4) Guanine

Ans. (2)

Sol. In DNA thymine bind with adenine by to hydrogen bonding.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029 INCOMPRESENTATION OF THE PROPERTY OF THE PROPE Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222 To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
3 7340010333
4 besteek conflectionedds with state conflection and the conflection of th This solution was download from Resonance JEE (MAIN) 2021 Solution portal PAGE#7 RESONANCE | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | CHEMISTRY 21. Statement-1: Aniline is less basic then Acetamide Statement-2: In Aniline, the lone pair delocalised so electron density reduce. (1) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1. © Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029 (4) Statement -1 is False, Statement-2 is False. Ans. 22. Which of the following given statements on Eutrophication are not correct? (1) Eutrophication decrease oxygen level in water. (2) <6ppm oxygen fishes can't survive. (3) Eutrophication involve anaerobic respiration. (4) Eutrophication increase oxygen level in water. Ans. 23. Match the column: Column-I Column-II Furacine Antiseptic (i) (a) (ii) Dimetane Synthetic antihistamine (b) © Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029 (1) (i) - b, (ii) - a, (iii) - c, (iv) - d (2) (i) - a, (ii) - b, (iii) - d, (iv) - c (3) (i) - a, (ii) - d, (iii) - c, (iv) - b (4) (i) - c, (ii) - d, (iii) - a, (iv) - b Ans. 24. What is formula of mustard gas-(2)© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029 Resonance Eduventures Ltd. Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222 To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
tolleck.com/Resonancedu: www.resonancedu: www This solution was download from Resonance JEE (MAIN) 2021 Solution portal PAGE#8

25. Stability of given molecules is:

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

(i) (ii) (iii) (iii) (iv)

(1) i > ii > iv > iii (2) i > ii > iv (3) i > ii > iii > iv (4) iv > iii > i > ii

Ans. (1)

26. Staggered and eclipsed form of ethane are

(1) Enantiomer (2) Anomer (3) Epimer (4) Rotamer

Ans. (4)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac in | CIN - U80302RJ2007PLC024029


Response Respon

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#9





COMPUTER BASED TEST (CBT)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029







ADMISSION OPEN for Session 2021-22

ONLINE + OFFLINE PROGRAMS

CLASS 11, 12 & 12+

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Scholarship Upto 90%*

Toll Free: 1800 258 5555 | Visit us: www.resonance.ac.in

RESONANCE* | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- If $\vec{a} = \vec{i} + \vec{j} + \vec{k}$, $\vec{b} = 2\vec{i} + \vec{j} \vec{k}$ and $\vec{c} = 4\vec{i} + 3\vec{j} + \vec{k}$ then the value of $((\vec{a} + \vec{b}) \times (\vec{a} (\vec{a} \vec{b}) \times \vec{b})) \times \vec{c}$ is:
- (1) $(30\hat{i} 34\hat{j} + 36\hat{k})$ (2) $(30\hat{i} + 34\hat{j} + 36\hat{k})$

(3) $(30\hat{i} + 34\hat{j} - 36\hat{k})$ (4) None of these

Ans. (1)

 $\ddot{a} + \ddot{b} = 3\ddot{i} + 2\ddot{j} = \ddot{r}_1$

 $\vec{r}_2 = \vec{a} - (\vec{a} - \vec{b}) \times \vec{b} = \vec{r}_2 = (\hat{i} + \hat{j} + \hat{k}) - (-\hat{i} + 2\hat{k}) \times (2\hat{i} + \hat{j} - \hat{k})$

 $(\hat{i} + \hat{j} + \hat{k}) - (-\hat{k} - \hat{j} + 4\hat{j} - 2\hat{i}) = (3\hat{i} - 2\hat{j} + 2\hat{k})$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$= (\vec{r}_1 \cdot \vec{c})\vec{r}_2 - (\vec{r}_2 \cdot \vec{c})\vec{r}_1$$

$$= 18[3\hat{i} - 2\hat{j} + 2\hat{k}) - 8[3\hat{i} + 2\hat{j})$$

$$=(30\hat{i}-34\hat{j}+36\hat{k})$$
 Ans.

Which of the following is logically equivalent to $(p \lor q) \land (\neg p \rightarrow q)$ is:

 $(1)(p \vee q)$ (2) (p ∧ q) $(3) \sim p \vee q$ (4) p ∧ ~q Ans. (1) Sol. We know $p \rightarrow q = p \vee q$ © Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029 3. If $S_1 = \{z : |z-3-2i|^2 = 8\}$ and $S_2 = \{z : |z-\overline{z}| = 8\}$ and $S_3 = \{z : re(z) \ge 5\}$ then $S_1 \cap S_2 \cap S_3$ has

(1) Infinite many element

(2) Only one element

(3) No any element

(4) Two element

Ans. (2)

Resonance Eduventures Ltd.


Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029 Tall Free : 1800 258 5555 🔯 7340010333 🚮 backook own/Reparamoteta 💟 witter con/Resonanceta 🚵 www.youtube.com/resonance

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

RESONANCE | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS

Sol. Let z = x+iy



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302R.J2007PLC024029



$$S_2 \Rightarrow |2iy| = 8 \Rightarrow |y| = 4$$

 $y = \pm 4$

$$S_3 \Rightarrow x \geq 5$$

$$S_1 \Rightarrow |z - 3 - 2i|^2 = 8 = (x - 3)^2 + (y - 2)^2 = 8$$

Is a circle with centre (3,2) and radius = $2\sqrt{2}$

· Circle passes through (5,4)

 \Rightarrow There is exactly one point (5,4) in $S_1 \cap S_2 \cap S_3$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

.... αe^β - 224 .

Sol. Put
$$x = 3 \Rightarrow f(3) = 0$$

 $e^{-x} [1^3 + 1^2 + 4(F(t))] = F(x)$
 $e^{-x} (x^3 + x^2 + 4F(x)) = (27 + 9 + 4F(3))$

$$e^{-x}(x^3 + x^2 + 4F(x) - (27 + 9 + 4F(3) = F(x))$$

$$\Rightarrow F(x) = e^{-x} (x^3 + x^2 - 36 + 4F(x))$$

\Rightarrow e^x F(x) = x^3 + x^2 - 36 + 4F(x)

$$F(x) = \frac{x^3 + x^2 - 36}{(e^x - 4)}$$

$$F'(x) = \frac{(3x^2 + 2x)(e^x - 4) - (e^x)(x^3 + x^2 - 36)}{(2x^4 + 4)^2}$$

$$F'(4) = \frac{56(e^4 - 4) - e^4(44)}{(e^4 - 4)^2}$$

$$F'(4) = \frac{12e^4 - 224}{(e^4 - 4)^2}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555
7840010333
**Tooltoik contribution fluoremental with the contribution fl

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

5. The coefficient of x^7 and x^{-7} in the expansion of $\left| x^2 + \frac{1}{bx^2} \right|$ and $\left| x + \frac{1}{bx^2} \right|$ respectively are equal then

the value of b is:

- (1) 1
- (2) 1
- (3) 0
- (4) 2

Ans. (1)

Sol. $\left(x^2 + \frac{1}{bx}\right)$

 $T_{r+1} = {}^{11}C_r (x^2)^{11-r} \left(\frac{1}{bx}\right)^r = {}^{11}C_r (x)^{22-3r} \left(\frac{1}{b}\right)^r$

for $x^4 \Rightarrow 22 - 3r = 7 \Rightarrow r = 5$

 \therefore coefficient of $x^7 = {}^{11}\text{Cs} \left(\frac{1}{b}\right)^5$

similarly $\left(x + \frac{1}{bx^2}\right)^{tt}$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

For $x^{-7} \Rightarrow 11 - 3r = -7 \Rightarrow r = 6$

 \therefore coefficient $x^{-7} = {}^{11}C_6 \left(\frac{1}{h}\right)^8$

 ${}^{11}C_6\left(\frac{1}{b}\right)^6 = {}^{11}C_5\left(\frac{1}{b}\right)^5 \Rightarrow b = 1$

6. Evaluate $\lim_{x\to 2} \frac{x^2f(2) - 4f(x)}{x-2}$, if f(2) = 4 & f'(2) = 1

Ans. (12

Sol. $\lim_{x \to 2} \frac{x^2 f(2) - 4 f(x)}{x - 2} = \lim_{x \to 2} \frac{2 x f(2) - 4 f'(x)}{1} = 2 \cdot (2) \cdot f(2) - 4 f'(2) = 16 - 4 = 12$

7. If $\sin\theta + \cos\theta = \frac{1}{2}$, then the value of $16(\sin 2\theta + \cos 4\theta + \sin 6\theta)$ is :

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol. $(\sin\theta + \cos\theta)^2 = \frac{1}{4}$

 $1 + \sin 2\theta = \frac{1}{4}$

 $\sin 2\theta = \frac{-3}{4}$

 $16(\sin 2\theta + \cos 4\theta + \sin 6\theta) = 16(\sin 2\theta + 1 - 2\sin^2 2\theta + 3\sin 2\theta - 4\sin^3 2\theta)$

 $= 16 (4\sin 2\theta + 1 - 2\sin^2 2\theta - 4\sin^3 2\theta) = -23$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

| JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS

If mean and variance of eight observation 10, 13, 6, 7, a, 12, b, 12 are 9 and $\frac{37}{4}$ respectively, then the 8. value of (a-b)2 is s

(1)25

(2)36

(3) 16

(4)49

Ans.

9 = 10 + 13 + 6 + 7 + a + 12 + b + 12Sol.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\sum x_i^2 = 100 + 169 + 36 + 49 + a^2 + 144 + b^2 + 144$$

$$= a^2 + b^2 + 642$$

$$\sigma^2 = \frac{\sum x_i^2}{8} - (\bar{x})^2 = \frac{37}{4}$$

$$\frac{a^2 + b^2 + 642}{8} - 81 = \frac{37}{4}$$

$$\Rightarrow a^2 + b^2 = 80 \Rightarrow a^2 + (12-a)^2 = 80 \Rightarrow a^2 - 12a + 32 = 0$$

 \Rightarrow a = 4 or a = 8 b=8 or b = 4 \Rightarrow $(a - b)^2 = 16$ Ans.

If matrix $A = \begin{bmatrix} 2 & 4 \\ -1 & 1 \end{bmatrix}$ and $A^{-1} = \alpha I + \beta A$, α , $\beta \in R$. Then the value of $(\alpha - 6\beta)$ is equal to :

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Ans. (1)

Sol. Equation of $|A-\lambda I| = 0$

$$\begin{vmatrix} 2-\lambda & 4 \\ -1 & 1-\lambda \end{vmatrix} = 0$$

$$\Rightarrow$$
 $(2-\lambda)(1-\lambda)+4=0 \Rightarrow \lambda^2-3\lambda+2+4=0 \Rightarrow \lambda^2-3\lambda+6=0$

$$A^2 - 3A + 6I = 0$$

Multiply both side by A-1

$$A - 3I + 6A^{-1} = 0$$

$$A^{-1} = \frac{-A}{6} + \frac{I}{2}$$
 (1)

$$A^{-1} = \alpha 1 + \beta A$$
(2)

Comparing equation (1) and (2)

$$\alpha = \frac{1}{2}$$
, $\beta = \frac{-1}{6}$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555
7340010333
to extend confliction confliction contact with the contact with the confliction con

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#4

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

If α and β are the roots of the equation $x^2-20^{1/4}x+5^{1/2}=0$, then the value of $\alpha^8+\beta^8$ is equal to :

(1) 25 (2) 50

(3) 75

(4) 100

Ans.

Sol. $x^2 - 20^{1/4}x + 5^{1/2} = 0$

$$(x^2 + \sqrt{5})^2 = \sqrt{20}x^2$$

 $x^4 + 5 + 2\sqrt{5}x^2 = 2\sqrt{5}x^2$

- 11. Let n be a two digit natural number. If n is selected at random then find the probability such that 21 - 2 is a multiple of 3.
 - $(1) \frac{1}{4}$
- (3) $\frac{1}{2}$

Ans. (3)

Sol. Total number of cases = 90C1 = 90

Now $2^n - 2 = (3 - 1)^n - 2$

=
$${}^{n}C_{0} 3^{n} - {}^{n}C_{1} 3^{n-1} + \dots + (-1)^{n-1} {}^{n}C_{n-1} 3 + (-1)^{n} {}^{n}C_{n} - 2$$

 $= 3/3^{n-1} - n 3^{n-2} + (-1)^{n-1} + (-1)^{n-2}$

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

So number of favourable cases = 45, Hence required probability = $\frac{45}{90} = \frac{1}{2}$

- 12. If tangents to the circle $(x 1)^2 + (y 3)^2 = 2^2$ are drawn from the point (-1, 1) such that the points of contact of tangents are A and B. Also a point P lies on the circle such that AB = AP, then the area of ∆ABP is:

- (1) 2 (2) √3 (3) 4 (4) 2√2

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#5


| JEE MAIN-2021 | DATE: 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS

Ans. (3)

Sol.

Q

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029




$$AQ = \sqrt{S_1} = \sqrt{(-1-1)^2 + (1-3)^2 - 4} = 2$$

AD = 2sin459

AB = 4sin45º

AP = 4sin45°



Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#6

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

MEMORY BASED | MATHEMATICS

- 13. If log_32 , $log_3(2^x-5)$, $log_3(2^x-\frac{7}{2})$ are in A.P. then the value of x is
 - (1) 4
- (2)2
- (3) 3
- (4) 5

Ans. (3

Sol.
$$2\log_3(2^x - 5) = \log_3 2 + \log_3(2^x - \frac{7}{2})$$

$$\Rightarrow \log_3(2^x - 5)^2 = \log_3(2(2^x - \frac{7}{2}))$$

$$\Rightarrow$$
 $(2^{\times})^2 - 10(2^{\times}) + 25 = 2(2^{\times}) - 7$

$$\Rightarrow$$
 $(2^{x})^{2} - 12(2^{x}) + 32 = 0$

$$\Rightarrow$$
 2× = 4 or 8

$$\Rightarrow$$
 x = 2 or

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

⇒ X = 3

14. If
$$\sec y \frac{dy}{dx} = \sin(x+y) + \sin(x-y)$$
; $y(0) = 0$ then the value of $5y'\left(\frac{\pi}{2}\right)$ is

- (1) 3
- (2)
- (3) 2
- (4) 1

Ans. (3)

Sol.
$$\sec y \frac{dy}{dx} = 2\sin(x) \cos(y) \dots (1)$$

$$\int \sec^2 y \, dy = \int 2\sin x \, dx$$

$$tany = -2cosx + c$$

When
$$x = 0$$
, $y = 0 \Rightarrow c = 2$

$$tany = -2cosx + 2$$
(2)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$(1 + \tan^2 y) \frac{dy}{dx} = 2\sin x$$
(3)

By equation (2) & (3)

$$\frac{dy}{dx} = \frac{2\sin x}{1 + (2 - 2\cos x)^2}$$

$$f'\left(\frac{\pi}{2}\right) = \frac{2}{1+(2)^2} = \frac{2}{5} \Rightarrow 5f'\left(\frac{\pi}{2}\right) = 2$$

(1) (-1, 5) & (5, 1)

(2) (-2, 5) & (-5, 1)

(3) (3, 6) & (5, 1)

(4) (-1, 1) & (5, -1)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#7

RESONANCE | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS

Ans.

Sol. $(x-2)^2 + (y-3)^3 = 13$



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$y = \frac{3}{2}x$$

line perpendicular to the above line and passing through (2,3) is 3y +2x = 13

Coordinates of P,Q \Rightarrow $(2 \pm \sqrt{13}\cos\theta, 3 \pm \sqrt{13}\sin\theta)$

$$\Rightarrow \left(2\pm\sqrt{13}\!\left(\!\frac{-3}{\sqrt{13}}\right)\!\!3\pm\!\sqrt{13}\!\left(\!\frac{2}{\sqrt{13}}\right)\!\right)$$

Evaluate $\lim_{n\to\infty} \frac{1}{n} \sum_{j=1}^{n} \frac{((2j-1)+8n)}{((2j-1)+4n)}$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$\lim_{n\to\infty} \frac{1}{n} \sum_{j=1}^{n} \frac{2^{j} - \frac{1}{n} + 8}{2^{j} - \frac{1}{n} + 4} \frac{1}{n}$$

$$=\int\limits_0^1\frac{2x+8}{2x+4}dx$$

$$= \int_0^1 dx + \int \frac{4}{2x+4} dx$$

$$= 1 + 4.\frac{1}{2} [\ln(2x + 4)]_0^{1}$$

= 1 + 2[ln6 - ln4]

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
tolteck.com/Reconancedu: velocities with youtube.com/resonance.ac.in | Smith | Sm

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

Resonance | Jee Main-2021 | Date : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS

17. If
$$f(x) = \begin{cases} (1+|\sin x|)^{\frac{a}{|\sin x|}} & -\frac{\pi}{4} < x < 0 \\ b & x = 0 \\ \frac{\cos(4x)}{e^{\cos(2x)}} & 0 < x < \frac{\pi}{4} \end{cases}$$
 is continuous, then find the value of $(a + \ell nb)$.

Ans.

Sol. RHL =
$$f(0+) = \lim_{x \to 0^+} e^{\frac{\cot 4x}{\cot 2x}} = \lim_{x \to 0^+} e^{\frac{\tan 2x}{\tan 4x}} = e^{\frac{1}{2}}$$

LHL =
$$f(0^-) = \lim_{x \to 0^-} (1 - \sin x)^{\frac{a}{-\sin x}}$$

= $e^{x \cdot o^- \sin x} = e^a$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)7/4/27777777, 27777700 | contact@resonance.ac.in | CIN - U80302R.J2007PLC024029

$$\Rightarrow \frac{1}{e^2} = e^a = b \Rightarrow a = \frac{1}{2} \& b = e^{\frac{1}{2}} \Rightarrow \ell nb = \frac{1}{2}$$
$$\Rightarrow a + \ell nb = 1$$

Evaluate $I = \int_{-\pi/4}^{\pi/4} \frac{dx}{(1 + e^{x\cos x})(\sin^4 x + \cos^4 x)}$

 $(4) \frac{\pi}{4}$

Ans. (2)

f(a+b-x)=f(x)

$$I = \int_{1/4}^{1/4} \frac{e^{x \cos x}}{4 - x \cos x} dx$$

© Resonance Eduventures Limited | Toli-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

 $2I = \int_{-\pi/4} \frac{1}{\sin^4 x + \cos^4 x}$

$$2I = \int_{-\pi/4}^{\pi/4} \frac{1}{1 - \frac{1}{2} \sin^2 2x} dx$$

$$2I = \int_{-\pi/4}^{\pi/4} \frac{2 \sec^2 2x}{\sec^2 2x + 1} dx$$
 (put tan2x = t

$$2I = \frac{4}{2} \int_{0}^{\infty} \frac{dt}{2 + t^{2}} = \left(\frac{1}{\sqrt{2}} \tan^{-1} \frac{1}{\sqrt{2}}\right)_{0}^{\infty}$$

$$I = \frac{\pi}{2\sqrt{2}}$$

Resonance Eduventures Ltd.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Toll Free: 1800 258 5555 🔯 7340010333 🔣 Societa Com/Resonance Edu 💟 rwitter.com/Resonance Edu 🖺 www.youtube.com/

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#9

Resonance* | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS

19. If the domain of the function $f(x) = \log_8(\log_4(\log_3(16x-x^2-77)))$ is (a,b), then find the value of

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Sol. $\log_4 \log_3 (18x - x^2 - 77) > 0$

$$18x - x^2 - 77 > 3$$

$$x^2 - 18x + 80 < 0$$

$$(x-8)(x-10) < 0$$

$$x \in (8, 10)$$


$$\therefore a = 8 \text{ and } b = 10$$

$$\therefore 1 = \int_{a}^{b} \frac{\sin^{3} x \, dx}{\sin^{3} x + \sin^{3}(a + b - x)} = \int_{a}^{10} \frac{\sin^{3} x \, dx}{\sin^{3} x + \sin^{3}(a + b - x)}$$
Using the property
$$\int_{a}^{b} f(x) dx = \int_{a}^{b} f(a + b - x) dx, \text{ we get}$$

Find area bounded by y = max $\{0, \ell nx\}$ and y < 2^x where $\frac{1}{2} < x < 1$:

(1) $\frac{2+\sqrt{2}}{\ell n2}$ (2) $\frac{2-\sqrt{2}}{\ell n2}$ (3) $\frac{1-\sqrt{2}}{\ell n2}$ (4) $\frac{1+\sqrt{2}}{\ell n2}$

Ans. (2) Sol.



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029 Toll Free : 1800 258 5555 🔞 7340010333 🔣 tooleoxic own ResonanceEds 🗾 with tercon/ResonanceEds 🔠 www.ycutube.com/resovatch 🕒 biog resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

| JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS

$$\sin^2 x -2 + \cos^2 x \cos 2x$$

If $f(x) = 2 + \sin^2 x$ $\cos^2 x$ $\cos^2 x$ $\cos^2 x$, $x \in (0, \pi]$. Then the maximum value of f(x) is ? 21. $\cos^2 x$ 1+cos 2x

Ans. 06.00

Sol. $R_1 \rightarrow R_1 - R_2$

R₂→ R₂ - R₃

-2 -2

₌ 2 0 -1 sin2 x cos2 x 1+cos2x

 $=-2(\cos^2 x) + 2(2+2\cos 2x + \sin^2 x)$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

= -2(COS-X-SIN-X) +4 COS2X + 4

 $=4+2\cos 2x=2(2+\cos 2x)$, maximum value of f(x)=6