

2021

COMPUTER BASED TEST (CBT) Memory Based Questions & Solutions

Date: 25 July, 2021 (SHIFT-2) | TIME: (3.00 p.m. to 6.00 p.m)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

SUBJECT: PHYSICS Resonance Eduventures Ltd. Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222 To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029 Toll Free: 1800 258 5555 🔘 7340010333 📝 treetonic confessionate del 💆 twitter com Resonance del 🖺 www.youtube.com/resonance del This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

PART: PHYSICS

Value of force $F = A\sin(Bt) + C\cos(Dx)$ find dimension of $\frac{AB}{D}$

(2) ML²T⁻³ (3) ML¹T⁻³

Dimension of $A = MLT^{-2}$, $B = T^{-1}$, $D = L^{-1}$

 $Dim = \frac{AB}{D} = \frac{MLT^{-2}T^{-1}}{L^{-1}} = ML^{2}T^{-3}$

- Force is given by F = (5y + 20)j Find work done for moving particle from y = 0 to y = 5:
 - (1) 162.5 J
- (2) 165 J
- (3) 132.5 J
- (4) 140.5 J

Ans. (1)

 $w = \int F.dy$ Sol.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029



- A hot air balloon is ascending with constant velocity of 10 m/s, when balloon reaches a height of 75 m, a stone is dropped from balloon, what will be the height of balloon, when stone reaches earth?
- (2) 135 m.
- (3) 140 m.

Ans. (1)


For stone Sol.

$$75 = -10t + \frac{1}{2}gt^2$$

$$75 = -10t + 5t^2$$

$$t^2 - 2t - 15 = 0$$

 $t = 5$ sec.



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$H = vt + 75$$

$$H = 10 \times 5 + 75 = 125 \text{ m}.$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555 S 7340010333 enter with contact of the contribution of the contact of the contribution of the contact of the contribution of the cont

This solution was download from Resonance JEE (MAIN) 2021 Solution portal PAGE#1

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Resonance* | Jee Main-2021 | Date : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

- Relation between position and time of a particle moving along straight line is given by $t = x + 3x^2$. Find acceleration of particle at t = 10s
 - $(1) \frac{-5}{1331}$
- $(2) \frac{6}{1331}$

Ans.

 $t = x + 3x^2$...(1) Sol.

$$1 = \frac{dx}{dt} + 6x \frac{dx}{dt} \implies v = \frac{1}{(1+6x)}$$

$$0 = \frac{d^2x}{dt^2} + 6\left(x\frac{d^2x}{dt^2} + \left(\frac{dx}{dt}\right)^2\right)$$

$$0 = a + 6xa + 6v^2$$

$$a = \frac{-6v^2}{(1+6x)}$$
 ...(2)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

From equation(1)

$$10 = x + 3x^2$$

$$3x^2 + x - 10 = 0$$

$$3x^2 + 6x - 5x - 10 = 0$$

$$3x(x+2) - 5(x+2)$$

$$(3x-5)(x+2) \Rightarrow x = \frac{5}{3}$$

$$a = \frac{-6}{\left(1 + 6 \times \frac{5}{3}\right)^3} = \frac{-6}{1331}$$

- Two particle of same mass & charges Q₁ and Q₂ are moving perpendicular to an uniform magnetic field where the ratio of charges is $\frac{Q_1}{Q_2} = \frac{1}{2}$ and ratio of velocities is $\frac{V_1}{V_2} = \frac{3}{2}$ then find the ratio of the radius $\frac{R_1}{R_2}$:

$$\frac{Q_1}{Q_2} = \frac{1}{2} \& \frac{V_1}{V_2} = \frac{3}{2}$$

$$R = \frac{mv}{aB}$$

$$\frac{R_1}{R_2} = \frac{V_1}{V_2} \times \frac{Q_2}{Q_1} = \frac{3}{2} \times \frac{2}{1} = \frac{3}{1}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029 Tall Free : 1800 258 5555 🔯 7340010333 🚮 bootook own/Reparamotedu 💟 witter com/Resonancedu. 🚵 www.youtube.com/resonance

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

@ Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

RESONANCE | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

- A particle performing SHM with amplitude A. Find the ratio of kinetic energy and total energy when particle
 - $(1)\frac{3}{4}$

Ans.

 $V_{A/2} = \omega \sqrt{A^2 - x^2}$

$$= \omega \sqrt{A^2 - \left(\frac{A}{2}\right)^2} = \omega \left(\frac{\sqrt{3}}{2}A\right)$$

$$=\frac{\sqrt{3}}{2}V_{\text{max}}$$

$$KE = \frac{1}{2} m \left(\frac{\sqrt{3}}{2} V_{max} \right)^2$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\frac{KE}{TE} = \frac{3}{4}$$

Ans.

- In photoelectric effect stopping potential is $3V_0$ for incident wave length λ_0 and stopping potential V_0 for incident wavelength 2λο. Find threshold wavelength.
- (1) 3λο
- (2) 2λο
- (3) 420
- (4) 8λο

Ans. (3)

Sol. KE = hv - W

$$eV = \frac{hc}{\lambda} - W$$

For first case

$$e(3V_0) = \frac{hc}{\lambda_0} - W \qquad \dots (i)$$

For second case

$$eV_0 = \frac{hc}{2\lambda_0} - W \qquad \dots (i$$

From equation (i) and (ii)

For
$$\lambda_1$$

or
$$\lambda_{th}$$

$$W = \frac{hc}{\lambda_{th}}$$

Resonance Eduventures Ltd.

4λ₀ λ_{th} Pesonence Pesononice

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555
7340010333
acceptable contracts of the contract of t

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#3

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance | Jee Main-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

- 8. Efficiency of heat engine is $\eta = 1/6$. If temperature of sink is decreased by 62K, then efficiency becomes 1/3. Find temperature of source :
 - (1) 372K
- (2) 272K
- (3) 350K
- (4) 450K

Ans. (1)

Sol. $\eta = \left(1 - \frac{T_2}{T_1}\right)$

 $\frac{T_2}{T_1} = 1 - \eta = 1 - \frac{1}{6}$

....(1)

 $\frac{T_2 - 62}{T_2 - 62} = 1 - \frac{1}{2}$ (2)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Equation (2)

 $\Rightarrow \frac{T_2}{T_2 - 62} = \frac{5}{6} \times \frac{3}{2} = \frac{5}{4}$

 \Rightarrow $T_2 = 5 \times 62$

From eq. (1)

 $T_1 = \frac{T_2}{1 - \eta} = \frac{5 \times 62}{1 - \frac{1}{6}} = 5 \times 62 \times \frac{6}{5} = 372K$

- 9. Activity of an element x becomes 1/8 of initial in 30 years. Find half-life:
 - (1) 10 Year.
- (2) 12 Year
- (3) 15 Year
- (4) 17 Year

Ans. (1)

Sol. $A = A_0e^{-\lambda t}$

For half life

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

 $\frac{1}{2} = e^{\lambda t_{1/2}}$

....(1)

Given 1/8 = e-230

....(2)

Solving (1) and (2)

 $e^{-3\lambda t_{1/2}} = e^{-\lambda 30}$

 $T_{1/2} = 10 \text{ Yrs.}$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-45 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U803028J2007PLC024029

Tall Free: 1800 258 5555
7340010333
to toolesk own Resonancedds with the same way you take contributions at an incidence of the same way you take contributions at an incidence of the same way.

IDII FERR : 1900 539 3333 🔯 13400 10333 🎹 antidoparating and antidoparating

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

RESONANCE | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS


- If De-Broglie wavelengths of photon and electron are equal, what will be the ratio of kinetic energy of electron and energy of photon? Given that velocity of electron is v and velocity of light is c :

Ans.


- De-Broglie wavelength is given by $\lambda = \frac{h}{}$ Sol.
 - $KE_{pn} = mc^2 = pc$
- $KE_0 = \frac{1}{m} mv^2 = \frac{pv}{m}$...(2)
- © Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN U80302RJ2007PLC024029

- 11. A square loop of total resistance 16 Ω . If a battery of 2V and 1 Ω internal resistance is connected across one of its side then find potential difference across its diagonal :
 - (1) 1V
- (2) 2V
- (3) 3V
- (4) 4V

Ans. (1) Sol.



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



 $\frac{r_2}{r_1}i = \frac{r_2}{3+1}$

 $V_{AB} = \frac{1}{8} \times 8 = 1V$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

RESONANCE | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

- 12. \vec{A} and \vec{B} are two vectors such that $|\vec{A}| = 2$ and $|\vec{B}| = 5$. If $|\vec{A} \times \vec{B}| = 8$, then $|\vec{A}.\vec{B}| = 7$

Ans.

 $\vec{A} \times \vec{B} = \vec{A} \vec{B} \sin \theta$ Sol.

 $\Rightarrow 10\sin\theta = 8$

Now
$$|\vec{A}.\vec{B}| = |\vec{A}| |\vec{B}| \cos\theta = 10 \times \frac{3}{5} = 6$$

Find significant figure for the value 0.00346.

(2)4

(3) 3

(4)2

Ans.

Sol. There are 3 non zero digit after the decimal point so significant number is 3. 0.00346

14 For a prism, if angle of minimum deviation is equal to angle of prism. If refractive index of prism material is μ. Then angle of prism should be?

(1)
$$2\sin^{3}\left(\frac{\mu}{2}\right)$$
 (2) $2\cos^{3}\left(\frac{\mu}{2}\right)$ (3) $3\cos^{3}\left(\frac{\mu}{2}\right)$ (4) $3\sin^{3}\left(\frac{\mu}{2}\right)$

Ans.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$\mu = \frac{A}{\sin(\frac{A}{2})}$$

$$\mu = \frac{\sin A}{\sin A/2}$$

$$\mu = 2\cos\frac{A}{2}$$

$$A = 2\cos^{-1}\left(\frac{\mu}{2}\right)$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#6

Resonance | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

A photon of wavelength 500 nm falls on a metal surface of work function 1.3eV. An electron releases from metal moved in a perpendicular magnetic field. In a circular path of radius 30 cm. Then the magnitude of magnetic field will be ?

(1) 12.2 µT

(2) 10.2 μT (3) 8.2 μT (4) 6.2 μT

Ans. (1)


$$\frac{1240}{500} = 1.3 + KE_{max}$$

Now R =
$$\frac{mv}{qB} = \frac{\sqrt{2mKE}}{qB}$$

$$B = \frac{\sqrt{2m KE}}{aB}$$

$$B = \frac{\sqrt{2 \times 9.1 \times 10^{-31} \times 1.18 \times 1.6 \times 10^{-19}}}{1.6 \times 10^{-19} \times 30 \times 10^{-2}}$$

$$B = 12.2 \times 10^{-6}$$



Decemence Educentures Ltd

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555 ◎ 7340010333 달 beekeek own/fleorencedds ☑ witter con/fleorencedds ☑ www.youtube.com/reconstain ⓒ biog-resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

Resonance* | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

Sol.



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

7E



Electric field due to P1 at axis point S

$$E_{abdis} = \frac{2KP_1}{r^3}$$

$$\Rightarrow Ecos37^\circ = \frac{2KP_1}{r^3} \qquad ...(1)$$

Electric field due to \vec{P}_2 at perpendicular bisector at point S.


$$E_{\perp} = \frac{KP_{2}}{r^{3}}$$

$$\Rightarrow \qquad E \sin 37^{\circ} = \frac{KP_{2}}{r^{3}} \dots (2)$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow \frac{2P_1}{P_2} = \frac{4}{3} \Rightarrow \frac{P_1}{P_2} = \frac{2}{3}$$

17. Power in both the given circuit are same then find angular frequency of AC source.



220V (1)200(2)300(4)500© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029 nesonance Eudventures Etd. Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222 This solution was download from Resonance JEE (MAIN) 2021 Solution portal PAGE#8 RESONANCE | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS © Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029 $10 = \sqrt{\left(\omega L - \frac{1}{\omega C}\right)^2 + R^2}$ $100 = \left[\omega(0.1) - \frac{1}{\omega(40 \times 10^{-6})}\right]^2 + 100$ $\omega^2 = \frac{1}{4} \times 10^6$ For the given circuit, find the potential drop across 2Ω resistance? 20V @ Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029 The wire AB is of length 10 cm, and its resistance is 1Ω /cm. Point D is mid-point of wire AB. (1) 2.44 V (2) 4.44 V (3) 3.44 V (4) 10.44V Ans. (2) Sol. 20V 20V <u>10</u> Ω ww 20 10 10 + 5 Von - A AA V © Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 55555 | 7340010333 | sociotok om/Reconancedd: www.resonance.ac.in | Cin: U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 9

Mass of a planet is double the mass of earth. Both the planet have same mass density. A body has weight W on surface of earth, then weight of the same body on surface of planet?

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol. 2ME = MP

$$2\rho \times \frac{4}{3}R_E^3 = \rho \times \frac{4}{3}\pi R_P^3$$
 (same density)

 $g_P = \frac{GM_P}{R_P^2}$ (acceleration due to gravity)

$$g_P = \frac{G2M_E}{(2^{1/3}R_E)^2} = \frac{G2M_E}{2^{2/3}R_E^2}$$

 $g_P = 2^{1/3} g_e$

weight on planet = 21/3 weight on earth

Wp = 21/3 W

A force $\vec{F} = 40i + 10j$ is applied on a stationary object of mass 5kg. What will be the position of object 20. after 10e if initially chiect was at origin ?

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302R.J2007PLC024029

Ans.


Sol. $\vec{a} = 8i + 2j$

$$\vec{s} = \vec{u}t + \frac{1}{2}\vec{a}t^2$$

$$\vec{s} = \frac{1}{2} (8\hat{i} + 2\hat{j}) \times 100$$

$$\bar{s} = 400\hat{i} + 100\hat{j}$$

An AC Source with $V_{max} = 200 \text{ V}$ and f = 50 Hz connected across 10Ω resistance. Find the time in which source voltage changes from maximum to rms value.



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- (1) $\frac{1}{200}$ s

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555 | 7340010333 | sectors confidence of the confidence of the

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#10

Resonance | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

$$\omega = 2\pi f$$

= $100 \pi \text{ rad/s}$.

$$V = V_0 \sin(\omega t + \frac{\pi}{2})$$

© Resonance Eduventures Limited | Toll-Free 1800-250-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\omega t = \frac{\pi}{4}$$

thus
$$t = \frac{\pi/4}{100\pi} = \frac{1}{400}$$
 s

- A Disc of mass 2 kg and radius 2m is rotating with angular velocity ω = 600 rpm. If this disc stops under the action of a constant Torque in 10 sec then if Torque is $n\pi$ then 'n' is.
- (2) 6

Ans. (3)

Sol.
$$\omega = \frac{600 \times 2\pi}{60} = 20\pi \text{ rad/s}$$

 $\omega_t = \omega_t + \alpha_t$

$$0 = 20 \pi - \alpha(10)$$

$$\alpha = 2\pi \text{ rad/s}^2$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$= \frac{mR^2}{2} \times 2\pi = \frac{2 \times 4}{2} \times 2\pi = 8\pi$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302R12007PLC024029 Toll Free : 1800 258 5555 🙆 7340010333 🌃 bookook own/Reconsciedus 🗾 twitter con/Reconsciedus 🔠 www.ycutube.com/recovatch 🔼 biog reconscie ac i

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302R.J2007PLC024029

For two vector X and Y, |X| = |Y| and |X - Y| = n |X + Y|. Then find angle between X and Y

(1)
$$\cos^{-1}\frac{1-n^2}{1-n^2}$$

(1)
$$\cos^{-1}\frac{1-n^2}{1+n^2}$$
 (2) $\cos^{-1}\frac{1+n^2}{1-n^2}$ (3) $\cos^{-1}\frac{2-n^2}{2+n^2}$ (4) $\cos^{-1}\frac{2+n^2}{2-n^2}$

(3)
$$\cos^{-1} \frac{2-n^2}{2+n^2}$$

(4)
$$\cos^{-1}\frac{2+n^2}{n}$$

Ans. (1)

Sol.
$$|\vec{X} - \vec{Y}| = n|\vec{X} + \vec{Y}|$$

$$\left|\vec{X}\right|^2 + \left|\vec{Y}\right|^2 - 2\left|\vec{X}\right| \vec{Y} \cos\theta = n^2 \left|\vec{X}\right|^2 + \left|\vec{Y}\right|^2 + 2\left|\vec{X}\right| \vec{Y} \cos\theta$$

As
$$|\vec{X}| = |\vec{Y}|$$

$$2|\vec{X}|^2 - 2|\vec{X}|^2\cos\theta = 2n^2|\vec{X}|^2 + 2n^2|\vec{X}|^2\cos\theta$$

$$1 - \cos\theta = n^2 + n^2 \cos\theta$$

$$\cos\theta = \frac{1 - n^2}{1 + n^2}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Find energy required to break an Aluminium nucleus into its constituent nucleons.

- $(m_n = 1.00867 \text{ u}, m_p = 1.00783 \text{ u}, m_{Al} = 26.98154 \text{ u})$ (1) 225 MeV
 - (2) 230 MeV
- (3) 235 MeV
- (4) 245 MeV

Ans.

Binding Energy = ΔmC^2

$$\Delta m = [13 \times 1.00783 + 14 \times 1.00867 - 26.98154]$$

A Cell of Voltage 'Vo' is connected across a capacitor of capacitance 'C'. Now the space between the plates is filled with a material of dielectric constant K. Find the ratio of charge appear on the plates of

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222


To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302R12007PLC024029

Toll Free: 1800 258 5555 | 7340010333 | sectes contilesconanced: www.resonanced. www.resonanced

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

RESONANCE | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS



$$Q_1 = CV$$

$$Q_2 = KCV$$

$$\frac{Q_1}{Q_2} = \frac{1}{K}$$

Pure S_i at room temperature has equal electron (n_0) and hole (n_0) concentration of $1.5 \times 10^{16} \, \text{m}^{-3}$. Doping 26. by indium increases n_h to 3 × 10²² m⁻³. Calculate n_e in the doped S_i.

(1)
$$7.5 \times 10^9 \,\mathrm{m}^{-3}$$

$$(4) 7.5 \times 10^7 \,\mathrm{m}^{-3}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow \qquad n_e = \frac{n_L^2}{n_h} = \frac{(1.5 \times 10^{16})^2}{3 \times 10^{22}} = 7.5 \times 10^9 \text{ m}^{-3}$$

A particle starts from rest and moves with a variable acceleration $a = \alpha t + \beta t^2$, where α and β are positive 27. constants. Find the distance covered by particle in t = 1sec to t = 2sec ?

$$(1) \frac{11}{6} \alpha + \frac{15}{12} \beta$$

$$(2) \frac{7}{6} \alpha + \frac{17}{12}$$

(2)
$$\frac{7}{6}\alpha + \frac{17}{12}\beta$$
 (3) $\frac{7}{6}\alpha + \frac{15}{12}\beta$

(4)
$$\frac{1}{3}\alpha + \frac{15}{12}\beta$$

Ans.

Sol.
$$\int_{0}^{v} dv = \int_{0}^{1} adt$$

$$V = \frac{\alpha t^2}{2} + \frac{\beta t^2}{3}$$

$$S = \left[\frac{\alpha t^3}{6} + \frac{\beta t^4}{12}\right]^2 \Rightarrow S = \frac{7}{6}\alpha + \frac{15}{12}\beta$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 14

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

RESONANCE | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

28. A carrier frequency of 1 MHz and peak value of 10 v is amplitude modulated with a signal frequency of 10 KHz with peak value of 0.5 v. Find modulation index.

(1) 0.02 (2) 0.03

(4) 0.05

Ans. (4)

Sol. $A_{\text{max.}} = 10 + 0.5 = 10.5$

 $A_{min} = 10 - 0.5 = 9.5$

L - 1-1-


$$m_a = \frac{A_{max.} - A_{min.}}{A_{max.} + A_{min.}} = \frac{10.5 - 9.5}{10.5 + 9.5} = 0.05$$

Two soap bubbles of radius r₁ and r₂ in vacuum are combined isothermally to form a new bubble. Find the radius of this new bubble?

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Ans. (1)

Sol.





By surface energy conservation

 $\sigma A_1 + \sigma A_2 = \sigma A$

 $\sigma[2 \times 4\pi r_1^2] + \sigma[2 \times 4\pi r_2^2] = \sigma[2 \times 4\pi r_2^2]$

 $r_1^2 + r_2^2 = r^2$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.


Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029 Toll Free: 1800 258 5555 🚳 7340010333 📝 frontesk own/Resonancefdu 💟 twitter com/Resonancefdu 🚵 www.youtube.com/resonance

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

RESCHARCE | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

- 30. A ray is incident on a slab of refractive index $\frac{5}{4}$ at an angle θ as shown in figure. Find maximum angle
 - 9. So that TIR occur at surface AD.



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

./5

Ans. (3)

Sol.
$$1 \times \sin\theta = \frac{5}{4} \sin (90 - C)$$

$$\sin\theta = \frac{5}{4}\cos C$$

./E

but sinC =
$$\frac{1}{11} = \frac{4}{5}$$

$$\cos C = \frac{3}{5}$$

$$\sin\theta = \frac{5}{4} \times \frac{3}{5} = \frac{3}{4}$$

For T.I.R.
$$\sin \theta < \frac{3}{4}$$

$$\theta = \sin^{-1} \frac{3}{4}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029


Resonance Eduventures Ltd.


Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 15









COMPUTER BASED TEST (CBT)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Date: 25 July, 2021 (SHIFT-2) | TIME: (3.00 p.m. to 6.00 p.m) Duration: 3 Hours | Max. Marks: 300

SUBJECT: CHEMISTRY

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RU2007PLC024029

Toll Free : 1800 258 5555 📵 7340010333 👔 techtosk ours Resonanceldu 💟 twitter com/Resonanceldu 🔁 www.you.fuide.com/resonanceldu 🔁 biog. resaminou ac. i

This solution was download from Resonance JEE (MAIN) 2021 Solution portal







ADMISSION OPEN for Session 2021-22

ONLINE + OFFLINE PROGRAMS

CLASS 11, 12 & 12+

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Scholarship Upto 90%*

Toll Free: 1800 258 5555 | Visit us: www.resonance.ac.in

RESONANCE" | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | CHEMISTRY

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

3 gm of 'X' dissolve in 100 gm of CCI4 which increases the boiling point by 0.6. Find molar mass of 'X'. Given Kb of CCl4 = 5 K kg/mol.

Ans. 250

Sol. $\Delta T_b = K_b m$

= 5 ×
$$\left(\frac{\text{Vt.} \times 1000}{\text{M.M} \times \text{Mass of solvent(g)}}\right)$$

 $= 5 \times \frac{3 \times 1000}{\text{M.M} \times 100}$

 $0.6 = \frac{150}{M.M}$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

In the following ions, The spin only magnetic moment of Ti3+; Sc3+; V2+ respectively are

(1) 1.73, 0, 3.87 (2) 1.73, 3.87, 0 (3) 3.87, 0, 1.73 (4) 0, 1.73, 3.87

Ans. (1)

Ti3+ (Unpaired electron = 1)

Sc3+ {Unpaired electron = 0}

V*4 (Unpaired electron = 3)

 $\mu = \sqrt{n(n+2)} B.M.$

Heat given to a system is 150 joules and work done by the system is 200 joules. The magnitude of the change in the internal energy is

Ans. (50)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

 $\Delta E = q + w$

(First law of thermodynamics

 $\Delta E = 150 + (-200)$

 $\Delta E = -50$ Joule

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

© Resonance Eduventures Limited | Toll-Frée 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

RESONANCE" | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | CHEMISTRY

- The covalent radii of FT = 1.33 Å, O2T = 1.40 Å and for N = 0.74 Å. Then which of the following in correct.
 - (1) Ionic radius of N3-is in between F- and O2- but greater than 'N'.
 - (2) Ionic radius of N3-is greater than both F- and O2-and greater than 'N'.
 - (3) Ionic radius of N3-is less than both F- and O2-and less than 'N'.
 - (4) Innic radius of N3-is less than both F1 and O2-but less than 'N'

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

$$N = 0.74 \text{ Å}$$

Ionic radius of N3-is greater than F- and O2-

size of Anion

Magnitude of -ve change

For the following ions correct order of their Bond order is:

$$(1) O_2^- > O_2^- > O_2^+ > O_2^{2-}$$

$$(2) O_2^+ > O_2 > O_2^- > O_2^{2-}$$

$$(3) O_2^+ > O_2 > O_2^- > O_2^-$$

$$(4) O_2^+ > O_2^{2-} > O_2 > O_2^-$$

Ans. (2)

According to Molecular orbital theory

DO - TINE NET

© Resoriance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$O_2^+ = 2.5$$

$$O_2^- = 1.5$$

$$O_2^{2-}=1.0$$

| Colloid | Dispersion Medium | |
|-----------------|----------------------|--|
| a) Pumice Stone | (i) Liquid in Liquid | |
| b) Cloud | (ii) Gas in Solid | |
| c) Cheese | (iii) Liquid in Gas | |
| d) Hair Cream | (iv) Liquid in solid | |

The Correct option is

Ans.

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555 | 7340010333 | develop contribution of the contact of the contac

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

| | Dispersed phase | Disporators medium | Type of entired | |
|------|--------------------|-----------------------|--------------------|--------------------------------------|
| Sol. | Solid | Solid | Solid sol | Some coloured glasses and gem stone |
| | Solid | Liquid | Sol | Paints, cell fluids |
| | Solid | Cas | Aerosol | Smoke, dust |
| | Liquid | Solidi | Gel | Cheese, butter, jellies |
| | Liquid | Liquid | Emulsion | Milk, hair cream |
| | Liquid | Gan | Agrosol | Fog. mist. cloud, insecticide sprays |
| | Gas | Solid | Selid sol | Pumice stone, foam rubber |
| | Gas | Liquid- | Feam | Freth, whipped cream, sosp lather |

In Ho3+ [Atomic No. = 67), number of 4f electron are:

Ans. 10

Sol. Holmium (Z = 67): 4f11, 6s2

Ho+3: 4f 10

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

| Column - I | I Column - II | | |
|------------|---|--|--|
| a) Li | (i) I- is least soluble | | |
| b) Na | (ii) Bicarbonate is used in fire extinguisher | | |
| c) K | (iii) Carbonate easily decomposed on heating | | |
| d) Cs | (iv) Has vital role in biological system | | |

The Correct option is:

(1) a-(iv) b-(iii) c-(ii) d-(i)

(2) a-(iii) b-(ii) c-(iv) d-(i)

(3) a-(i) b-(iv) c-(ii) d-(iii)

(4) a-(iv) b-(ii) c-(iii) d- (i)

Ans. (2)

(i) Li₂CO₃

△ Li₂O + CO₂ Sol.

(ii) NaHCO₃ is used in dry fire extinguishers.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

The concentration of H_3O^+ ions in 0.005 M solution of $Ba(OH)_2$ at 298 K is $[x] \times 10^{-12}$. Assume that Ba(OH)2 is completely ionized under given conditions.

Ans. 1

Ba(OH)₂ ── Ba+2 + 2OH-Sol. 0.005 M 0.005 M 0.010 M

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Respondence" | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | CHEMISTRY

Now [H+] [OH-] = Kw

$$[H_3O^*] = \frac{K_w}{[OH^-]}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- Which form interstitial hydride easily?
 - (1) Fe
- (2) Cr
- (3) Ni
- (4) Co

Ans. (2)

- These are formed by many d-block and f-block elements. However, the metals of group 7, 8 and 9 do Sol. not form hydride. Even from group 6, only chromium forms CrH.
- 11. Match List-I with List-II

Column-

(a) Froth Floatation

- (b) Bessemer convertor
- (c) Blast furnance
- Column-II
- (i) Sulphide ore

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- (ii) Pig iron
- (iii) Ag

- (3) a-(iv) b-(ii) c-(iii) d-(i)
- (4) a-(iii) b-(iv) c-(ii) d-(i)

Ans. (2)

- In which of the following reaction oxidation state changes by 5.
 - (1) $Cr_5O_7^{2-} \rightarrow Cr^{+3}$

(2) $MnO_4^- \rightarrow Mn^{2+}$

(3) $C_2O_4^2 \rightarrow CO_2$

(4) $CrO_4^{2-} \rightarrow Cr^{+3}$

- Ans.
- Sol.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.acin | E-mail: contact@resonance.acin | CIN: U80302RU2007PLC024029

Toll Free: 1800 258 5555 📓 7340010333 👔 teostonic com/fleschenceddo 💟 twitter, com/fleschencedd. 🛗 www.youtube.com/fr

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#4

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- In which of the following compounds one π bond is present and maximum canonical structures possible.
 - (1) 503
- (2) CO₃²
- (3) O₂

Ans. (2)

3 canonical structures

An e-moving with a velocity of 2 × 10° m/s. If the speed can be measured with an accuracy of 0.02% calculate the uncertainty in its position is 1.45 x 10-x. The value of x

Ans. (7)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\Delta x.m\Delta V = \frac{h}{4\pi}$$

$$\Delta v = 2 \times 10^6 \times \frac{0.02}{100}$$

 $\Delta v \rightarrow 400 \text{ m/s}$

$$\Delta x = \frac{h}{4\pi \times m \ \Delta V} = \frac{6.63 \times 10^{-34}}{4 \times 3.14 \times 9.1 \times 10^{-31} \times 400 \ m/s} = 1.45 \times 10^{-7} s$$

15.

In this reaction, concentration of B changes by 0.2 in 30 minutes. The average rate of the reaction is x × 10-1 moles per litre hour. The value of x is:

Ans. 4

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

0.4 mole/L hr.

- Which among the following compounds is most stable: 16.
 - (1) [Cr(en)₂(NH₃)₂]Cl₃ (2) [Cr(en)₃]Cl₃
- (3) [Cr(en) (NH₃)₄]Cl₃ (4) [Cr (NH₃)₆]Cl₃

Ans. (2)

Chelation due to bidentateligand. Greater the chelation greater is the stability. Sol.

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

RESONANCE* | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | CHEMISTRY

In Kjeldahl's method, 0.8g of organic compound is used. The percentage of Nitrogen came out to be 42%. The _____ ml of 1M H₂SO₄ used to neutralize ammonia.

(1) 17 (2) 20

Ans.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302R.J2007PLC024029

$$42 = \frac{1.4 \times (1 \times 2) \times 1}{0.9}$$

 $V = 12 \, ml$

18 Find the product 'P'.

$$NO_2$$
 $Sn + HCI \rightarrow (A)$
 (P)

$$(1) H_2 N \longrightarrow N = N$$

$$(2) H_2N \longrightarrow N = N \longrightarrow$$

- Ans. (2)

(Racemic mixture)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#6

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

| JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | CHEMISTRY

- 20. Biodegradable polyamide is formed by-
 - (1) Glycine + isoprene
- (2) Glycine + Aminocaproic acid
- (3) Alanine + chloroprene
- (4) Acrylonitrile + Aminocaproic acid

- Ans. (2)
- Sol. Nylon 2-Nylon 6 (Polyamide copolymer) is biodegradable polymer.

Its monomer units are : Glycine + Aminocaproic acid

[H₂N - CH₂ - COOH + NH₂(CH₂)₅ - COOH]

Glycine

A minocaproic acid

- 21. Benzenenitrile with grignard reagent form product (P), which of the following chemical test given by product (P)
 - (1) Schiffe research (2) Ladeform (2) Nin

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol.
$$C \equiv N$$
 CH_3 C


22. Correct order of acidic strength form following compounds

- (1) a > b > c > d
- (2) d > c > b > a (3) b > c > d > a
- (4) c > b > a > d

- Ans. (2)
- Sol. Acidic strength & stability of conjugate base

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac_in | CIN - U80302RJ2007PLC024029

23. Structure of cytosine is



Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal PAGE#7

Resonance | Jee Main-2021 | Date : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | CHEMISTRY

CH₃ NH₂ (1) NaNO₂+HCl > Product (P) is: (2)H2O

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Ans. (3)

CH₃ (1)NaNO2+HCI (2)H₂O

25. S1: CFCs are dissociated with to CI radical by radiation of visible region.

CHa

S2: O3 reacts with nitric oxide to form N2 & O2

(1) False, True

(2) False, False

(3) True, False

(4) True, True

Ans. (2)

CFCs + UV --→CI* Sol.

0-110 -10-10-

@ Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- increasing order or density
 - (I) Benzene

(II) 1,3-Dichlorobenzene

(III) Chloro benzene

(IV) 1-Bromo-3-chlorobenzene

- (1) |V > || > || > |
- (2) | V > | | > | > |
- (3) ||| > || > |V > | (4) || > || > |V |

Ans. (1)

Higher the molecular weight higher will be density

- 27. Maleic anhydride can be prepared by.
 - (1) Treating cis but-2-ene-1,4-dioic acid with alcohol.
 - (2) Heating cis but-2-ene-1,4-dioic acid
 - (3) Treating trans but-2-ene-1,4-dioic acid with alcohol and acid
 - (4) Heating trans but-2-ene-idiric acid

(2) Ans.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Sol. COOH Maleic acid

Resonance Eduventures Ltd.


Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.acin | E-mail: contact@resonance.acin | CIN: U80302RJ2007PLC024029

Toll Free : 1800 258 5555 🔘 7340010333 📓 feeterkoom/Recommenda 💟 twittencom/Recommenda. 🔝 www.youtube.com/recovate. 🕒 biog recommenda.e.e.

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#8





COMPUTER BASED TEST (CBT)

Date: 25 July, 2021 (SHIFT-2) | TIME: (3.00 p.m. to 6.00 p.m)

Duration: 3 Hours | Max. Marks: 300

SUBJECT: MATHEMATICS

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005
Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: Sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302R J2007PL C024029

Toll Free: 1800 258 5555 340010333 westerscord in Max. Mill. Jhalawar Road | Max. Marks: 300

This solution was download from Resonance JEE (MAIN) 2021 Solution portal







ADMISSION OPEN for Session 2021-22

ONLINE + OFFLINE PROGRAMS

CLASS 11, 12 & 12+

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Scholarship Upto 90%*

Toll Free: 1800 258 5555 | Visit us: www.resonance.ac.in



Resonance³ | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

- If matrix $P = \begin{bmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 1 \end{bmatrix}$, then the matrix P^{50} is equal to

Sol.
$$P^{2} = \begin{bmatrix} \frac{1}{2} & 0 \\ \frac{1}{2} & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ \frac{1}{2} & 1 \end{bmatrix}$$
$$P^{3} = \begin{bmatrix} \frac{1}{2} & 0 \\ \frac{1}{2} & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & 1 \\ \frac{1}{2} & 1 \end{bmatrix}$$
$$P^{3} = \begin{bmatrix} \frac{1}{3} & 0 \\ \frac{3}{2} & 1 \end{bmatrix}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$P^{50} = \begin{bmatrix} 1 & 0 \\ 25 & 1 \end{bmatrix}$$

If the first sample A of 100 items has the mean 15 and standard deviation 3 and second sample B has 150 items. If the combined mean and standard deviation of items of both the sample is 15.6 and √13.44 . Then the standard deviation of items of sample B is :

Ans. (4) Sol. Combined mean = 15.6 $\therefore 15.6 = \frac{100 \times 15 + 150 \times \overline{x}}{15.00 \times 15}$ $\Rightarrow \overline{x}_{B} = 16$ (mean of sample B) Combined standard deviation = √13.44 \Rightarrow combined variance (σ^2) = 13.44 © Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$13.44 = \frac{\sum x_i^2}{250} - 243.36$$

$$\Rightarrow \sum x_i^2 = 64200$$
(1)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029 Toll Free : 1800 258 5555 🕲 7340010333 🌃 bonlook con/Reconancedul 💟 twitter.com/Resonancedul 🚵 www.youtube.com/resonance.ac.n

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 27777777, 27777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

RESONANCE* | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

$$9 = \frac{\sum x_{A}^{2}}{100} - 225$$

$$\Rightarrow \sum x_{A}^{2} = 23400$$

$$\Rightarrow \sum x_{ss}^{2} = 64200 - 23400 = 40800$$
standard deviation of sample B will be

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\sqrt{\frac{1}{n_B}} - (x_B)^2 = \sqrt{\frac{43000}{150}} - 256 = 4 \text{ Ans}$$

The value of $x \in \left[\frac{\pi}{4}, \frac{\pi}{4}\right]$ for which

$$(1) -\frac{\pi}{4}$$

$$(2) - \frac{\pi}{8}$$

(3)
$$\frac{\pi}{4}$$

(4)
$$\frac{\pi}{8}$$

Ans.

 $R_1 \rightarrow R_1 + R_2 + R_3$

$$\cos x \qquad \cos x \qquad \sin x$$

$$\left(\sin x + 2\cos x\right) \cos x \qquad \sin x \qquad \cos x$$

$$\cos x \cos x \qquad \sin x \qquad \cos x$$

$$\cos x \cos x \qquad \sin x$$

$$C_2 \rightarrow C_2 - C_1, C_3 \rightarrow C_3 - C_1$$

$$\left(\sin x + 2\cos x\right) \cos x \qquad \sin x - \cos x$$

$$\cos x \qquad 0 \qquad \sin x - \cos x$$

$$\left(\sin x + 2\cos x\right) \left(\sin x - \cos x\right)^2 = 0$$

$$\sin x = \cos x \qquad \text{or} \qquad \sin x = -2\cos x$$

$$\tan x = 1 \qquad \text{or} \qquad \tan x = -2$$

$$\therefore x \in \left[-\frac{\pi}{12}, \frac{\pi}{12}, \frac{\pi}{12}\right]$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
3 7340010333
Toolkok contribusional Resonanced with a way you also contribusion ac. in | www.you also contribusion ac. in | CIN: U80302RJ2007PLC024029 | Toll Free: 1800 258 5555
3 7340010333
Toolkok contribusional Resonanced with a way you also contribusion ac. in | CIN: U80302RJ2007PLC024029 | Toll Free: 1800 258 5555
3 7340010333
Toolkok contribusion ac. in | CIN: U80302RJ2007PLC024029 | Toll Free: 1800 258 5555
3 7340010333
Toolkok contribusion ac. in | CIN: U80302RJ2007PLC024029 | Toll Free: 1800 258 5555
3 7340010333
Toolkok contribusion ac. in | CIN: U80302RJ2007PLC024029 | Toll Free: 1800 258 5555
3 7340010333
Toolkok contribusion ac. in | CIN: U80302RJ2007PLC024029 | Toll Free: 1800 258 5555
3 7340010333
Toolkok contribusion ac. in | CIN: U80302RJ2007PLC024029 | Toll Free: 1800 258 5555
3 7340010333
Toolkok contribusion ac. in | CIN: U80302RJ2007PLC024029 | Toll Free: 1800 258 5555
3 7340010333
Toolkok contribusion ac. in | CIN: U80302RJ2007PLC024029 | Toll Free: 1800 258 5555
3 7340010333
Toolkok contribusion ac. in | CIN: U80302RJ2007PLC024029 | Toll Free: 1800 258 5555 | Toll Free: 1800 258

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

If a and b are two vectors such that $|\vec{a} \times \vec{b}| = 8$, $|\vec{a}| = 2$, $|\vec{b}| = 5$, then the value of $|\vec{a}|$ is:

(1) 6

(2)3

(3)12

Ans. (1)

Sol. a×b = absin⊕

 $\sin \theta = \frac{4}{5} \Rightarrow \cos \theta = \pm \frac{3}{5} \Rightarrow |\cos \theta| = \frac{3}{5}$

 $|\vec{a}.\vec{b}| = |\vec{a}| |\vec{b}| |\cos \theta| = 2 \times 5 \times \frac{3}{5} = 6$

If function $f(x):A\to B$, and $g(x):B\to C$ are defined such that $(g(f(x)))^{-1}$ exist then f(x) and g(x) are f(x) are f(x) and f(x) are f(x) are f(x) are f(x) and f(x) are f(x) are f(x) and f(x) are f(x) and f(x) are f(x)

(1) one-one and onto

(2) many-one and onto

(3) one-one and into

(4) many-one and into

Ans. (1)

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

If a + b + c = 1, ab + bc + ca = 2 and abc = 3, then the value of $a^4 + b^4 + c^4$ is:

Ans. 13

Sol. $(a + b + c)^2 = 1$

 \Rightarrow a² + b² + c² + 2(ab + bc + ca) = 1

 $\Rightarrow a^2 + b^2 + c^2 = -3 \qquad(i)$

 \Rightarrow ab + bc + ca = 2 Squaring of equation (ii),

 \Rightarrow a²b² + b²c² + c²a² + 2(ab²c + bc²a + ca²b) = 4

 $\Rightarrow a^2b^2 + b^2c^2 + c^2a^2 + 2abc(a + b + c) = 4$

 \Rightarrow $a^2b^2 + b^2c^2 + c^2a^2 + 6 = 4$

 $\Rightarrow a^2b^2 + b^2c^2 + c^2a^2 = -2$

Squaring of equation (i), \Rightarrow a⁴ + b⁴ + c⁴ + 2(a²b² + b²c² + c²a²) = 9

 \Rightarrow a⁴ + b⁴ + c⁴ - 4 = 9

 \Rightarrow a⁴ + b⁴ + c⁴ = 13

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

So,
$$\left(1+\frac{1}{n}\right)^n = {}^nC_0 + {}^nC_1\left(\frac{1}{n}\right) + {}^nC_2\left(\frac{1}{n}\right)^2 + {}^nC_3\left(\frac{1}{n}\right)^3 + \dots$$

$$= 1 + 1 + \frac{n(n-1)}{2n^2} + \frac{n(n-1)(n-2)}{6n^3} + \dots$$

$$\Rightarrow \left(1+\frac{1}{n}\right)^n > 2$$

Also $\lim_{n \to \infty} \left(1 + \frac{1}{n} \right)^n = e < 3$.

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Resonance* | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

- If a rectangle is inscribed in an equilateral triangle of side $2\sqrt{2}$, then the square of maximum area of rectangle is:
- Ans. (3)

Sol.



© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029



Area of rectangle = x. h

from ABDE

h = BE tan 60

$$h = \frac{\left(2\sqrt{2} - x\right)}{2} \sqrt{3}$$

So area, $A = \frac{\sqrt{3}}{2} (2\sqrt{2} \times -x^2)$

For maxima $\frac{dA}{dx} = \frac{\sqrt{3}}{2}(2\sqrt{2} - 2x) = 0$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

From (ii) h =
$$\sqrt{\frac{3}{3}}$$

 $(Area)^2 = 3$

- 9. If the coefficients of x^7 and x^8 in the expansion of $\left(2+\frac{x}{3}\right)^n$ are equal then the value of n is :
- (2) 54

Ans. (3)

Sol.
$$\left(2+\frac{x}{3}\right)^n = \sum_{r=0}^n {^nC_r} 2^{n-r} \left(\frac{x}{3}\right)^r$$

Coefficient of
$$\sqrt{7} = 0.0220-7$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744-2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Coefficient of
$$x^8 = {}^nC_8 2^{n-8}$$
. $\left(\frac{1}{3}\right)^n$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Tall Free: 1800 258 5555
7340010333
To extent confliction models with the confliction models with

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#4

$$\therefore {^{11}C_7} \frac{2^{n-7}}{3^7} = {^{11}C_8} \frac{2^{n-8}}{3^8}$$

$$\Rightarrow {^{n}C_{7}} \cdot 6 = {^{n}C_{8}}$$

$$\Rightarrow \frac{6n!}{7! \cdot (n-7)!} = \frac{n!}{8! \cdot (n-8)!}$$

$$\Rightarrow$$
 48 = n $-7 \Rightarrow$ n = 55

10. If
$$f(x) = \begin{cases} \frac{P(x)}{x-2} & |x| \neq 2 \\ x-2 & \text{and } P(x) \text{ is a polynomial such that } P''(x) \text{ is constant and } P(3) = 9. \text{ If } f(x) \text{ is continuous} \end{cases}$$

 $17 \quad x = 2$

at x = 2 then find the value of P(5)

Ans. (39)

<u>ana Bali ka awa sa</u>

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\lim_{x \to 2} f(x) = \lim_{x \to 2} f(x-2)$$

$$\Rightarrow K(2-\beta) = 7 \dots (1)$$

and
$$P(3) = K(3-2)(3-\beta) = 9$$

$$K(3-\beta) = 9 \dots (2)$$

Divide equation (1) by (2)

$$\frac{2-\beta}{3-\beta} = \frac{7}{9} \Rightarrow \beta = \frac{-3}{2}$$

Then P(x) =
$$2(x-2)\left(x+\frac{3}{2}\right)$$

$$P(5) = 2 \times (5-2) \times \left(5 + \frac{3}{2}\right) = 39$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777770 | contact@resonance ac.in | CIN - U80302R.J2007PLC024029

- 11. The number of real solutions of the equation $x^2 |x| 12 = 0$ is:
 - (1)0
- (2)1
- (3) 3
- (4) 2

Ans. (4)

Sol. $|x|^2 - |x| - 12 = 0$

|x| = 4, -3 (not possible)

$$\Rightarrow$$
 |x| = 4 \Rightarrow x = ± 4

.: Number of real solutions = 2

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 5

Resonance* | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

12. A coin is tossed natimes. If the probability of getting at least one head is greater than 0.9, then the minimum value of n is:

valu

- (1)3
- (2)5
- (3)4
- (4)2

Ans. (3)

(1)ⁿ

© Resonance Eduventures Limited | Toil-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow 0.1 > \left(\frac{1}{2}\right)^n \Rightarrow n = 4$$

- 13. Negation of the statement :
 - "We will play football only if ground is not wet and there is no sunlight" is:
 - (1) We will play football if ground is wet and there is no sunlight.
 - (2) We will play football if ground is wet and there is sunlight.
 - (3) There is no sunlight and ground is not wet and we will not play football.
 - (4) There is sunlight or ground is wet and we will play football.

Ans. (4

- Sol. p: We will play football.
 - q: Ground is not wet.
 - r: There is no sunlight.
 - ∴ Given statement is p → (q ∧ r)
 - Negation is p_∧ ~ (q_∧ r)
 - p ^ (~ q v ~ r)

(4)100

Ans. (4)

Sol.
$${}^{n}C_{0} + 3{}^{n}C_{1} + 5{}^{n}C_{2} + 7{}^{n}C_{3} + \dots (n+1) \text{ terms} = \sum_{r=0}^{n} (2r+1)^{n}C_{r}$$

$$=2\sum_{r=0}^{n}r^{r_{1}}C_{r}+\sum_{r=0}^{n}{}^{r_{1}}C_{r}$$

$$= 2n \cdot 2^{n-1} + 2^n = (n + 1) \cdot 2^n$$

15. Evaluate
$$\int_{1}^{1} \log \left(x + \sqrt{x^2 + 1} \right) dx$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

(3)96

Ans. (1)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
weekerk.om/flecoracedds
www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
weekerk.om/flecoracedds
www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac_in | CIN - U80302RJ2007PLC024029

TRESCITION ICE | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

Sol.
$$I = \int_{1}^{1} \log \left(x + \sqrt{x^2 + 1} \right) dx$$

$$f(x) = \log\left(\sqrt{x^2 + 1} + x\right)$$

$$f(-x) = \log\left(\sqrt{x^2 + 1} - x\right)$$

$$= -f(x)$$

So f(x) is a odd function.

$$\rightarrow 1 = 0$$

If ${}^nP_r = {}^nP_{r+1}$ and ${}^nC_r = {}^nC_{r-1}$, then the value of n is : 16.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029 Sol. ${}^{n}C_{r} = {}^{n}C_{r-1} \Rightarrow \frac{1}{r} = 1 \Rightarrow n+1 = 2r$ (1)

Sol.
$${}^{n}C_{r} = {}^{n}C_{r-1} \Rightarrow \frac{n+r-1}{r} = 1 \Rightarrow n+1 = 2r \dots (1)$$

and
$${}^{n}P_{r} = {}^{n}P_{r+1} \Rightarrow \frac{n!}{(n-r)!} = \frac{n!}{(n-r-1)!}$$

Solving (1) & (2)
$$n + 1 = 2(n - 1) \Rightarrow n = 3$$

17. Value of
$$\sum_{n=8}^{100} \left[(-1)^n \frac{n}{2} \right]$$
 is : (Where [.] represent greatest integer function)

Sol.
$$\sum_{n=0}^{100} \left[(-1)^n \frac{n}{2} \right]$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow$$
 -1 × 46 + 50 = 4

18. Evaluate
$$\cot\left(\frac{\pi}{24}\right)$$

(1)
$$\sqrt{6} - \sqrt{3} + \sqrt{2} - 2$$
 (2) $\sqrt{6} + \sqrt{3} - \sqrt{2} + 2$ (3) $\sqrt{6} + \sqrt{3} + \sqrt{2} - 2$ (4) $\sqrt{6} + \sqrt{3} + \sqrt{2} + 2$

Ans. (4)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE#7

RESONANCE | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

 $\cot\theta = \frac{\cos\theta}{\sin\theta} = \frac{2\cos^2\theta}{2\sin\theta\cos\theta}$

$$= \frac{1 + \cos 2\theta}{\sin 2\theta}$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow \frac{\frac{1+\sqrt{3}+1}{2\sqrt{2}}}{\frac{\sqrt{3}-1}{2\sqrt{2}}} = \frac{2\sqrt{2}+\sqrt{3}+1}{\sqrt{3}-1}$$

$$\Rightarrow \frac{\left(2\sqrt{2}+\sqrt{3}+1\right)\left(\sqrt{3}+1\right)}{2}$$

$$\Rightarrow \frac{2\sqrt{6} + 2\sqrt{2} + 3 + \sqrt{3} + \sqrt{3} + 1}{2} = \sqrt{6} + \sqrt{2} + \sqrt{3} + 2$$

19. If two lines $L_1 = \frac{x-k}{1} = \frac{y-2}{2} = \frac{z-3}{3}$ and $L_2 = \frac{x+1}{3} = \frac{y+2}{2} = \frac{z+3}{1}$ are coplanar. Then the value of k is:

Ans. (2) k+1 2+2 3+3

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow (k + 1) (-4) - 4(-8) + 6 (2-6) = 0$$
$$\Rightarrow (k + 1) (-4) = -8$$

20. If y = f(x) is the solution of differential equation $xdy = (y + x^3 \cos x)dx$ and $f(\pi) = 0$ then $f(\frac{\pi}{2})$ is:

(1) $\frac{\pi^2}{4} + \frac{\pi}{6}$ (2) $\frac{\pi^2}{4} + \frac{\pi}{2}$ (3) $\frac{\pi^2}{6} + \frac{\pi}{4}$ (4) $\frac{\pi^2}{6} + \frac{\pi}{6}$

Sol. $\frac{xdy - ydx}{x^2} = x \cos x dx$

 $\Rightarrow \int d\left(\frac{y}{x}\right) = \int x \cos x \, dx$

 $\Rightarrow \frac{y}{x} = x \sin x + \cos x + c$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

$$\Rightarrow f\left(\frac{\pi}{2}\right) = \frac{\pi^2}{4} + 0 + \frac{\pi}{2} = \frac{\pi^2}{4} + \frac{\pi}{2}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
Toll bree: vww.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
Toll bree: vww.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302RJ2007PLC024029

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

Ans. (4)

Sol.
$$|\vec{x} - \vec{y}| = n|\vec{x} + \vec{y}|$$

$$x^2 + y^2 - 2xy \cos\theta = n^2 (x^2 + y^2 + 2xy\cos\theta)$$

$$x^2(1+1-2\cos\theta) = n^2 x^2(1+1+2\cos\theta)$$

$$2(1-n^2) = 2\cos\theta (n^2 + 1)$$

$$\cos\theta = \frac{1-n^2}{n^2+1}$$

$$\theta = \cos^{-1}\left(\frac{1-n^2}{n^2+1}\right)$$

23. If combined equation of line y = p(x) and y = q(x) can be written as (y - p(x)) (y - q(x)) = 0 then angle bisector of $x^2 - 4xy - 5y^2 = 0$ is:

(1)
$$x^2 + 3xy + y^2 = 0$$

(2)
$$x^2 + 3xy - y^2 = 0$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance ac in | CIN - U80302RJ2007PLC024029

Ans. (2)

Sol. Equation of angle bisector of homogeneous equation of pair of straight line ax² + 2hxy + by² is

$$\frac{x^2 - y^2}{a - b} = \frac{xy}{h}$$

for
$$x^2 - 4xy - 5y^2 = 0$$

$$a = 1, h = -2, b = -5$$

so, equation of angle bisector is

$$\frac{x^2 - y^2}{1 - (-5)} = \frac{xy}{-2}$$

$$x^2 - y^2 + 3xy = 0$$

so, combined equation of angle bisector is $x^2 + 3xy - y^2 = 0$

Resonance Eduventures Ltd.

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 27777777, 2777700 | contact@resonance ac.in | CIN - U80302RJ2007PLC024029

Toll Free: 1800 258 5555
7340010333
Tollers conv. Hesonancelds
This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 9

RESONANCE | JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

24. Equation of a circle is $Re(z^2) + 2(Im(z))^2 + 2Re(z) = 0$ where z = x + iy and a line passes through the vertex of parabola $x^2 - 6x + y + 13 = 0$ and the centre of circle, then y intercept of the line is:

So,
$$z^2 = x^2 - y^2 + i2xy$$

Now
$$x^2-y^2 + 2y^2 + 2x = 0$$

$$x^2 + y^2 + 2x = 0 \Rightarrow$$
 centre = (-1, 0) and $x^2 - 6x + y + 13 = 0$

$$(x-3)^2 = -(y+4)$$

Vertex (3, -4)

$$\therefore \text{ Equation of line is } (y-0) = \frac{-4-0}{3+1} (x+1) \implies 4y = -4(x+1)$$

$$x + y + 1 = 0 \Rightarrow \frac{x}{-1} + \frac{y}{-1} = 1$$

25. If
$$f(x) = \begin{cases} 5x+1 & ; x < 2 \\ \int_0^x (5+|1-t|)dt & ; x \ge 2 \end{cases}$$

(1) f(x) is differentiable ∀x∈R

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

(3) f(x) is continuous at x = 2 but not differentiable at x = 1.

(4) f(x) is neither continuous nor differentiable at x = 2.

Ans. (2)

Sol. LHL =
$$\lim_{x \to 2^-} (5x+1) = 11$$

$$RHL = \lim_{x \to 2^+} \int_0^x (5+|1-t|) dt = \int_0^1 (5+(1-t)) dt + \int_1^2 (5-(1-t)) dt = 11$$

$$f(2) = 1$$

So, f(x) is continuous at x = 2

LHD at x = 2 is
$$\frac{d}{dx}(5x+1)\Big|_{x=2} = 5$$

RHD at x = 2 is
$$\frac{d}{dx} \int_0^x (5+|1-t|) dt \Big|_{x=2} = 6$$

© Resonance Eduventures Limited | Toll-Free 1800-258-5555 | (0)744 2777777, 2777700 | contact@resonance.ac.in | CIN - U80302RJ2007PLC024029

Resonance Eduventures Ltd.

Reg. Office & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005 Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more: sms RESO at 56677 | Website: www.resonance.ac.in | E-mail: contact@resonance.ac.in | CIN: U80302R12007PLC024029

Toll Free: 1800 258 5555
3 7340010333
Tooltok cont-Resonancetdi to bwtter com/Resonancetdi www.youtule.com/resonance ac.e.

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 10