


Math Refresher for DS

Practical Session 5



girafe
ai



Coordinate Change for Linear Transforms

Change of Basis for Vectors

- V – a vector space.
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis.
- $x \in V$ – some vector.
- We know already that

Change of Basis for Vectors

- V – a vector space.
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis.
- $x \in V$ – some vector.
- We know already that

$$x_B = Mx_S, \quad M = M_{B \rightarrow S} = \left[[s_1]_B \mid \dots \mid [s_n]_B \right] \text{ – transition matrix.}$$

Change of Basis for Vectors

- V – a vector space.
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis.
- $x \in V$ – some vector.
- We know already that

$$x_B = Mx_S, \quad M = M_{B \rightarrow S} = \left[[s_1]_B \mid \dots \mid [s_n]_B \right] \text{ – transition matrix.}$$

- We also established that

$$\exists M^{-1} = M_{S \rightarrow B}, \quad x_S = M^{-1}x_B$$

Change of Basis for Vectors

- V – a vector space.
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis.
- $x \in V$ – some vector.
- We know already that

$$x_B = Mx_S, \quad M = M_{B \rightarrow S} = \left[[s_1]_B \mid \dots \mid [s_n]_B \right] \text{ – transition matrix.}$$

- We also established that

$$\exists M^{-1} = M_{S \rightarrow B}, \quad x_S = M^{-1}x_B$$

- But vectors aren't the only things with coordinates...

Change of Basis for Linear Transforms

- Consider a linear transform A .
- It's defined by its matrix: columns = what happens to basis vectors.
- Example: rotation

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

Change of Basis for Linear Transforms

- Consider a linear transform A .
- It's defined by its matrix: columns = what happens to basis vectors.
- Example: rotation

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

- $S = \left\{ s_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$ – another basis.

How would A look like in this basis?

Change of Basis for Linear Transforms

- A – linear transform;
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis;
- $M = M_{B \rightarrow S}$ – transition matrix.
- x – some vector.

Change of Basis for Linear Transforms

- A – linear transform;
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis;
- $M = M_{B \rightarrow S}$ – transition matrix.
- x – some vector.

$$[x']_B = [A]_B \cdot [x]_B$$

Change of Basis for Linear Transforms

- A – linear transform;
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis;
- $M = M_{B \rightarrow S}$ – transition matrix.
- x – some vector.

$$[x']_B = [A]_B \cdot [x]_B$$

$$[x']_S = [A]_S \cdot [x]_S$$

Change of Basis for Linear Transforms

- A – linear transform;
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis;
- $M = M_{B \rightarrow S}$ – transition matrix.
- x – some vector.

$$[x']_B = [A]_B \cdot [x]_B$$

$$[x']_S = \quad \mathbf{[A]_S} \quad \cdot [x]_S$$

Change of Basis for Linear Transforms

- A – linear transform;
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis;
- $M = M_{B \rightarrow S}$ – transition matrix.
- x – some vector.

$$[x']_B = [A]_B \cdot [x]_B$$
$$[x']_S = \quad \cdot [x]_S$$

Change of Basis for Linear Transforms

- A – linear transform;
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis;
- $M = M_{B \rightarrow S}$ – transition matrix.
- x – some vector.

$$[x']_B = [A]_B \cdot [x]_B$$

$$[x']_S = M \cdot [x]_S$$

Change of Basis for Linear Transforms

- A – linear transform;
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis;
- $M = M_{B \rightarrow S}$ – transition matrix.
- x – some vector.

$$[x']_B = [A]_B \cdot [x]_B$$

$$[x']_S = [A]_B M \cdot [x]_S$$

Change of Basis for Linear Transforms

- A – linear transform;
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis;
- $M = M_{B \rightarrow S}$ – transition matrix.
- x – some vector.

$$[x']_B = [A]_B \cdot [x]_B$$

$$[x']_S = M^{-1} [A]_B M \cdot [x]_S$$

Change of Basis for Linear Transforms

- A – linear transform;
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis;
- $M = M_{B \rightarrow S}$ – transition matrix.
- x – some vector.

$$[x']_B = [A]_B \cdot [x]_B$$

$$[x']_S = M^{-1} [A]_B M \cdot [x]_S$$

Change of Basis for Linear Transforms

- A – linear transform;
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis;
- $M = M_{B \rightarrow S}$ – transition matrix.
- x – some vector.

$$[x']_B = [A]_B \cdot [x]_B$$

$$[x']_S = M^{-1} [A]_B M \cdot [x]_S$$

$$[A]_S =$$

Change of Basis for Linear Transforms

- A – linear transform;
- $B = \{b_1, \dots, b_n\}$ – current basis, $S = \{s_1, \dots, s_n\}$ – new basis;
- $M = M_{B \rightarrow S}$ – transition matrix.
- x – some vector.

$$[x']_B = [A]_B \cdot [x]_B$$

$$[x']_S = M^{-1} [A]_B M \cdot [x]_S$$

$$[A]_S = M^{-1} [A]_B M$$

Change of Basis for Linear Transforms

- Back to our example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad S \left\{ s_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis.}$$

$$[A]_S = M^{-1}AM =$$

Change of Basis for Linear Transforms

- Back to our example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad S \left\{ s_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis.}$$

$$[A]_S = M^{-1}AM = \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix} =$$

Change of Basis for Linear Transforms

- Back to our example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad S \left\{ s_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis.}$$

$$[A]_S = M^{-1}AM = \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1/3 & 1/3 \\ -1/3 & 2/3 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix} =$$

Change of Basis for Linear Transforms

- Back to our example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad S \left\{ s_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis.}$$

$$\begin{aligned} [A]_S &= M^{-1}AM = \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1/3 & 1/3 \\ -1/3 & 2/3 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix} = \\ &= \begin{bmatrix} 1/3 & -1/3 \\ 2/3 & 1/3 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = \end{aligned}$$

Change of Basis for Linear Transforms

- Back to our example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad S \left\{ s_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis.}$$

$$\begin{aligned} [A]_S &= M^{-1}AM = \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1/3 & 1/3 \\ -1/3 & 2/3 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix} = \\ &= \begin{bmatrix} 1/3 & -1/3 \\ 2/3 & 1/3 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 1/3 & -2/3 \\ 5/3 & -1/3 \end{bmatrix} \end{aligned}$$

Change of Basis for Linear Transforms

$$[A]_E = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, [A]_S = \begin{bmatrix} 1/3 & -2/3 \\ 5/3 & -1/3 \end{bmatrix}, S = \left\{ s_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis};$$

$$x_S = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad x'_S = [A]_S \cdot x_S =$$

Change of Basis for Linear Transforms

$$[A]_E = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, [A]_S = \begin{bmatrix} 1/3 & -2/3 \\ 5/3 & -1/3 \end{bmatrix}, S = \left\{ s_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis};$$

$$x_S = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad x'_S = [A]_S \cdot x_S = \begin{bmatrix} 1/3 & -2/3 \\ 5/3 & -1/3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} =$$

Change of Basis for Linear Transforms

$$[A]_E = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, [A]_S = \begin{bmatrix} 1/3 & -2/3 \\ 5/3 & -1/3 \end{bmatrix}, S = \left\{ s_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis};$$

$$x_S = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad x'_S = [A]_S \cdot x_S = \begin{bmatrix} 1/3 & -2/3 \\ 5/3 & -1/3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Change of Basis for Linear Transforms

$$[A]_E = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, [A]_S = \begin{bmatrix} 1/3 & -2/3 \\ 5/3 & -1/3 \end{bmatrix}, S = \left\{ s_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis};$$

$$x_S = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad x'_S = [A]_S \cdot x_S = \begin{bmatrix} 1/3 & -2/3 \\ 5/3 & -1/3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$x_E =$$

Change of Basis for Linear Transforms

$$[A]_E = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, [A]_S = \begin{bmatrix} 1/3 & -2/3 \\ 5/3 & -1/3 \end{bmatrix}, S = \left\{ s_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis};$$

$$x_S = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad x'_S = [A]_S \cdot x_S = \begin{bmatrix} 1/3 & -2/3 \\ 5/3 & -1/3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$x_E = \begin{bmatrix} 0 \\ 3 \end{bmatrix}, \quad x'_E =$$

Change of Basis for Linear Transforms

$$[A]_E = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, [A]_S = \begin{bmatrix} 1/3 & -2/3 \\ 5/3 & -1/3 \end{bmatrix}, S = \left\{ s_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis};$$

$$x_S = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad x'_S = [A]_S \cdot x_S = \begin{bmatrix} 1/3 & -2/3 \\ 5/3 & -1/3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$x_E = \begin{bmatrix} 0 \\ 3 \end{bmatrix}, \quad x'_E = [A]_E \cdot x_E = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 3 \end{bmatrix} = \begin{bmatrix} -3 \\ 0 \end{bmatrix}$$

Change of Basis for Linear Transforms

- Another example:

$$A = \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix}, \quad S = \left\{ s_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis}$$

$$[A]_S = M^{-1}AM =$$

Change of Basis for Linear Transforms

- Another example:

$$A = \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix}, \quad S = \left\{ s_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis}$$

$$[A]_S = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} =$$

Change of Basis for Linear Transforms

- Another example:

$$A = \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix}, \quad S = \left\{ s_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis}$$

$$[A]_S = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} =$$

Change of Basis for Linear Transforms

- Another example:

$$A = \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix}, \quad S = \left\{ s_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis}$$

$$[A]_S = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix}$$

Change of Basis for Linear Transforms

- Another example:

$$A = \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix}, \quad S = \left\{ s_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, s_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} - \text{new basis}$$

$$[A]_S = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix}$$

We get a diagonal matrix, it's easier to work with it!

Coordinate Change for Linear Transforms

Linear Transformations

- Every $n \times n$ matrix A represents a linear transformation of \mathbb{R}^n .
- Columns of A = what happens to the basis vectors.

Linear Transformations

- Another example:

$$A = \begin{bmatrix} 2 & 4 \\ 0 & 0 \end{bmatrix}$$

Linear Transformations

- Another example:

$$A = \begin{bmatrix} 2 & 4 \\ 0 & 0 \end{bmatrix}$$

Every vector in \mathbb{R}^2 is mapped onto a line, a one-dimensional subspace of \mathbb{R}^2
(but we still stay in \mathbb{R}^2)

$$Ax = \begin{bmatrix} 2 & 4 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2x_1 + 4x_2 \\ 0 \end{bmatrix}$$

Linear Transformations

- Another example:

$$A = \begin{bmatrix} 2 & 4 \\ 0 & 0 \end{bmatrix}$$

Every vector in \mathbb{R}^2 is mapped onto a line, a one-dimensional subspace of \mathbb{R}^2
(but we still stay in \mathbb{R}^2)

$$Ax = \begin{bmatrix} 2 & 4 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2x_1 + 4x_2 \\ 0 \end{bmatrix}$$

Many vectors are mapped onto the same one (no inverse!):

$$\text{Example: } \begin{bmatrix} 2 & 4 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 3 \\ 2 \end{bmatrix} = \begin{bmatrix} 14 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 5 \\ 1 \end{bmatrix} = \begin{bmatrix} 14 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 14 \\ 0 \end{bmatrix}$$

Linear Transformations

So far: only square matrices.

But about non-square ones?

Linear Transformations

$$A = \begin{bmatrix} 3 & 1 \\ 4 & 1 \\ 5 & 9 \end{bmatrix} - 3 \times 2 \text{ matrix.}$$

Linear Transformations

$$A = \begin{bmatrix} 3 & 1 \\ 4 & 1 \\ 5 & 9 \end{bmatrix} - 3 \times 2 \text{ matrix.}$$

$$x \in \mathbb{R}^2, \quad Ax = \begin{bmatrix} 3 & 1 \\ 4 & 1 \\ 5 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} =$$

Linear Transformations

$$A = \begin{bmatrix} 3 & 1 \\ 4 & 1 \\ 5 & 9 \end{bmatrix} - 3 \times 2 \text{ matrix.}$$

$$x \in \mathbb{R}^2, \quad Ax = \begin{bmatrix} 3 & 1 \\ 4 & 1 \\ 5 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ 14 \end{bmatrix} \in \mathbb{R}^3$$

Linear Transformations

$$A = \begin{bmatrix} 3 & 1 \\ 4 & 1 \\ 5 & 9 \end{bmatrix} - 3 \times 2 \text{ matrix.}$$

$$x \in \mathbb{R}^2, \quad Ax = \begin{bmatrix} 3 & 1 \\ 4 & 1 \\ 5 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ 14 \end{bmatrix} \in \mathbb{R}^3$$

A is a linear transformation from \mathbb{R}^2 to \mathbb{R}^3 !

Linear Transformations

$$A = \begin{bmatrix} 3 & 1 \\ 4 & 1 \\ 5 & 9 \end{bmatrix} - 3 \times 2 \text{ matrix.}$$

$$x \in \mathbb{R}^2, \quad Ax = \begin{bmatrix} 3 & 1 \\ 4 & 1 \\ 5 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ 14 \end{bmatrix} \in \mathbb{R}^3$$

A is a linear transformation from \mathbb{R}^2 to \mathbb{R}^3 !

$\text{rank}(A) = 2$: vectors that were independent in \mathbb{R}^2
will be mapped on independent vectors in \mathbb{R}^3 .

Linear Transformations

$$A = \begin{bmatrix} 3 & 1 & 4 \\ 1 & 5 & 9 \end{bmatrix} - 2 \times 3 \text{ matrix.}$$

Linear Transformations

$$A = \begin{bmatrix} 3 & 1 & 4 \\ 1 & 5 & 9 \end{bmatrix} - 2 \times 3 \text{ matrix.}$$

$$x \in \mathbb{R}^3, \quad Ax = \begin{bmatrix} 3 & 1 & 4 \\ 1 & 5 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} =$$

Linear Transformations

$$A = \begin{bmatrix} 3 & 1 & 4 \\ 1 & 5 & 9 \end{bmatrix} - 2 \times 3 \text{ matrix.}$$

$$x \in \mathbb{R}^3, \quad Ax = \begin{bmatrix} 3 & 1 & 4 \\ 1 & 5 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 8 \\ 15 \end{bmatrix} \in \mathbb{R}^2$$

Linear Transformations

$$A = \begin{bmatrix} 3 & 1 & 4 \\ 1 & 5 & 9 \end{bmatrix} - 2 \times 3 \text{ matrix.}$$

$$x \in \mathbb{R}^3, \quad Ax = \begin{bmatrix} 3 & 1 & 4 \\ 1 & 5 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 8 \\ 15 \end{bmatrix} \in \mathbb{R}^2$$

A is a linear transformation from \mathbb{R}^3 to \mathbb{R}^2 !

Linear Transformations

$$A = \begin{bmatrix} 3 & 1 & 4 \\ 1 & 5 & 9 \end{bmatrix} - 2 \times 3 \text{ matrix.}$$

$$x \in \mathbb{R}^3, \quad Ax = \begin{bmatrix} 3 & 1 & 4 \\ 1 & 5 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 8 \\ 15 \end{bmatrix} \in \mathbb{R}^2$$

A is a linear transformation from \mathbb{R}^3 to \mathbb{R}^2 !

$\text{rank}(A) = 2$: vectors that were independent in \mathbb{R}^3
may be mapped on dependent vectors in \mathbb{R}^2 .

Linear Transformations

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} - m \times n \text{ matrix.}$$

$$x \in \mathbb{R}^n, \quad Ax = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix} \in \mathbb{R}^m$$

A is a linear transformation from \mathbb{R}^n to \mathbb{R}^m !

Linear Transformations From \mathbb{R}^n to \mathbb{R}^m

Why this is useful?

Dimensionality reduction!

Dimensionality Reduction

Dimensionality Reduction


- Imagine that you have some data: m features, n examples.
- Each example $x = (x_1, \dots, x_m)$ – a point in \mathbb{R}^m .
- How to visualize this data?

Dimensionality Reduction


- Imagine that you have some data: m features, n examples.
- Each example $x = (x_1, \dots, x_m)$ – a point in \mathbb{R}^m .
- How to visualize this data?

Map into onto a lower-dimensional space!
But preserve as much variance in data as possible.


Dimensionality Reduction




Dimensionality Reduction



Dimensionality Reduction





Dimensionality Reduction



Dimensionality Reduction

- How to find this direction?



Let's review the theory around it

Eigenvalues & Eigenvectors

- Consider an $n \times n$ matrix A .
- A – linear transformation of \mathbb{R}^n . Every vector gets scaled and rotated:

$$x' = Ax$$

Eigenvalues & Eigenvectors

- Consider an $n \times n$ matrix A .
- A – linear transformation of \mathbb{R}^n . Every vector gets scaled and rotated:

$$x' = Ax$$

- Some vectors only get scaled:

$$Av = \lambda v, \quad \lambda \in \mathbb{R}, \quad v \neq 0$$

$v \neq 0$ – eigenvector, λ – corresponding eigenvalue.

Eigenvalues & Eigenvectors

- Consider an $n \times n$ matrix A .
- $v \neq 0$ – eigenvector, λ – corresponding eigenvalue:

$$Av = \lambda v$$

How to find λ and v ?

$$Av = \lambda v \Leftrightarrow Av - \lambda v = 0 \Leftrightarrow (A - \lambda E)v = 0 \Leftrightarrow$$

Eigenvalues & Eigenvectors

- Consider an $n \times n$ matrix A .
- $v \neq 0$ – eigenvector, λ – corresponding eigenvalue:

$$Av = \lambda v$$

How to find λ and v ?

$$Av = \lambda v \Leftrightarrow Av - \lambda v = 0 \Leftrightarrow (A - \lambda E)v = 0 \Leftrightarrow \det(A - \lambda E) = 0$$

$\det(A - \lambda E)$ – characteristic polynomial of A .

Eigenvalues & Eigenvectors

- Example: $A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$
- Let's find eigenvalues and eigenvectors of A .
- Characteristic polynomial:

$$\det(A - \lambda E) =$$

Eigenvalues & Eigenvectors

- Example: $A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$
- Let's find eigenvalues and eigenvectors of A .
- Characteristic polynomial:

$$\det(A - \lambda E) = \begin{vmatrix} -5 - \lambda & 2 \\ -7 & 4 - \lambda \end{vmatrix} =$$

Eigenvalues & Eigenvectors

- Example: $A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$
- Let's find eigenvalues and eigenvectors of A .
- Characteristic polynomial:

$$\det(A - \lambda E) = \begin{vmatrix} -5 - \lambda & 2 \\ -7 & 4 - \lambda \end{vmatrix} = (-5 - \lambda)(4 - \lambda) + 14 = 0 \Leftrightarrow$$

$$\lambda^2 + \lambda - 6 = 0 \Leftrightarrow$$

Eigenvalues & Eigenvectors

- Example: $A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$
- Let's find eigenvalues and eigenvectors of A .
- Characteristic polynomial:

$$\det(A - \lambda E) = \begin{vmatrix} -5 - \lambda & 2 \\ -7 & 4 - \lambda \end{vmatrix} = (-5 - \lambda)(4 - \lambda) + 14 = 0 \Leftrightarrow$$

$$\lambda^2 + \lambda - 6 = 0 \Leftrightarrow$$

$$\lambda_1 = 2, \lambda_2 = -3 \quad - \text{eigenvalues of } A.$$

Eigenvalues & Eigenvectors

- Example: $A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$
- Let's find eigenvalues and eigenvectors of A .

$\lambda_1 = 2, \lambda_2 = -3$ – eigenvalues of A .

$$Av_1 = \lambda_1 v_1 \Leftrightarrow$$

Eigenvalues & Eigenvectors

- Example: $A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$
- Let's find eigenvalues and eigenvectors of A .

$\lambda_1 = 2, \lambda_2 = -3$ – eigenvalues of A .

$$Av_1 = \lambda_1 v_1 \Leftrightarrow \begin{bmatrix} -5-2 & 2 \\ -7 & 4-2 \end{bmatrix} \begin{bmatrix} v_{11} \\ v_{21} \end{bmatrix} = 0 \Leftrightarrow$$

Eigenvalues & Eigenvectors

- Example: $A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$
- Let's find eigenvalues and eigenvectors of A .

$\lambda_1 = 2, \lambda_2 = -3$ – eigenvalues of A .

$$Av_1 = \lambda_1 v_1 \Leftrightarrow \begin{bmatrix} -5-2 & 2 \\ -7 & 4-2 \end{bmatrix} \begin{bmatrix} v_{11} \\ v_{21} \end{bmatrix} = 0 \Leftrightarrow v_{11} = \frac{2}{7} v_{21}$$

For example, $v_1 = \begin{bmatrix} 2 \\ 7 \end{bmatrix}$.

Eigenvalues & Eigenvectors

- Example: $A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$
- Let's find eigenvalues and eigenvectors of A .

$\lambda_1 = 2, \lambda_2 = -3$ – eigenvalues of A .

$$Av_2 = \lambda_2 v_2 \Leftrightarrow$$

Eigenvalues & Eigenvectors

- Example: $A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$
- Let's find eigenvalues and eigenvectors of A .

$\lambda_1 = 2, \lambda_2 = -3$ – eigenvalues of A .

$$Av_2 = \lambda_2 v_2 \Leftrightarrow \begin{bmatrix} -5 + 3 & 2 \\ -7 & 4 + 3 \end{bmatrix} \begin{bmatrix} v_{12} \\ v_{22} \end{bmatrix} = 0$$

Eigenvalues & Eigenvectors

- Example: $A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$
- Let's find eigenvalues and eigenvectors of A .

$\lambda_1 = 2, \lambda_2 = -3$ – eigenvalues of A .

$$Av_2 = \lambda_2 v_2 \Leftrightarrow \begin{bmatrix} -5 + 3 & 2 \\ -7 & 4 + 3 \end{bmatrix} \begin{bmatrix} v_{12} \\ v_{22} \end{bmatrix} = 0 \Leftrightarrow v_{12} = v_{22}$$

Eigenvalues & Eigenvectors

- Example: $A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$
- Let's find eigenvalues and eigenvectors of A .

$\lambda_1 = 2, \lambda_2 = -3$ – eigenvalues of A .

$$Av_2 = \lambda_2 v_2 \Leftrightarrow \begin{bmatrix} -5 + 3 & 2 \\ -7 & 4 + 3 \end{bmatrix} \begin{bmatrix} v_{12} \\ v_{22} \end{bmatrix} = 0 \Leftrightarrow v_{12} = v_{22}$$

For example, $v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

Eigenvalues & Eigenvectors

- Example: $A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$
- Eigenvalues and eigenvectors of A :

$$\lambda_1 = 2, \quad v_1 = \begin{bmatrix} 2 \\ 7 \end{bmatrix}$$

$$\lambda_2 = -3, \quad v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Eigenvalues & Eigenvectors

- Example: $A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$
- Eigenvalues and eigenvectors of A :

$$\lambda_1 = 2, \quad v_1 = \begin{bmatrix} 2 \\ 7 \end{bmatrix}$$

$$\lambda_2 = -3, \quad v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Note that v_1 and v_2 are linearly independent.

Eigenbasis

$$A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$$

- Eigenvalues and eigenvectors: $\lambda_1 = 2$, $v_1 = \begin{bmatrix} 2 \\ 7 \end{bmatrix}$, $\lambda_2 = -3$, $v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
- v_1 and v_2 are linearly independent. They form **eigenbasis**.
What happens to A if we change to basis $\{v_1, v_2\}$?

Eigenbasis

$$A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$$

- Eigenvalues and eigenvectors: $\lambda_1 = 2$, $v_1 = \begin{bmatrix} 2 \\ 7 \end{bmatrix}$, $\lambda_2 = -3$, $v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
- v_1 and v_2 are linearly independent. They form **eigenbasis**.
What happens to A if we change to basis $\{v_1, v_2\}$?

$[A]_V = \{\text{what happens to basis vectors } v_1, v_2 \text{ after applying } A\} =$

Eigenbasis

$$A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$$

- Eigenvalues and eigenvectors: $\lambda_1 = 2$, $v_1 = \begin{bmatrix} 2 \\ 7 \end{bmatrix}$, $\lambda_2 = -3$, $v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
- v_1 and v_2 are linearly independent. They form **eigenbasis**.
What happens to A if we change to basis $\{v_1, v_2\}$?

$$\begin{aligned} [A]_V &= \{\text{what happens to basis vectors } v_1, v_2 \text{ after applying } A\} = \\ &= \begin{bmatrix} 2 & 0 \\ 0 & -3 \end{bmatrix} = \Lambda - \text{it becomes diagonal!} \end{aligned}$$

Eigendecomposition

$$A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$$

- Eigenvalues and eigenvectors: $\lambda_1 = 2$, $v_1 = \begin{bmatrix} 2 \\ 7 \end{bmatrix}$, $\lambda_2 = -3$, $v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
- How do we change to basis $\{v_1, v_2\}$?

Eigendecomposition

$$A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$$

- Eigenvalues and eigenvectors: $\lambda_1 = 2$, $v_1 = \begin{bmatrix} 2 \\ 7 \end{bmatrix}$, $\lambda_2 = -3$, $v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
- How do we change to basis $\{v_1, v_2\}$?

$V = [v_1 \mid v_2]$ – transition from standard to eigenbasis.

$$A = V[A]_V V^{-1}$$

$A = V\Lambda V^{-1}$ – eigendecomposition.

Eigendecomposition

$$A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$$

- Eigenvalues and eigenvectors: $\lambda_1 = 2$, $v_1 = \begin{bmatrix} 2 \\ 7 \end{bmatrix}$, $\lambda_2 = -3$, $v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
- How do we change to basis $\{v_1, v_2\}$?

$$A = V\Lambda V^{-1} \text{ – eigendecomposition.}$$

Let's check this:

$$\begin{bmatrix} 2 & 1 \\ 7 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & -3 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 7 & 1 \end{bmatrix}^{-1} =$$

Eigendecomposition

$$A = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$$

- Eigenvalues and eigenvectors: $\lambda_1 = 2$, $v_1 = \begin{bmatrix} 2 \\ 7 \end{bmatrix}$, $\lambda_2 = -3$, $v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
- How do we change to basis $\{v_1, v_2\}$?

$$A = V\Lambda V^{-1} \text{ – eigendecomposition.}$$

Let's check this:

$$\begin{bmatrix} 2 & 1 \\ 7 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & -3 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 7 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 2 & 1 \\ 7 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & -3 \end{bmatrix} \begin{bmatrix} -0.2 & 0.2 \\ 1.4 & -0.4 \end{bmatrix} = \begin{bmatrix} -5 & 2 \\ -7 & 4 \end{bmatrix}$$

Eigendecomposition

Is it always possible to find an eigenbasis?

No ☹️ (see lectures for examples).

But there are good news 😊

The Spectral Theorem

If A is an $n \times n$ symmetric matrix,
then A always n linearly independent eigenvectors v_1, \dots, v_n .

What's more, v_1, \dots, v_n are mutually orthogonal!
Since we are choosing the scaling, we can make them orthonormal.

The Spectral Theorem

If A is an $n \times n$ symmetric matrix,
then A always n linearly independent eigenvectors v_1, \dots, v_n .

What's more, v_1, \dots, v_n are mutually orthogonal!
Since we are choosing the scaling, we can make them orthonormal.

So, we can always decompose square symmetric matrices as

$$A = V\Lambda V^{-1}, \text{ where}$$

V – orthogonal matrix (columns = eigenvectors),

Λ – diagonal matrix (diagonal elements = eigenvalues)

Orthogonal Matrices

- A matrix where all columns are mutually orthonormal:

$$A^T A = A A^T = E$$

Orthogonal Matrices

- A matrix where all columns are mutually orthonormal:

$$A^T A = A A^T = E$$

- That means that orthogonal matrices are easy to invert:

$$A^{-1} = A^T.$$

The Spectral Theorem

If A is an $n \times n$ symmetric matrix,
then A always n linearly independent eigenvectors v_1, \dots, v_n .

What's more, v_1, \dots, v_n are mutually orthogonal!
Since we are choosing the scaling, we can make them orthonormal.

So, we can always decompose square symmetric matrices as

$$A = V \Lambda V^{-1} = V \Lambda V^T, \text{ where}$$

V – orthogonal matrix (columns = eigenvectors),

Λ – diagonal matrix (diagonal elements = eigenvalues)

Back to Dimensionality Reduction



girafe
ai

Dimensionality Reduction

- Imagine that you have some data: m features, n examples.
- Each example $x = (x_1, \dots, x_m)$ – a point in \mathbb{R}^m .
- How to visualize this data?
- Map into onto a lower-dimensional space!
But preserve as much variance in data as possible.

Dimensionality Reduction

- How to find this direction?



PCA

- Let's construct data covariance matrix.

$$S = \frac{1}{n-1}XX^T - m \times m \text{ symmetric matrix.}$$

s_{ij} – covariance between features i and j ,

s_{ii} – variance of feature i

PCA

- Let's construct data covariance matrix.

$$S = \frac{1}{n-1}XX^T - m \times m \text{ symmetric matrix.}$$

s_{ij} – covariance between features i and j ,

s_{ii} – variance of feature i

- We can apply eigendecomposition to S :

$$S = V\Lambda V^T$$

PCA

$$S = V\Lambda V^{-1} = V\Lambda V^T$$

$$\begin{bmatrix} \mathbf{s}_{11} & s_{12} & \cdots & s_{1m} \\ s_{21} & \mathbf{s}_{22} & \cdots & s_{2m} \\ \vdots & \vdots & \cdots & \vdots \\ s_{m1} & s_{m2} & \cdots & \mathbf{s}_{mm} \end{bmatrix} =$$

$$= \begin{bmatrix} v_{11} & v_{12} & \cdots & v_{1m} \\ v_{21} & v_{22} & \cdots & v_{2m} \\ \vdots & \vdots & \cdots & \vdots \\ v_{m1} & v_{m2} & \cdots & v_{mm} \end{bmatrix} \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & \cdots & \lambda_m \end{bmatrix} \begin{bmatrix} v_{11} & v_{12} & \cdots & v_{1m} \\ v_{21} & v_{22} & \cdots & v_{2m} \\ \vdots & \vdots & \cdots & \vdots \\ v_{m1} & v_{m2} & \cdots & v_{mm} \end{bmatrix}^T$$

PCA

$$S = V\Lambda V^{-1} = V\Lambda V^T$$

$$\begin{bmatrix} \mathbf{s}_{11} & s_{12} & \cdots & s_{1m} \\ s_{21} & \mathbf{s}_{22} & \cdots & s_{2m} \\ \vdots & \vdots & \cdots & \vdots \\ s_{m1} & s_{m2} & \cdots & \mathbf{s}_{mm} \end{bmatrix} =$$

$$= \begin{bmatrix} v_{11} & v_{12} & \cdots & v_{1m} \\ v_{21} & v_{22} & \cdots & v_{2m} \\ \vdots & \vdots & \cdots & \vdots \\ v_{m1} & v_{m2} & \cdots & v_{mm} \end{bmatrix} \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & \cdots & \lambda_m \end{bmatrix} \begin{bmatrix} v_{11} & v_{12} & \cdots & v_{1m} \\ v_{21} & v_{22} & \cdots & v_{2m} \\ \vdots & \vdots & \cdots & \vdots \\ v_{m1} & v_{m2} & \cdots & v_{mm} \end{bmatrix}^T$$

Total variance of the data

$$T = \text{tr}(S) = s_{11} + \cdots + s_{nn} = \lambda_1 + \cdots + \lambda_m$$

Projecting the Data

- What happens if we project the data onto the eigenvectors of S ?
Remember: $\{v_1, \dots, v_m\}$ is an orthonormal basis!

Projecting the Data

- What happens if we project the data onto the eigenvectors of S ?
Remember: $\{v_1, \dots, v_m\}$ is an orthonormal basis!

$$x = x'_1 v_1 + x'_2 v_2 + \dots + x'_m v_m,$$

Projecting the Data

- What happens if we project the data onto the eigenvectors of S ?
Remember: $\{v_1, \dots, v_m\}$ is an orthonormal basis!

$$x = x'_1 v_1 + x'_2 v_2 + \dots + x'_m v_m, \quad (x, v_i) = x'_i (v_i, v_i) + 0 = x_i$$

Coordinates in an orthonormal basis = dot products!

Projecting the Data

- What happens if we project the data onto the eigenvectors of S ?
Remember: $\{v_1, \dots, v_m\}$ is an orthonormal basis!

$$x = x'_1 v_1 + x'_2 v_2 + \dots + x'_m v_m, \quad (x, v_i) = x'_i (v_i, v_i) + 0 = x_i$$

Coordinates in an orthonormal basis = dot products!

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_m \end{bmatrix} \text{ -- one example,} \quad x_{proj} = V^T x$$

Projecting the Data

- What happens if we project the data onto the eigenvectors of S ?
Remember: $\{v_1, \dots, v_m\}$ is an orthonormal basis!

$$x = x'_1 v_1 + x'_2 v_2 + \dots + x'_m v_m, \quad (x, v_i) = x'_i (v_i, v_i) + 0 = x_i$$

Coordinates in an orthonormal basis = dot products!

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_m \end{bmatrix} \text{ -- one example,} \quad x_{proj} = V^T x$$

$$\text{The whole data: } X_{proj} = V^T X$$

Projecting the Data

- How to project the data onto $p < m$ eigenvectors of S ?

Projecting the Data

- How to project the data onto $p < m$ eigenvectors of S ?

$$V_p = \{v_1 \mid \dots \mid v_p\}$$

Projecting the Data

- How to project the data onto $p < m$ eigenvectors of S ?

$$V_p = \{v_1 \mid \dots \mid v_p\}$$

$$X_p = V_p^T X$$

Projecting the Data

- How to project the data onto $p < m$ eigenvectors of S ?

$$V_p = \{v_1 \mid \dots \mid v_p\}$$

$$X_p = V_p^T X$$

- What happens to the covariance matrix?

$$S_{proj} = \frac{1}{n-1} X_p X_p^T = \frac{1}{n-1} V_p^T X X^T V_p$$

Let's practice!