


Math Refresher for DS

Lecture 2



girafe
ai



Last Time

- Vector spaces
- Euclidian spaces (= vector spaces + dot product)
- Length of a vector
- Distances and angles between the vectors
- Orthogonality
- Orthogonal projections

Today



- Back to vector spaces
 - Linear independence
 - Basis
- Basic operations with matrices.



Back to Vector Spaces



(Reminder) Vector Space: Definition

- A real-valued vector space $(V, +, \cdot)$ is a set of vectors V with two operations

$$(1) +: V \times V \rightarrow V, \quad (2) \cdot: \mathbb{R} \times V \rightarrow V$$

that satisfy the following properties (axioms):

	Property	Meaning
1.	Associativity of addition	$x + (y + z) = (x + y) + z$
2.	Commutativity of addition	$x + y = y + x$
3.	Identity element of addition	$\exists 0 \in V: \forall x \in V \quad 0 + x = x$
4.	Identity element of scalar multiplication	$\forall x \in V \quad 1 \cdot x = x$
5.	Inverse element of addition	$\forall x \in V \exists -x \in V: x + (-x) = 0$
6.	Compatibility of scalar multiplication	$\alpha(\beta x) = (\alpha\beta)x$
7.	Distributivity	$(\alpha + \beta)x = \alpha x + \beta x$
8.		$\alpha(x + y) = \alpha x + \alpha y$

(Reminder) Examples of Vector Spaces

- \mathbb{R}^n - a set of vectors with n real entries.
 $(\mathbb{R}^n, +, \cdot)$ is a vector space.

(Reminder) Examples of Vector Spaces

- \mathbb{R}^n - a set of vectors with n real entries.
 $(\mathbb{R}^n, +, \cdot)$ is a vector space.
- \mathbb{P}^n - a set of polynomials of degree $\leq n$ with real coefficients
 $(\mathbb{P}^n, +, \cdot)$ is also a vector space!
“Vectors” here are polynomials.

Vector Subspaces



Vector Subspace

- $V = (\mathbb{V}, +, \cdot)$ - a vector space.

Vector Subspace

- $V = (\mathbb{V}, +, \cdot)$ - a vector space.
- Consider $\mathbb{U} \neq \emptyset$ - a subset of \mathbb{V} ($\mathbb{U} \subseteq \mathbb{V}$).

Vector Subspace

- $V = (\mathbb{V}, +, \cdot)$ - a vector space.
- Consider $\mathbb{U} \neq \emptyset$ - a subset of \mathbb{V} ($\mathbb{U} \subseteq \mathbb{V}$).
- $U = (\mathbb{U}, +, \cdot)$ - a *vector subspace* ($U \subseteq V$) if U is a vector space with operations
 - $+: \mathbb{U} \times \mathbb{U} \rightarrow \mathbb{U}$
 - $\cdot: \mathbb{R} \times \mathbb{U} \rightarrow \mathbb{U}$

Vector Subspace

- How do we check if $U = (\mathbb{U}, +, \cdot)$ is a vector space?

Vector Subspace

- How do we check if $U = (\mathbb{U}, +, \cdot)$ is a vector space?
- Since $\mathbb{U} \subseteq \mathbb{V}$ and $V = (\mathbb{V}, +, \cdot)$ is a vector space, many properties of $+$ and \cdot hold automatically.

Vector Subspace

- How do we check if $U = (\mathbb{U}, +, \cdot)$ is a vector space?
- Since $\mathbb{U} \subseteq \mathbb{V}$ and $V = (\mathbb{V}, +, \cdot)$ is a vector space, many properties of $+$ and \cdot hold automatically.
- In fact, we only need to check:
 1. that $0 \in \mathbb{U}$


Vector Subspace

- How do we check if $U = (\mathbb{U}, +, \cdot)$ is a vector space?
- Since $\mathbb{U} \subseteq \mathbb{V}$ and $V = (\mathbb{V}, +, \cdot)$ is a vector space, many properties of $+$ and \cdot hold automatically.
- In fact, we only need to check:
 1. that $0 \in \mathbb{U}$
 2. closure of $+$ and \cdot :
 - $\forall x, y \in \mathbb{U} \ x + y \in \mathbb{U}$
 - $\forall x \in \mathbb{U}, \lambda \in \mathbb{R} \ \lambda x \in \mathbb{U}$

Vector Subspace: Examples




- Consider \mathbb{R}^2 .
- Is A a vector subspace?



Vector Subspace: Examples




- Consider \mathbb{R}^2 .
- Is A a vector subspace?



Vector Subspace: Examples



- Consider \mathbb{R}^2 .
- Is **A** a vector subspace?




Vector Subspace: Examples



- Consider \mathbb{R}^2 .
- Is A a vector subspace?
- No!


$x \in A$ but $2x \notin A \rightarrow$
• operation isn't closed.



Vector Subspace: Examples



- Consider \mathbb{R}^2 .
- Is B a vector subspace?




Vector Subspace: Examples



- Consider \mathbb{R}^2 .
- Is B a vector subspace?
- No!


$$0 \notin B$$



Vector Subspace: Examples




- Consider \mathbb{R}^2 .
- Is D a vector subspace?



Vector Subspace: Examples



- Consider \mathbb{R}^2 .
- Is D a vector subspace?
- Yes!






Vector Subspace: Examples

- Consider \mathbb{R}^2 .
- Is D a vector subspace?
- Yes!

$\{0, +, \cdot\}$ is a trivial vector subspace of any vector space



Vector Subspace: Examples

- Consider $P^n = (\mathbb{P}^n, +, \cdot)$ - a vector space.

Vector Subspace: Examples

- Consider $P^n = (\mathbb{P}^n, +, \cdot)$ - a vector space.
- $\forall m \leq n \ P^m = (\mathbb{P}^m, +, \cdot) \subseteq P^n$ - a vector subspace:

Vector Subspace: Examples

- Consider $P^n = (\mathbb{P}^n, +, \cdot)$ - a vector space.
- $\forall m \leq n \ P^m = (\mathbb{P}^m, +, \cdot) \subseteq P^n$ - a vector subspace:
 - $0 \in \mathbb{P}^m$

Vector Subspace: Examples

- Consider $P^n = (\mathbb{P}^n, +, \cdot)$ - a vector space.
- $\forall m \leq n \ P^m = (\mathbb{P}^m, +, \cdot) \subseteq P^n$ - a vector subspace:
 - $0 \in \mathbb{P}^m$
 - Closure: when we add up polynomials of degree $m \leq n$ or multiply them by a scalar, we always get a polynomial of degree $m \leq n$.

Linear Combinations



Linear Combinations

- We can add vectors together and multiply them by scalars.
Let's combine these two operations!

Linear Combinations

- We can add vectors together and multiply them by scalars.
Let's combine these two operations!
- V – a vector space.

Linear Combinations

- We can add vectors together and multiply them by scalars. Let's combine these two operations!
- V – a vector space.
- $x_1, x_2, \dots, x_k \in V$ – some vectors.

Linear Combinations

- We can add vectors together and multiply them by scalars. Let's combine these two operations!
- V – a vector space.
- $x_1, x_2, \dots, x_k \in V$ – some vectors.
- $\lambda_1, \lambda_2, \dots, \lambda_k \in \mathbb{R}$ - some scalars.

Linear Combinations

- We can add vectors together and multiply them by scalars. Let's combine these two operations!
- V – a vector space.
- $x_1, x_2, \dots, x_k \in V$ – some vectors.
- $\lambda_1, \lambda_2, \dots, \lambda_k \in \mathbb{R}$ – some scalars.

$$v = \lambda_1 x_1 + \lambda_2 x_2 + \dots + \lambda_k x_k = \sum_{i=1}^k \lambda_i x_i \in V -$$

a *linear combination* of x_1, x_2, \dots, x_k .

Linear Combinations: Examples

- In $(\mathbb{R}^2, +, \cdot)$, consider vectors $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

Linear Combinations: Examples

- In $(\mathbb{R}^2, +, \cdot)$, consider vectors $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

$u = 2e_1 - e_2 = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ is a linear combination of e_1 and e_2 .

Linear Combinations: Examples

- In $(\mathbb{R}^2, +, \cdot)$, consider vectors $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

$u = 2e_1 - e_2 = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ is a linear combination of e_1 and e_2 .

- In $(\mathbb{P}^2, +, \cdot)$, consider vectors $e_0 = 1, e_1 = t, e_2 = t^2$.

Linear Combinations: Examples

- In $(\mathbb{R}^2, +, \cdot)$, consider vectors $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

$u = 2e_1 - e_2 = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ is a linear combination of e_1 and e_2 .

- In $(\mathbb{P}^2, +, \cdot)$, consider vectors $e_0 = 1, e_1 = t, e_2 = t^2$.

$u = 2t^2 - t + 1 = 2e_2 - e_1 + e_0$ is a linear combination of e_0, e_1 and e_2 .

Linear Combinations: Examples

- In $(\mathbb{R}^2, +, \cdot)$, consider vectors $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

$u = 2e_1 - e_2 = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ is a linear combination of e_1 and e_2 .

- In $(\mathbb{P}^2, +, \cdot)$, consider vectors $e_0 = 1, e_1 = t, e_2 = t^2$.

$u = 2t^2 - t + 1 = 2e_2 - e_1 + e_0$ is a linear combination of e_0, e_1 and e_2 .

$v = 3t + 3 = 3e_1 + 3e_0$ is a linear combination of e_0 and e_1 .

Span



Span

- V – a vector space, $\mathbb{A} = \{x_1, x_2, \dots, x_k\} \subseteq V$ – a set of vectors.

Span

- V – a vector space, $\mathbb{A} = \{x_1, x_2, \dots, x_k\} \subseteq V$ – a set of vectors.
- A set of all linear combinations of vectors in \mathbb{A} is called a *span* of \mathbb{A} :

Span

- V – a vector space, $\mathbb{A} = \{x_1, x_2, \dots, x_k\} \subseteq V$ – a set of vectors.
- A set of all linear combinations of vectors in \mathbb{A} is called a *span* of \mathbb{A} :

$$\text{span}[x_1, \dots, x_n] = \{\lambda_1 x_1 + \dots + \lambda_n x_n \mid \lambda_1, \dots, \lambda_n \in \mathbb{R}\}$$

Span

- V – a vector space, $\mathbb{A} = \{x_1, x_2, \dots, x_k\} \subseteq V$ – a set of vectors.
- A set of all linear combinations of vectors in \mathbb{A} is called a *span* of \mathbb{A} :

$$\text{span}[x_1, \dots, x_n] = \{\lambda_1 x_1 + \dots + \lambda_n x_n \mid \lambda_1, \dots, \lambda_n \in \mathbb{R}\}$$


- If \mathbb{A} spans vector space V , we write

$$V = \text{span}[\mathbb{A}] \text{ or } V = \text{span}[x_1, \dots, x_n].$$

Span: Example 1




- Consider \mathbb{R}^2 .



Span: Example 1




- Consider \mathbb{R}^2 .
- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$



Span: Example 1




- Consider \mathbb{R}^2 .
- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$
 $\text{span}[e_1] = \left\{ \begin{bmatrix} \lambda \\ 0 \end{bmatrix} \mid \lambda \in \mathbb{R} \right\}$ - only
vectors parallel to the first axis.



Span: Example 1




- Consider \mathbb{R}^2 .
- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$
 $\text{span}[e_1] = \left\{ \begin{bmatrix} \lambda \\ 0 \end{bmatrix} \mid \lambda \in \mathbb{R} \right\}$ - only vectors parallel to the first axis.
- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$



Span: Example 1



- Consider \mathbb{R}^2 .
- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$
 $\text{span}[e_1] = \left\{ \begin{bmatrix} \lambda \\ 0 \end{bmatrix} \mid \lambda \in \mathbb{R} \right\}$ - only vectors parallel to the first axis.
- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
 $\text{span}[e_1] = \left\{ \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix} \mid \lambda_{1,2} \in \mathbb{R} \right\} = \mathbb{R}^2$.



Span: Example 2

- Consider $P^3 = (\mathbb{P}^3, +, \cdot)$.

Span: Example 2

- Consider $P^3 = (\mathbb{P}^3, +, \cdot)$.
- $e_0 = 1, e_1 = t, e_2 = t^2$

$$\text{span}[e_0, \dots, e_2] =$$

Span: Example 2

- Consider $P^3 = (\mathbb{P}^3, +, \cdot)$.
- $e_0 = 1, e_1 = t, e_2 = t^2$

$$\text{span}[e_0, \dots, e_2] = \{at^2 + bt + c \mid a, b, c \in \mathbb{R}\} = P^2$$

Span: Example 2

- Consider $P^3 = (\mathbb{P}^3, +, \cdot)$.
- $e_0 = 1, e_1 = t, e_2 = t^2$

$$\text{span}[e_0, \dots, e_2] = \{at^2 + bt + c \mid a, b, c \in \mathbb{R}\} = P^2 \subseteq P^3$$

Span: Example 2

- Consider $P^3 = (\mathbb{P}^3, +, \cdot)$.
- $e_0 = 1, e_1 = t, e_2 = t^2$

$$\text{span}[e_0, \dots, e_2] = \{at^2 + bt + c \mid a, b, c \in \mathbb{R}\} = P^2 \subseteq P^3$$

- $e_0 = 1, e_1 = t, e_2 = t^2, e_3 = t^3$

$$\text{span}[e_0, \dots, e_3] =$$

Span: Example 2

- Consider $P^3 = (\mathbb{P}^3, +, \cdot)$.
- $e_0 = 1, e_1 = t, e_2 = t^2$

$$\text{span}[e_0, \dots, e_2] = \{at^2 + bt + c \mid a, b, c \in \mathbb{R}\} = P^2 \subseteq P^3$$

- $e_0 = 1, e_1 = t, e_2 = t^2, e_3 = t^3$

$$\text{span}[e_0, \dots, e_3] = \{at^3 + bt^2 + ct + d \mid a, b, c, d \in \mathbb{R}\} = P^3$$

Generating Set

- V – a vector space.
- $\mathbb{A} = \{x_1, x_2, \dots, x_k\} \subseteq V$ – a set of vectors.

Generating Set

- V – a vector space.
- $\mathbb{A} = \{x_1, x_2, \dots, x_k\} \subseteq V$ – a set of vectors.
- If every vector $v \in V$ can be expressed as a linear combination of x_1, x_2, \dots, x_k , \mathbb{A} is called a *generating set* for V .

Linear independence



Linear Combinations

- A zero vector can always be represented as a trivial linear combination of x_1, x_2, \dots, x_k :

$$0 = \sum_{i=1}^k 0 \cdot x_i$$

Linear Combinations

- A zero vector can always be represented as a trivial linear combination of x_1, x_2, \dots, x_k :

$$0 = \sum_{i=1}^k 0 \cdot x_i$$

- We are mostly interested in *non-trivial linear combinations* of x_1, x_2, \dots, x_k where not all λ_i are 0.

Linear (In)dependence

- Consider a vector space V .
- $x_1, x_2, \dots, x_k \in V$ – some vectors.

Linear (In)dependence

- Consider a vector space V .
- $x_1, x_2, \dots, x_k \in V$ – some vectors.
- If there is a non-trivial linear combination of x_1, x_2, \dots, x_k such that $\sum_{i=1}^k \lambda_i x_i = 0$ with at least one $\lambda_i \neq 0$, vectors x_1, x_2, \dots, x_k are *linearly dependent*.

Linear (In)dependence

- Consider a vector space V .
- $x_1, x_2, \dots, x_k \in V$ – some vectors.
- If there is a non-trivial linear combination of x_1, x_2, \dots, x_k such that $\sum_{i=1}^k \lambda_i x_i = 0$ with at least one $\lambda_i \neq 0$, vectors x_1, x_2, \dots, x_k are *linearly dependent*.
- If only trivial solution exists, vectors x_1, x_2, \dots, x_k are *linearly independent*.

Linear (In)dependence

- If there is a non-trivial linear combination of x_1, x_2, \dots, x_k such that $\sum_{i=1}^k \lambda_i x_i = 0$ with at least one $\lambda_i \neq 0$, vectors x_1, x_2, \dots, x_k are *linearly dependent*.

Linear (In)dependence

- If there is a non-trivial linear combination of x_1, x_2, \dots, x_k such that $\sum_{i=1}^k \lambda_i x_i = 0$ with at least one $\lambda_i \neq 0$, vectors x_1, x_2, \dots, x_k are *linearly dependent*.

\Leftrightarrow

- A set of vectors x_1, x_2, \dots, x_k is linearly dependent if and only if (at least) one of the vectors is a linear combination of the others

$$x_i = \alpha_1 x_1 + \dots + \alpha_k x_k$$

Linear (In)dependence

- If there is a non-trivial linear combination of x_1, x_2, \dots, x_k such that $\sum_{i=1}^k \lambda_i x_i = 0$ with at least one $\lambda_i \neq 0$, vectors x_1, x_2, \dots, x_k are *linearly dependent*.

\Leftrightarrow

- A set of vectors x_1, x_2, \dots, x_k is linearly dependent if and only if (at least) one of the vectors is a linear combination of the others

$$x_i = \alpha_1 x_1 + \dots + \alpha_k x_k$$

Why is this so? Try to prove this yourself.

Linear (In)dependence: Example 1

- Consider \mathbb{R}^2 .

Linear (In)dependence: Example 1

- Consider \mathbb{R}^2 .
- Vectors $u = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$ and $v = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$ are not linearly independent: $u = -2v$.

Linear (In)dependence: Example 1

- Consider \mathbb{R}^2 .
- Vectors $u = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$ and $v = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$ are not linearly independent: $u = -2v$.
- Vectors $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ are linearly independent:
there are no $\lambda_1, \lambda_2 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that
$$\lambda_1 e_1 + \lambda_2 e_2 = \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

Linear (In)dependence: Example 1

- Consider \mathbb{R}^2 .
- Vectors $u = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$ and $v = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$ are not linearly independent: $u = -2v$.
- Vectors $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ are linearly independent:
there are no $\lambda_1, \lambda_2 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that
$$\lambda_1 e_1 + \lambda_2 e_2 = \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

(Or: you cannot represent e_1 as λe_2 or vice versa).

Linear (In)dependence: Example 2

- Consider $P = (\mathbb{P}^3, +, \cdot)$.
- $1, t, t^2 \in P$ – vectors. Are they linearly independent?

Linear (In)dependence: Example 2

- Consider $P = (\mathbb{P}^3, +, \cdot)$.
- $1, t, t^2 \in P$ – vectors. Are they linearly independent?
- Yes!

There is no way we can represent one of those vectors as a linear combination of the others.

Linear (In)dependence: Example 3

- Are the following vectors in \mathbb{R}^4 linearly independent?

$$x_1 = \begin{bmatrix} 1 \\ 2 \\ -3 \\ 4 \end{bmatrix}, \quad x_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 2 \end{bmatrix}, \quad x_3 = \begin{bmatrix} -1 \\ -2 \\ 1 \\ 1 \end{bmatrix}$$

Linear (In)dependence: Example 3

- Are the following vectors in \mathbb{R}^4 linearly independent?

$$x_1 = \begin{bmatrix} 1 \\ 2 \\ -3 \\ 4 \end{bmatrix}, \quad x_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 2 \end{bmatrix}, \quad x_3 = \begin{bmatrix} -1 \\ -2 \\ 1 \\ 1 \end{bmatrix}$$

- Are there $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that

$$\lambda_1 x_1 + \lambda_2 x_2 + \lambda_3 x_3 = 0$$

Linear (In)dependence: Example 3

- Are the following vectors in \mathbb{R}^4 linearly independent?

$$x_1 = \begin{bmatrix} 1 \\ 2 \\ -3 \\ 4 \end{bmatrix}, \quad x_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 2 \end{bmatrix}, \quad x_3 = \begin{bmatrix} -1 \\ -2 \\ 1 \\ 1 \end{bmatrix}$$

- Are there $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that

$$\lambda_1 x_1 + \lambda_2 x_2 + \lambda_3 x_3 = 0 \leftrightarrow$$

$$\lambda_1 \begin{bmatrix} 1 \\ 2 \\ -3 \\ 4 \end{bmatrix} + \lambda_2 \begin{bmatrix} 1 \\ 1 \\ 0 \\ 2 \end{bmatrix} + \lambda_3 \begin{bmatrix} -1 \\ -2 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} \lambda_1 + \lambda_2 - \lambda_3 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 \\ -3\lambda_1 + \lambda_3 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

Linear (In)dependence: Example 3

- Are there $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases}$$

Linear (In)dependence: Example 3

- Are there $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} \lambda_3 = 3\lambda_1 \end{cases}$$

Linear (In)dependence: Example 3

- Are there $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} -2\lambda_1 + \lambda_2 = 0 \\ -4\lambda_1 + \lambda_2 = 0 \\ \lambda_3 = 3\lambda_1 \\ 7\lambda_1 + 2\lambda_2 = 0 \end{cases}$$

Linear (In)dependence: Example 3

- Are there $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} -2\lambda_1 + \lambda_2 = 0 \\ -4\lambda_1 + \lambda_2 = 0 \\ \lambda_3 = 3\lambda_1 \\ 7\lambda_1 + 2\lambda_2 = 0 \end{cases} \Leftrightarrow \begin{cases} \lambda_2 = 2\lambda_1 \\ \lambda_3 = 3\lambda_1 \end{cases}$$

Linear (In)dependence: Example 3

- Are there $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} -2\lambda_1 + \lambda_2 = 0 \\ -4\lambda_1 + \lambda_2 = 0 \\ \lambda_3 = 3\lambda_1 \\ 7\lambda_1 + 2\lambda_2 = 0 \end{cases} \Leftrightarrow \begin{cases} \lambda_2 = 2\lambda_1 \\ -2\lambda_1 = 0 \\ \lambda_3 = 3\lambda_1 \\ 11\lambda_1 = 0 \end{cases}$$

Linear (In)dependence: Example 3

- Are there $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} -2\lambda_1 + \lambda_2 = 0 \\ -4\lambda_1 + \lambda_2 = 0 \\ \lambda_3 = 3\lambda_1 \\ 7\lambda_1 + 2\lambda_2 = 0 \end{cases} \Leftrightarrow \begin{cases} \lambda_2 = 2\lambda_1 \\ -2\lambda_1 = 0 \\ \lambda_3 = 3\lambda_1 \\ 11\lambda_1 = 0 \end{cases} \Leftrightarrow \lambda_{1,2,3} = 0$$

Linear (In)dependence: Example 3

- Are there $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} -2\lambda_1 + \lambda_2 = 0 \\ -4\lambda_1 + \lambda_2 = 0 \\ \lambda_3 = 3\lambda_1 \\ 7\lambda_1 + 2\lambda_2 = 0 \end{cases} \Leftrightarrow \begin{cases} \lambda_2 = 2\lambda_1 \\ -2\lambda_1 = 0 \\ \lambda_3 = 3\lambda_1 \\ 11\lambda_1 = 0 \end{cases} \Leftrightarrow \lambda_{1,2,3} = 0$$

- Only a trivial linear combination of x_1, x_2 and x_3 gives a zero vector.

Linear (In)dependence: Example 3

- Are there $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} -2\lambda_1 + \lambda_2 = 0 \\ -4\lambda_1 + \lambda_2 = 0 \\ \lambda_3 = 3\lambda_1 \\ 7\lambda_1 + 2\lambda_2 = 0 \end{cases} \Leftrightarrow \begin{cases} \lambda_2 = 2\lambda_1 \\ -2\lambda_1 = 0 \\ \lambda_3 = 3\lambda_1 \\ 11\lambda_1 = 0 \end{cases} \Leftrightarrow \lambda_{1,2,3} = 0$$

- Only a trivial linear combination of x_1, x_2 and x_3 gives a zero vector.

Therefore, x_1, x_2 and x_3 are linearly independent.

Linear (In)dependence: Example 3

- Are there $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} -2\lambda_1 + \lambda_2 = 0 \\ -4\lambda_1 + \lambda_2 = 0 \\ \lambda_3 = 3\lambda_1 \\ 7\lambda_1 + 2\lambda_2 = 0 \end{cases} \Leftrightarrow \begin{cases} \lambda_2 = 2\lambda_1 \\ -2\lambda_1 = 0 \\ \lambda_3 = 3\lambda_1 \\ 11\lambda_1 = 0 \end{cases} \Leftrightarrow \lambda_{1,2,3} = 0$$

- Only a trivial linear combination of x_1, x_2 and x_3 gives a zero vector.

Therefore, x_1, x_2 and x_3 are linearly independent.

Next lecture: a better way to solve such systems of equations

Dimension of a Linear Space

- The linear space V is called *n -dimensional* if the maximal number of linearly independent vectors in it is equal to n .

Dimension of a Linear Space

- The linear space V is called *n -dimensional* if the maximal number of linearly independent vectors in it is equal to n .

We denote this as $\dim(V) = n$.

Dimension: Example

- \mathbb{R}^n is a n -dimensional vector space. Why?

Dimension: Example

- \mathbb{R}^n is a n -dimensional vector space. Why?
- Consider n vectors e_1, \dots, e_n :

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \quad \dots, \quad e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$

Dimension: Example

- \mathbb{R}^n is a n -dimensional vector space. Why?
- Consider n vectors e_1, \dots, e_n :

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \quad \dots, \quad e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$

- e_1, \dots, e_n are linearly independent $\rightarrow \dim(\mathbb{R}^n) \geq n$.

Dimension: Example

- Can there be more than n linearly independent vectors in \mathbb{R}^n ?

Dimension: Example

- Can there be more than n linearly independent vectors in \mathbb{R}^n ?

No!

Explanation: next lecture.

Dimension: Another Example

- What is the dimensionality of $P^3 = (\mathbb{P}^3, +, \cdot)$?

Dimension: Another Example

- What is the dimensionality of $P^3 = (\mathbb{P}^3, +, \cdot)$?
- We can find 4 linearly independent vectors in P^3 :

$$1, \quad t, \quad t^2, \quad t^3$$

Dimension: Another Example

- What is the dimensionality of $P^3 = (\mathbb{P}^3, +, \cdot)$?
- We can find 4 linearly independent vectors in P^3 :

$$1, \quad t, \quad t^2, \quad t^3$$

- All the other vectors (polynomials) from P^3 are linear combinations of those.

Dimension: Another Example

- What is the dimensionality of $P^3 = (\mathbb{P}^3, +, \cdot)$?
- We can find 4 linearly independent vectors in P^3 :

$$1, \quad t, \quad t^2, \quad t^3$$

- All the other vectors (polynomials) from P^3 are linear combinations of those.

$$\rightarrow \dim(P^3) = 4.$$

Basis



Basis

- A set of n linearly independent vectors e_1, e_2, \dots, e_n in an n -dimensional space V is called a *basis* for V .

Basis

- A set of n linearly independent vectors e_1, e_2, \dots, e_n in an n -dimensional space V is called a *basis* for V .
- Basis is A set of vectors with which we can represent every vector in the vector space by adding them together and scaling them.

Basis: Example

Basis: Example

- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is a basis for \mathbb{R}^2 .

Basis: Example

- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is a basis for \mathbb{R}^2 .
- $e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \dots, e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$ is a basis for \mathbb{R}^n .

Basis: Example

- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is a basis for \mathbb{R}^2 .
- $e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \dots, e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$ is a basis for \mathbb{R}^n .
- $e_0 = 1, e_1 = t, \dots, e_n = t^n$ is a basis for P^n .

Basis: Example

- Find the basis of a vector space spanned by vectors

$$x = \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ -3 \\ 3 \end{bmatrix}, \quad z = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

Basis: Example

- Find the basis of a vector space spanned by vectors

$$x = \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ -3 \\ 3 \end{bmatrix}, \quad z = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

- Vectors x and y are linearly independent.

Basis: Example

- Find the basis of a vector space spanned by vectors

$$x = \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ -3 \\ 3 \end{bmatrix}, \quad z = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

- Vectors x and y are linearly independent.
- Vector z is a linear combination of x and y : $z = x - y$.

Basis: Example

- Find the basis of a vector space spanned by vectors

$$x = \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ -3 \\ 3 \end{bmatrix}, \quad z = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

- Vectors x and y are linearly independent.
- Vector z is a linear combination of x and y : $z = x - y$.
- Therefore, $V = \text{span}[x, y, z] = \text{span}[x, y]$.
 $B = \{x, y\}$ - basis of V .

Coordinates

- Let's fix the order of the vectors in the basis:

$$e_1, e_2, \dots, e_n$$

Coordinates

- Let's fix the order of the vectors in the basis:

$$e_1, e_2, \dots, e_n$$

- Every vector v in a vector space can be represented as a linear combination of e_1, e_2, \dots, e_n :

$$v = a_1e_1 + a_2e_2 + \dots + a_ne_n$$

Coordinates

- Let's fix the order of the vectors in the basis:

$$e_1, e_2, \dots, e_n$$

- Every vector v in a vector space can be represented as a linear combination of e_1, e_2, \dots, e_n :


$$v = a_1e_1 + a_2e_2 + \dots + a_ne_n$$

- a_1, a_2, \dots, a_n - *coordinates* of the vector v in the basis e_1, e_2, \dots, e_n .

Coordinates: Example




- Consider \mathbb{R}^2 .



Coordinates: Example



- Consider \mathbb{R}^2 .
- $x = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$



Coordinates: Example




- Consider \mathbb{R}^2 .

- $x = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$

- Canonical basis:

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



Coordinates: Example




- Consider \mathbb{R}^2 .

- $x = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$

- Canonical basis:

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$x = 4e_1 + 6e_2$$



Orthogonal Basis

- A basis e_1, \dots, e_n where $\forall i, j \ e_i \perp e_j$ is an *orthogonal basis*.

Orthogonal Basis

- A basis e_1, \dots, e_n where $\forall i, j \ e_i \perp e_j$ is an *orthogonal basis*.
- If, additionally, $\forall i \ \|e_i\| = 1$, then e_1, \dots, e_n is an *orthonormal basis*.

Orthogonal Basis

- A basis e_1, \dots, e_n where $\forall i, j \ e_i \perp e_j$ is an *orthogonal basis*.
- If, additionally, $\forall i \ \|e_i\| = 1$, then e_1, \dots, e_n is an *orthonormal basis*.

Example:

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \ e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \ \dots, \ e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix} \text{ is an orthonormal basis of } \mathbb{R}^n.$$

Orthogonal Basis

- A basis e_1, \dots, e_n where $\forall i, j \ e_i \perp e_j$ is an *orthogonal basis*.
- If, additionally, $\forall i \ \|e_i\| = 1$, then e_1, \dots, e_n is an *orthonormal basis*.

Example:

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \ e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \ \dots, \ e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix} \text{ is an orthonormal basis of } \mathbb{R}^n.$$

- *Gram-Schmidt process*: a way to convert any basis to an orthogonal one. More details: practical session.

Change of Basis



Basis

- A vector space has more than one basis.

Basis

- A vector space has more than one basis.
- Example: \mathbb{R}^2
 - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ - canonical basis;

Basis

- A vector space has more than one basis.
- Example: \mathbb{R}^2
 - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ - canonical basis;
 - $a = \left\{ a_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, a_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$ - another basis;

Basis

- A vector space has more than one basis.
- Example: \mathbb{R}^2
 - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ - canonical basis;
 - $a = \left\{ a_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, a_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$ - another basis;
 - $b = \left\{ b_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, b_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix} \right\}$ - yet another one.

Basis

- A vector space has more than one basis.
- Example: \mathbb{R}^2
 - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ - canonical basis;
 - $a = \left\{ a_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, a_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$ - another basis;
 - $b = \left\{ b_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, b_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix} \right\}$ - yet another one.
- Different basis = different coordinates.
How exactly do they change?

Coordinate Change: Example

- Consider \mathbb{R}^2 with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Coordinate Change: Example

- Consider \mathbb{R}^2 with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- New basis:

$$e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

Coordinate Change: Example

- Consider \mathbb{R}^2 with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- New basis:

$$e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad e'_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- $x_{old} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$

Coordinate Change: Example

- Consider \mathbb{R}^2 with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- New basis:

$$e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

- $x_{old} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$

- What are the coordinates in the new basis?

$$x_{new} = ?$$

Coordinate Change: General Case

- Consider a vector space V with basis e_1, e_2, \dots, e_n .

Coordinate Change: General Case

- Consider a vector space V with basis e_1, e_2, \dots, e_n .
- Imagine vector $x = [x_1, x_2, \dots, x_n] \in V$
 x_1, x_2, \dots, x_n - coordinates in basis e_1, e_2, \dots, e_n .

Coordinate Change: General Case

- Consider a vector space V with basis e_1, e_2, \dots, e_n .
- Imagine vector $x = [x_1, x_2, \dots, x_n] \in V$
 x_1, x_2, \dots, x_n - coordinates in basis e_1, e_2, \dots, e_n .
- New basis: e'_1, e'_2, \dots, e'_n .

Coordinate Change: General Case

- Consider a vector space V with basis e_1, e_2, \dots, e_n .
- Imagine vector $x = [x_1, x_2, \dots, x_n] \in V$
 x_1, x_2, \dots, x_n - coordinates in basis e_1, e_2, \dots, e_n .
- New basis: e'_1, e'_2, \dots, e'_n .
- What are the coordinates of x in this new basis?
$$x'_1, x'_2, \dots, x'_n = ?$$

Coordinate Change

- Old basis: e_1, e_2, \dots, e_n
New basis: e'_1, e'_2, \dots, e'_n
- $x_{old} = [x_1, x_2, \dots, x_n]$, $x_{new} = [x'_1, x'_2, \dots, x'_n] = ?$
- Coordinates of the new basis in the old one:

Coordinate Change

- Old basis: e_1, e_2, \dots, e_n
New basis: e'_1, e'_2, \dots, e'_n
- $x_{old} = [x_1, x_2, \dots, x_n]$, $x_{new} = [x'_1, x'_2, \dots, x'_n] = ?$
- Coordinates of the new basis in the old one:

$$e'_1 = \alpha_{11}e_1 + \alpha_{21}e_2 + \dots + \alpha_{n1}e_n$$

$$e'_2 = \alpha_{12}e_1 + \alpha_{22}e_2 + \dots + \alpha_{n2}e_n$$

$$\vdots$$

$$e'_i = \alpha_{1i}e_1 + \alpha_{2i}e_2 + \dots + \alpha_{ni}e_n$$

$$\vdots$$

$$e'_n = \alpha_{1n}e_1 + \alpha_{2n}e_2 + \dots + \alpha_{nn}e_n$$

Coordinate Change

$$x = x_1e_1 + x_2e_2 + \cdots + x_ne_n = x'_1\mathbf{e}'_1 + x'_2\mathbf{e}'_2 + \cdots x'_n\mathbf{e}'_n =$$

Coordinate Change

$$x = x_1e_1 + x_2e_2 + \cdots + x_ne_n = x'_1\mathbf{e}'_1 + x'_2\mathbf{e}'_2 + \cdots x'_n\mathbf{e}'_n =$$

Remember: $e'_i = \alpha_{1i}e_1 + \alpha_{2i}e_2 + \cdots + \alpha_{ni}e_n$

Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember: $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} = x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) = \end{aligned}$$

Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember: $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} &= x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ &\quad + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) = \end{aligned}$$

e_1, \dots, e_n linearly independent \rightarrow coefficients in front of them
should be the same on the both sides of the equality:

Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember: $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} &= x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ &\quad + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) = \end{aligned}$$

e_1, \dots, e_n linearly independent \rightarrow coefficients in front of them should be the same on the both sides of the equality:

$$x_1 = x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n}$$

$$x_2 = x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n}$$

$$\vdots$$

$$x_n = x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn}$$

Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember: $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} &= x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ &\quad + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) = \end{aligned}$$

x_{old}

$$\begin{aligned} x_1 &= x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n} \\ x_2 &= x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n} \\ &\vdots \\ x_n &= x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn} \end{aligned}$$

Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember: $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$= x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) =$$

$$\begin{array}{l} x_1 = x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n} \\ x_2 = x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n} \\ \vdots \\ x_n = x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn} \end{array}$$

x_{old}

x_{new}

Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

$$\text{Remember: } e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$$

$$= x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) =$$

$$\begin{array}{l} x_1 = x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n} \\ x_2 = x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n} \\ \vdots \\ x_n = x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn} \end{array}$$

x_{new}

Coordinate Change: Example

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

Coordinate Change: Example

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

$$\begin{matrix} & & & & e'_i \\ x_{old} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} & = & \begin{bmatrix} x'_1 \\ x'_1 \\ \vdots \\ x'_1 \end{bmatrix} \begin{bmatrix} \alpha_{11} \\ \alpha_{21} \\ \vdots \\ \alpha_{n1} \end{bmatrix} & + \cdots & + \begin{bmatrix} x'_i \\ x'_i \\ \vdots \\ x'_i \end{bmatrix} \begin{bmatrix} \alpha_{1i} \\ \alpha_{2i} \\ \vdots \\ \alpha_{ni} \end{bmatrix} & + \cdots & + \begin{bmatrix} x'_n \\ x'_n \\ \vdots \\ x'_n \end{bmatrix} \begin{bmatrix} \alpha_{1n} \\ \alpha_{2n} \\ \vdots \\ \alpha_{nn} \end{bmatrix} \end{matrix}$$

x_{new}

Coordinate Change: Example

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

$$\begin{matrix} x_{old} \\ \begin{matrix} x_1 = x'_1\alpha_{11} + \dots + x'_i\alpha_{1i} + \dots + x'_n\alpha_{1n} \\ x_2 = x'_1\alpha_{21} + \dots + x'_i\alpha_{2i} + \dots + x'_n\alpha_{2n} \\ \vdots \\ x_n = x'_1\alpha_{n1} + \dots + x'_i\alpha_{ni} + \dots + x'_n\alpha_{nn} \end{matrix} \end{matrix}$$

x_{new}

$$\begin{aligned} 2 &= 2x'_1 - 1x'_2 \\ -1 &= 1x'_1 - 1x'_2 \end{aligned}$$

Coordinate Change: Example

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

$$\begin{array}{l} x_1 = x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n} \\ x_2 = x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n} \\ \vdots \\ x_n = x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn} \end{array}$$

x_{old} (red box), x_{new} (green boxes), e'_i (blue line)

$$\begin{array}{l} 2 = 2x'_1 - 1x'_2 \\ -1 = 1x'_1 - 1x'_2 \end{array} \Leftrightarrow \begin{array}{l} x'_1 = 3 \\ x'_2 = 4 \end{array}$$

Coordinate Change: Example

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

$$\begin{array}{l} x_1 = x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n} \\ x_2 = x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n} \\ \vdots \\ x_n = x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn} \end{array}$$

x_{old} x_{new} e'_i

$$\begin{array}{l} 2 = 2x'_1 - 1x'_2 \\ -1 = 1x'_1 - 1x'_2 \end{array} \Leftrightarrow \begin{array}{l} x'_1 = 3 \\ x'_2 = 4 \end{array} \Leftrightarrow x_{new} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$$

Coordinate Change

- Going from one basis to the other:

The diagram shows the transformation of a vector x_{old} from an old basis to a new basis x_{new} . The vector x_{old} is represented by a red box containing the components x_1, x_2, \dots, x_n . The new basis vectors e'_i are shown at the top, with lines connecting them to the corresponding terms in the expansion of each x_i . The expansion of x_i is shown as a sum of products of the new basis components x'_j and the change of basis coefficients α_{ji} . The new basis components x'_j are highlighted in green boxes, and the coefficients α_{ji} are in blue boxes. The entire expansion is enclosed in a green box labeled x_{new} .

$$\begin{aligned} x_1 &= x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n} \\ x_2 &= x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n} \\ &\vdots \\ x_n &= x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn} \end{aligned}$$

Coordinate Change

- Going from one basis to the other:

$$\begin{array}{l} x_1 = x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n} \\ x_2 = x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n} \\ \vdots \\ x_n = x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn} \end{array}$$

x_{old}

x_{new}

e'_i

- There is a more compact way of writing this down using [matrices](#).

Matrices



A Matrix

- $A \in \mathbb{R}^{m \times n}$ - a matrix with m rows and n columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

A Matrix

- $A \in \mathbb{R}^{m \times n}$ - a matrix with m rows and n columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- *Examples:*

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}, \quad \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \in \mathbb{R}^{2 \times 3}$$

Special Matrices

- Diagonal matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ ($a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j$)

Special Matrices

- Diagonal matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ $(a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j)$
- Identity matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ $(a_{ii} = 1, a_{ij} = 0 \forall i \neq j)$

Special Matrices

- Diagonal matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ ($a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j$)
- Identity matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ ($a_{ii} = 1, a_{ij} = 0 \forall i \neq j$)
- Symmetric matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ ($a_{ij} = a_{ji}$)

Special Matrices

- Diagonal matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ ($a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j$)
- Identity matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ ($a_{ii} = 1, a_{ij} = 0 \forall i \neq j$)
- Symmetric matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ ($a_{ij} = a_{ji}$)
- Triangular matrix: $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 2 & 3 & 0 \\ 4 & 5 & 6 \end{bmatrix}$ ($a_{ij} = 0 \forall i > j \text{ or } \forall i < j$)

Vectors vs Matrices

- An n -dimensional vector can be considered a $n \times 1$ matrix:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^{n \times 1}$$

Operations with Matrices



Transpose of a Matrix

- Consider a matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

Transpose of a Matrix

- Consider a matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- Transpose = writing columns as rows:

$$A^T = \begin{bmatrix} a_{11} & a_{21} & \cdots & a_{m1} \\ a_{12} & a_{22} & \cdots & a_{m2} \\ \cdots & \cdots & \cdots & \cdots \\ a_{1n} & a_{2n} & \cdots & a_{mn} \end{bmatrix}, \quad \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}^T = [x_1, \dots, x_n]$$

Transpose of a Matrix: Example

- $\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix}^T = \begin{bmatrix} 0 & 2 \\ 1 & 1 \\ 2 & 5 \end{bmatrix}$

Transpose of a Matrix: Example

- $\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix}^T = \begin{bmatrix} 0 & 2 \\ 1 & 1 \\ 2 & 5 \end{bmatrix}$
- Transposing a symmetrical matrix = no changes:

$$\begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 2 \\ 3 & 2 & 1 \end{bmatrix}^T = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 2 \\ 3 & 2 & 1 \end{bmatrix}$$

Multiplying by a Scalar

- We can multiply matrix by a scalar:

$$\lambda A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} \lambda a_{11} & \lambda a_{12} & \cdots & \lambda a_{1n} \\ \lambda a_{21} & \lambda a_{22} & \cdots & \lambda a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ \lambda a_{m1} & \lambda a_{m2} & \cdots & \lambda a_{mn} \end{bmatrix}$$

Multiplying by a Scalar

- We can multiply matrix by a scalar:

$$\lambda A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} \lambda a_{11} & \lambda a_{12} & \cdots & \lambda a_{1n} \\ \lambda a_{21} & \lambda a_{22} & \cdots & \lambda a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ \lambda a_{m1} & \lambda a_{m2} & \cdots & \lambda a_{mn} \end{bmatrix}$$

- Example:

$$5 \cdot \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}$$

Sum of Two Matrices

- We can sum up matrices of the same size:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}, \quad B = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ b_{21} & b_{22} & \cdots & b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ b_{m1} & b_{m2} & \cdots & b_{mn} \end{bmatrix},$$

$$A + B = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \cdots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & \cdots & a_{2n} + b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \cdots & a_{mn} + b_{mn} \end{bmatrix}$$

Sum of Two Matrices

- We can sum up matrices of the same size:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}, \quad B = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ b_{21} & b_{22} & \cdots & b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ b_{m1} & b_{m2} & \cdots & b_{mn} \end{bmatrix},$$

$$A + B = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \cdots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & \cdots & a_{2n} + b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \cdots & a_{mn} + b_{mn} \end{bmatrix}$$

- Example:

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} -1 & 2 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 2 \\ 0 & 4 \end{bmatrix}$$

Matrices Also Form a Vector Space!

- $(\mathbb{R}^{m \times n}, +, \cdot)$ - a vector space.
“Vectors” = matrices.

Matrices Also Form a Vector Space!

- $(\mathbb{R}^{m \times n}, +, \cdot)$ - a vector space.
“Vectors” = matrices.
- You can check yourself that the necessary axioms hold.

Matrix Multiplication

- Consider two matrices $A = \{a_{ij}\}_{m \times n}$ and $b = \{b_{ij}\}_{n \times p}$.
- $C = AB$ – product of two matrices.

Matrix Multiplication

- Consider two matrices $A = \{a_{ij}\}_{m \times n}$ and $b = \{b_{ij}\}_{n \times p}$.
- $C = AB$ – product of two matrices.
- Each element c_{ij} of C is a dot product of i -th row of A and j -th column of B : $C = \{c_{ij}\}_{m \times p} = \{(A_i, B^j)\}_{m \times p}$

Matrix Multiplication

- Consider two matrices $A = \{a_{ij}\}_{m \times n}$ and $b = \{b_{ij}\}_{n \times p}$.
- $C = AB$ – product of two matrices.
- Each element c_{ij} of C is a dot product of i -th row of A and j -th column of B : $C = \{c_{ij}\}_{m \times p} = \{(A_i, B^j)\}_{m \times p}$
- Example $\mathbb{R}^{2 \times 2}$:
$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \cdot \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} = \begin{bmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{bmatrix}$$

Matrix Multiplication: Example

$$\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 3 & 6 & 4 \\ 2 & 5 & 8 \\ 7 & 1 & 9 \end{bmatrix} =$$

Matrix Multiplication: Example

$$\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 3 & 6 & 4 \\ 2 & 5 & 8 \\ 7 & 1 & 9 \end{bmatrix} =$$

$$= \begin{bmatrix} 0 + 2 + 14 & 0 + 5 + 2 & 0 + 8 + 18 \\ 6 + 2 + 35 & 12 + 5 + 5 & 8 + 8 + 45 \end{bmatrix} =$$

Matrix Multiplication: Example

$$\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 3 & 6 & 4 \\ 2 & 5 & 8 \\ 7 & 1 & 9 \end{bmatrix} =$$

$$= \begin{bmatrix} 0 + 2 + 14 & 0 + 5 + 2 & 0 + 8 + 18 \\ 6 + 2 + 35 & 12 + 5 + 5 & 8 + 8 + 45 \end{bmatrix} =$$

$$= \begin{bmatrix} 16 & 7 & 26 \\ 43 & 22 & 61 \end{bmatrix}$$

Coordinate Change: Matrix Notation



- Result obtained before:

e_1, \dots, e_n - old basis

e'_1, \dots, e'_n - new basis

$$x_{old} = [x_1, \dots, x_n], \quad x_{new} = [x'_1, \dots, x'_n]$$

x_{old}

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} x'_1 \alpha_{11} + \dots + x'_i \alpha_{i1} + \dots + x'_n \alpha_{1n} \\ x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n} \\ \vdots \\ x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn} \end{bmatrix}$$

e'_i

x_{new}

Coordinate Change: Matrix Notation



- Result obtained before:

e_1, \dots, e_n - old basis

e'_1, \dots, e'_n - new basis

$$x_{old} = [x_1, \dots, x_n], \quad x_{new} = [x'_1, \dots, x'_n]$$

- Transition matrix: columns = coordinates of the new basis in the old one.

$$A = \begin{bmatrix} \alpha_{11} & \alpha_{12} & \dots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} & \dots & \alpha_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{n1} & \alpha_{n2} & \dots & \alpha_{nn} \end{bmatrix}$$

x_{old}

$$\begin{array}{l} x_1 = x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n} \\ x_2 = x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n} \\ \vdots \\ x_n = x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn} \end{array}$$

x_{new}

e'_i

Coordinate Change: Matrix Notation



- Result obtained before:

e_1, \dots, e_n - old basis

e'_1, \dots, e'_n - new basis

$$x_{old} = [x_1, \dots, x_n], \quad x_{new} = [x'_1, \dots, x'_n]$$

x_{old}

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} x'_1 & x'_2 & \dots & x'_i & \dots & x'_n \end{bmatrix} \begin{bmatrix} \alpha_{11} & \alpha_{12} & \dots & \alpha_{1i} & \dots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} & \dots & \alpha_{2i} & \dots & \alpha_{2n} \\ \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ \alpha_{n1} & \alpha_{n2} & \dots & \alpha_{ni} & \dots & \alpha_{nn} \end{bmatrix}$$

x_{new}

e'_i

- Transition matrix: columns = coordinates of the new basis in the old one.

$$A = \begin{bmatrix} \alpha_{11} & \alpha_{12} & \dots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} & \dots & \alpha_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{n1} & \alpha_{n2} & \dots & \alpha_{nn} \end{bmatrix}$$

$$x_{old} = Ax_{new}$$

Coordinate Change: Example (again)

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = ?$

Coordinate Change: Example (again)

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = ?$

$$A = \begin{bmatrix} 2 & -1 \\ 1 & -1 \end{bmatrix},$$

Coordinate Change: Example (again)

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = ?$

$$A = \begin{bmatrix} 2 & -1 \\ 1 & -1 \end{bmatrix}, \quad \begin{bmatrix} 2 \\ -1 \end{bmatrix} = x_{old} = Ax_{new} =$$

Coordinate Change: Example (again)

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = ?$

$$A = \begin{bmatrix} 2 & -1 \\ 1 & -1 \end{bmatrix}, \quad \begin{bmatrix} 2 \\ -1 \end{bmatrix} = x_{old} = Ax_{new} = \begin{bmatrix} 2 & -1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} =$$

Coordinate Change: Example (again)

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = ?$

$$A = \begin{bmatrix} 2 & -1 \\ 1 & -1 \end{bmatrix}, \quad \begin{bmatrix} 2 \\ -1 \end{bmatrix} = x_{old} = Ax_{new} = \begin{bmatrix} 2 & -1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} 2x'_1 - x'_2 \\ x'_1 - x'_2 \end{bmatrix}$$

Coordinate Change: Example (again)

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = ?$

$$A = \begin{bmatrix} 2 & -1 \\ 1 & -1 \end{bmatrix}, \quad \begin{bmatrix} 2 \\ -1 \end{bmatrix} = x_{old} = Ax_{new} = \begin{bmatrix} 2 & -1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} 2x'_1 - x'_2 \\ x'_1 - x'_2 \end{bmatrix}$$

$$x_{new} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}.$$

To Sum Up

- Vector spaces
 - Linear (in)dependence
 - Span
 - Basis
- Matrices
 - Matrix operations
 - Change of coordinates

Next Time

- More on matrices
- Systems of linear equations