



Math Refresher for DS

Lecture 3



Last Time

- Vector Spaces
 - Linear combinations
 - Spans
 - Bases
 - Change of coordinates
- Matrices

Today

- More on matrices
 - matrix operations;
 - rank;
 - determinant.
- Linear transformations
- Systems of linear equations

Matrices: a small review



A Matrix

- $A \in \mathbb{R}^{m \times n}$ - a matrix with m rows and n columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- *Examples:*

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}, \quad \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \in \mathbb{R}^{2 \times 3}$$

Special Matrices

- Diagonal matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ ($a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j$)
- Identity matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ ($a_{ii} = 1, a_{ij} = 0 \forall i \neq j$)
- Symmetric matrix: $\begin{bmatrix} 1 & 4 & 5 \\ 4 & 2 & 6 \\ 5 & 6 & 3 \end{bmatrix}$ ($a_{ij} = a_{ji}$)
- Triangular matrix: $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 2 & 3 & 0 \\ 4 & 5 & 6 \end{bmatrix}$ ($a_{ij} = 0 \forall i > j \text{ or } \forall i < j$)

Basic Operations with Matrices

- Addition:

$$A = \{a_{ij}\}_{i=1,\dots,m,j=1,\dots,n'} \quad B = \{b_{ij}\}_{i=1,\dots,m,j=1,\dots,n'} \quad A + B = \{a_{ij} + b_{ij}\}_{i=1,\dots,m,j=1,\dots,n}$$

Basic Operations with Matrices

- Addition:

$$A = \{a_{ij}\}_{i=1,\dots,m,j=1,\dots,n'} \quad B = \{b_{ij}\}_{i=1,\dots,m,j=1,\dots,n'} \quad A + B = \{a_{ij} + b_{ij}\}_{i=1,\dots,m,j=1,\dots,n}$$

- Multiplication by a scalar:

$$A = \{a_{ij}\}_{i=1,\dots,m,j=1,\dots,n'} \quad \lambda \in \mathbb{R}, \quad \lambda A = \{\lambda a_{ij}\}_{i=1,\dots,m,j=1,\dots,n}$$

Matrix Multiplication

- Matrix multiplication:

$$A = \{a_{ij}\}_{i=1,\dots,m,j=1,\dots,n}, \quad B = \{b_{ij}\}_{i=1,\dots,n,j=1,\dots,k}$$

$$A \cdot B = \{(A_i, B^j)\}_{i=1,\dots,m,j=1,\dots,k} = \left\{ \sum_{l=1,\dots,n} a_{il} \cdot b_{lj} \right\}_{i=1,\dots,m,j=1,\dots,k}$$

Matrix Multiplication

- Matrix multiplication:

$$A = \{a_{ij}\}_{i=1,\dots,m,j=1,\dots,n}, \quad B = \{b_{ij}\}_{i=1,\dots,n,j=1,\dots,k}$$

$$A \cdot B = \{(A_i, B^j)\}_{i=1,\dots,m,j=1,\dots,k} = \left\{ \sum_{l=1,\dots,n} a_{il} \cdot b_{lj} \right\}_{i=1,\dots,m,j=1,\dots,k}$$

- Example $\mathbb{R}^{2 \times 2}$:

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \cdot \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} = \begin{bmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{bmatrix}$$

Matrix Multiplication

- For numbers: $2 \times 3 = 3 \times 2 = 6$.

Matrix Multiplication

- For numbers: $2 \times 3 = 3 \times 2 = 6$.
- Matrix multiplication is *(in general)* not commutative:

$$\mathbf{AB} \neq \mathbf{BA}$$

Matrix Multiplication

- For numbers: $2 \times 3 = 3 \times 2 = 6$.
- Matrix multiplication is *(in general)* not commutative:

$$\mathbf{AB} \neq \mathbf{BA}$$

- Example:

$$A = \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} 6 & 2 \\ 3 & 2 \end{bmatrix}, \quad AB = \begin{bmatrix} 30 & 14 \\ 12 & 6 \end{bmatrix}, \quad BA = \begin{bmatrix} 20 & 28 \\ 11 & 16 \end{bmatrix}$$

Matrix Multiplication

- Multiplication by identity matrix E :

$$AE = EA = A$$

Matrix Multiplication

- Multiplication by identity matrix E :

$$AE = EA = A$$

- Multiplication by zero matrix O :

$$AO = OA = O$$

Transposing a Matrix

- The transpose of a matrix results from “flipping” the rows and columns:

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \quad A^T = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}.$$

Transposing a Matrix

- The following properties of transposes are easily verified:
 - A – symmetric matrix $\Rightarrow A^T = A$
 - $(A^T)^T = A$
 - $(A + B)^T = A^T + B^T$
 - $(AB)^T = B^T A^T$

Linear Transforms

*A more interesting way of looking
at matrices.*



Linear Transformation



Linear Transformation

Linear Transformation



Linear Transformation

Linear Transformation




Linear Transformation

$$x_{input} \rightarrow A \rightarrow x_{output}$$


A – transformation

x_{input}, x_{output} – vectors





Transformation



Transformation



Transformation: Examples



Linear Transformation



Linear Transformation

$$x_{input} \rightarrow A \rightarrow x_{output}$$

A – transformation

x_{input}, x_{output} – vectors

Linear Transformation



Linear Transformation

A transformation that satisfies two properties:

1. $A(x + y) = A(x) + A(y)$
2. $A(\lambda x) = \lambda Ax$

$$x_{input} \rightarrow A \rightarrow x_{output}$$

A – transformation

x_{input}, x_{output} – vectors

Linear Transformation

- How to describe a linear transformation numerically?
-

Linear Transformation

- How to describe a linear transformation numerically?
- With matrices! How?

Linear Transformation

- How to describe a linear transformation numerically?
- With matrices! How?

$$x_{input} = x_1 e_1 + x_2 e_2 + \dots + x_n e_n, \quad e_1, \dots, e_n - \text{basis}, \quad x_1, \dots, x_n - \text{coordinates}$$

Linear Transformation

- How to describe a linear transformation numerically?
- With matrices! How?

$x_{input} = x_1e_1 + x_2e_2 + \dots + x_ne_n$, e_1, \dots, e_n – basis, x_1, \dots, x_n – coordinates

$$x_{output} = A(x_{input}) = A(x_1e_1 + x_2e_2 + \dots + x_ne_n) =$$

Linear Transformation

- How to describe a linear transformation numerically?
- With matrices! How?

$x_{input} = x_1e_1 + x_2e_2 + \dots + x_ne_n$, e_1, \dots, e_n – basis, x_1, \dots, x_n – coordinates

$$x_{output} = A(x_{input}) = A(x_1e_1 + x_2e_2 + \dots + x_ne_n) = x_1A(e_1) + x_2A(e_2) + \dots + x_nA(e_n)$$

Linear Transformation

- How to describe a linear transformation numerically?
- With matrices! How?

$x_{input} = x_1e_1 + x_2e_2 + \dots + x_ne_n$, e_1, \dots, e_n – basis, x_1, \dots, x_n – coordinates

$$x_{output} = A(x_{input}) = A(x_1e_1 + x_2e_2 + \dots + x_ne_n) = x_1A(e_1) + x_2A(e_2) + \dots + x_nA(e_n)$$

$$A := [A(e_1) \mid A(e_2) \mid \dots \mid A(e_n)]$$

Linear Transformation

- How to describe a linear transformation numerically?
- With matrices! How?

$x_{input} = x_1e_1 + x_2e_2 + \dots + x_ne_n$, e_1, \dots, e_n – basis, x_1, \dots, x_n – coordinates

$$x_{output} = A(x_{input}) = A(x_1e_1 + x_2e_2 + \dots + x_ne_n) = x_1A(e_1) + x_2A(e_2) + \dots + x_nA(e_n)$$


$$A := [A(e_1) \mid A(e_2) \mid \dots \mid A(e_n)]$$

$$\Rightarrow x_{output} = A(x_{input}) = A \cdot x_{input}$$

Example: Rotation



- Imagine that we want to rotate vectors in \mathbb{R}^2 90° anti-clockwise.




Example: Rotation



- Imagine that we want to rotate vectors in \mathbb{R}^2 90° anti-clockwise.
- What happens to the basis vectors?

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \quad , \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow$$




Example: Rotation



- Imagine that we want to rotate vectors in \mathbb{R}^2 90° anti-clockwise.
- What happens to the basis vectors?

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \quad , \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow$$




Example: Rotation



- Imagine that we want to rotate vectors in \mathbb{R}^2 90° anti-clockwise.
- What happens to the basis vectors?

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$



Example: Rotation




- Imagine that we want to rotate vectors in \mathbb{R}^2 90° anti-clockwise.
- What happens to the basis vectors?

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

- Therefore,

$$R = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$





Example: Rotation


- Imagine that we want to rotate vectors in \mathbb{R}^2 90° anti-clockwise.
- What happens to the basis vectors?

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

- Therefore,

$$R = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

- Consider $x = [1, 1]^T$.



Example: Rotation



- Imagine that we want to rotate vectors in \mathbb{R}^2 90° anti-clockwise.
- What happens to the basis vectors?


$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

- Therefore,

$$R = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

- Consider $x = [1, 1]^T$. After rotation:

$$x_{\text{rotated}} =$$



Example: Rotation



- Imagine that we want to rotate vectors in \mathbb{R}^2 90° anti-clockwise.
- What happens to the basis vectors?


$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

- Therefore,

$$R = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

- Consider $x = [1, 1]^T$. After rotation:

$$x_{rotated} =$$





Example: Rotation

- Imagine that we want to rotate vectors in \mathbb{R}^2 90° anti-clockwise.
- What happens to the basis vectors?


$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

- Therefore,

$$R = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

- Consider $x = [1, 1]^T$. After rotation:

$$x_{rotated} = Rx = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$



Linear Transformation

- Every linear transformation can be defined by its matrix.

Columns = how this transformation changes the vectors in the selected basis.

Linear Transformation

- Every linear transformation can be defined by its matrix.

Columns = how this transformation changes the vectors in the selected basis.

- Vice versa: every square matrix defines some linear transformation.

Common Transforms



Identity Transformation

- Doesn't change anything.
- Transformation matrix E :

$$Ex = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Identity Transformation

- Doesn't change anything.
- Transformation matrix E :

$$Ex = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

- Example:

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

Stretching / Squeezing

- Enlarge (compress) all distances in a particular direction by a constant factor.
- Transformation matrix:

$$Kx = \begin{bmatrix} k_1 & 0 \\ 0 & k_2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Stretching / Squeezing

- Enlarge (compress) all distances in a particular direction by a constant factor.
- Transformation matrix:

$$Kx = \begin{bmatrix} k_1 & 0 \\ 0 & k_2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

- Example: stretch x -axis ($\times 3$) and squeeze y -axis ($\times 0.5$):

$$\begin{bmatrix} 3 & 0 \\ 0 & 0.5 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

Projection on an Axis

- Consider \mathbb{R}^3 . Project on the XY –plane.
- Transformation matrix:

$$Px = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

Projection on an Axis

- Consider \mathbb{R}^3 . Project on the XY –plane.
- Transformation matrix:

$$Px = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

- Example:

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$$

Rotation

- Rotating points anticlockwise by θ .
- Rotation matrix R_θ :

$$R_\theta x = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Rotation

- Rotating points anticlockwise by θ .
- Rotation matrix R_θ :

$$R_\theta x = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

- Example: rotate by 45° anticlockwise:

$$\begin{bmatrix} \sqrt{2}/2 & -\sqrt{2}/2 \\ \sqrt{2}/2 & \sqrt{2}/2 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ \sqrt{2} \end{bmatrix}$$

Combining Transforms



Composition

- Let A and B be two linear transforms.
What if we first apply A and then B ?

Composition

- Let A and B be two linear transforms.
What if we first apply A and then B ?
- Example: first rotate by 90° , then squeeze.

Composition

- Let A and B be two linear transforms.
What if we first apply A and then B ?
- Example: first rotate by 90° , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} - \text{rotation}, \quad B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} - \text{squeezing}$$

Composition

- Let A and B be two linear transforms.
What if we first apply A and then B ?
- Example: first rotate by 90° , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} - \text{rotation}, \quad B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} - \text{squeezing}$$


$$B(Ax) =$$

Composition

- Let A and B be two linear transforms. What if we first apply A and then B ?
- Example: first rotate by 90° , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} - \text{rotation}, \quad B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} - \text{squeezing}$$

$$B(Ax) =$$




Composition

- Let A and B be two linear transforms. What if we first apply A and then B ?
- Example: first rotate by 90° , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} - \text{rotation}, \quad B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} - \text{squeezing}$$

$$B(Ax) =$$




Composition

- Let A and B be two linear transforms.
What if we first apply A and then B ?
- Example: first rotate by 90° , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} - \text{rotation}, \quad B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} - \text{squeezing}$$

$$B(Ax) =$$




Composition

- Let A and B be two linear transforms.
What if we first apply A and then B ?
- Example: first rotate by 90° , then squeeze.

$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ – rotation, $B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix}$ – squeezing

$$B(Ax) = (BA)x = Cx$$




Composition

- Let A and B be two linear transforms.
What if we first apply A and then B ?
- Example: first rotate by 90° , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} - \text{rotation}, \quad B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} - \text{squeezing}$$

$$B(Ax) = (BA)x = Cx$$

$$C = BA = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & -0.5 \\ 0.5 & 0 \end{bmatrix}$$



Composition


- Let A and B be two linear transforms.
What if we first apply A and then B ?
- Example: first rotate by 90° , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} - \text{rotation}, \quad B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} - \text{squeezing}$$

$$B(Ax) = (BA)x = Cx$$

$$C = BA = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & -0.5 \\ 0.5 & 0 \end{bmatrix}$$

= "rotate by 90° and squeeze"



Sum of Transforms

- Let A and B be two linear transforms.

Sum of Transforms

- Let A and B be two linear transforms.
- $C = A + B$ – also a linear transform:

$$C = Ax + Bx$$

Sum of Transforms

- Let A and B be two linear transforms.
- $C = A + B$ – also a linear transform:

$$C = Ax + Bx$$

- Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{ – rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{ – squeezing}$$

Sum of Transforms


- Let A and B be two linear transforms.
- $C = A + B$ – also a linear transform:

$$C = Ax + Bx$$

- Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{ – rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{ – squeezing}$$

$$x = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$



Sum of Transforms


- Let A and B be two linear transforms.
- $C = A + B$ – also a linear transform:

$$C = Ax + Bx$$

- Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{ – rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{ – squeezing}$$

$$x = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$



Sum of Transforms


- Let A and B be two linear transforms.
- $C = A + B$ – also a linear transform:

$$C = Ax + Bx$$

- Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{ – rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{ – squeezing}$$

$$x = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$



Sum of Transforms


- Let A and B be two linear transforms.
- $C = A + B$ – also a linear transform:

$$C = Ax + Bx$$

- Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{ – rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{ – squeezing}$$

$$x = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$



Sum of Transforms


- Let A and B be two linear transforms.
- $C = A + B$ – also a linear transform:

$$C = Ax + Bx$$

- Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{ – rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{ – squeezing}$$

$$x = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad Ax + Bx = \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \begin{bmatrix} 0.5 \\ 0.5 \end{bmatrix} = \begin{bmatrix} -0.5 \\ 1.5 \end{bmatrix}$$



Inverse Transform



Inverse transform

Inverse transform

- Consider a transform A .

Inverse transform

- Consider a transform A .
- Inverse transform A^{-1} “maps everything back to where it was”:

Inverse transform

- Consider a transform A .
- Inverse transform A^{-1} “maps everything back to where it was”:

$$A^{-1}(Ax) = x \Leftrightarrow A^{-1}A = E - \text{identity transformation.}$$

Inverse transform

- Consider a transform A .
- Inverse transform A^{-1} “maps everything back to where it was”:

$$A^{-1}(Ax) = x \Leftrightarrow A^{-1}A = E - \text{identity transformation.}$$

- Not every transform has an inverse!

Inverse transform

- Consider a transform A .
- Inverse transform A^{-1} “maps everything back to where it was”:

$$A^{-1}(Ax) = x \Leftrightarrow A^{-1}A = E - \text{identity transformation.}$$

- Not every transform has an inverse!
 - Rotation: yes (rotate it back)

Inverse transform

- Consider a transform A .
- Inverse transform A^{-1} “maps everything back to where it was”:

$$A^{-1}(Ax) = x \Leftrightarrow A^{-1}A = E - \text{identity transformation.}$$

- Not every transform has an inverse!
 - Rotation: yes (rotate it back)
 - Projection: no

Inverse of a Matrix

- An $n \times n$ matrix A has an inverse if there exists A^{-1} such that

$$AA^{-1} = A^{-1}A = E$$

Inverse of a Matrix

- An $n \times n$ matrix A has an inverse if there exists A^{-1} such that

$$AA^{-1} = A^{-1}A = E$$

- A matrix that doesn't have an inverse is called **singular** or **degenerate**.

Inverse of a Matrix

- An $n \times n$ matrix A has an inverse if there exists A^{-1} such that

$$AA^{-1} = A^{-1}A = E$$

- A matrix that doesn't have an inverse is called **singular** or **degenerate**.
- Which matrices have an inverse?

Determinant



Determinant

- A numerical way to characterize a linear transformation (and its matrix):
 - absolute value = how much area changes;
 - sign = change of orientation.
- More info on the interpretation: see [video](#).

Determinant

- A – linear transform.

Determinant


- A – linear transform.
- The determinant is nonzero $\Leftrightarrow A$ is invertible and the linear map represented by matrix A is an **isomorphism**:

Determinant

- A – linear transform.
- The determinant is nonzero $\Leftrightarrow A$ is invertible and the linear map represented by matrix A is an isomorphism:

- $\det A \neq 0$:

- A :




Determinant

- A – linear transform.
- The determinant is nonzero $\Leftrightarrow A$ is invertible and the linear map represented by matrix A is an **isomorphism**:


- $\det A \neq 0$:

- A :



- $\det A = 0$:

- Several vectors are mapped onto the same vector $\Leftrightarrow A$ maps original vector space onto a lower-dimensional space.



Computing Determinant

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

Computing Determinant

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

- Example:

$$R_{90} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = 0 - (-1) = 1 \Leftrightarrow$$

“there is a transform inverse to rotation by 90° anticlockwise”.

Computing Determinant

$$\begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix}$$

Computing Determinant

$$\begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix}$$

- Example:

$$\begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{vmatrix} = 1 \cdot \begin{vmatrix} 1 & 0 \\ 0 & 0 \end{vmatrix} - 0 + 0 = 0 \Leftrightarrow$$

“there is no transpose inverse to projection onto *XY*-plane”

Computing Determinant

- $A = \{a_{ij}\}$ – $n \times n$ matrix.

Computing Determinant

- $A = \{a_{ij}\}$ – $n \times n$ matrix.
- M_{ij} – its minor $\Leftrightarrow M_{ij}$ is an $(n - 1) \times (n - 1)$ matrix resulting from removing i -th row and j -th column from A .

Computing Determinant

- $A = \{a_{ij}\}$ – $n \times n$ matrix.
- M_{ij} – its minor $\Leftrightarrow M_{ij}$ is an $(n - 1) \times (n - 1)$ matrix resulting from removing i -th row and j -th column from A .

$$\det A = \sum_{i=1}^n (-1)^{i+j} a_{ij} \det M_{ij} = \sum_{j=1}^n (-1)^{i+j} a_{ij} \det M_{ij}$$

Computing Determinant

- $A = \{a_{ij}\}$ – $n \times n$ matrix.
- M_{ij} – its minor $\Leftrightarrow M_{ij}$ is an $(n - 1) \times (n - 1)$ matrix resulting from removing i -th row and j -th column from A .

$$\det A = \sum_{i=1}^n (-1)^{i+j} a_{ij} \det M_{ij} = \sum_{j=1}^n (-1)^{i+j} a_{ij} \det M_{ij}$$

- Laplace expansion.

Some Properties of the Determinant

- $\det A^T = \det A$
- $\det AB = \det A \cdot \det B$
- $\det A^{-1} = \frac{1}{\det A}$

Finding Inverse of a Matrix



Gaussian Elimination

- $A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 1 \\ 1 & 1 & 1 \end{pmatrix}$
- $\det A \neq 0 \Rightarrow$ there exists A^{-1} . Let's find it!



Gaussian Elimination

- $A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 1 \\ 1 & 1 & 1 \end{pmatrix}$
- $\det A \neq 0 \Rightarrow$ there exists A^{-1} . Let's find it!
- Augment the initial matrix:

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array} \right)$$



Gaussian Elimination

- $A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 1 \\ 1 & 1 & 1 \end{pmatrix}$
- $\det A \neq 0 \Rightarrow$ there exists A^{-1} . Let's find it!
- Augment the initial matrix:

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array} \right)$$

- Perform **elementary row operations** and obtain identity matrix on the left. The inverse will be on the right!



Gaussian Elimination

- Elementary row operations:
 - swap rows;
 - multiply rows by some number;
 - add / subtract one row to / from another.

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array} \right)$$

Gaussian Elimination

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array}\right) \rightarrow \{(3) - (1)\} \rightarrow \left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array}\right) \rightarrow$$

Gaussian Elimination

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array}\right) \rightarrow \{(3) - (1)\} \rightarrow \left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array}\right) \rightarrow$$

$$\rightarrow \{(2) - 2 \cdot (3)\} \rightarrow \left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 2 & 1 & -2 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array}\right)$$

Gaussian Elimination

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array}\right) \rightarrow \{(3) - (1)\} \rightarrow \left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array}\right) \rightarrow$$

$$\rightarrow \{(2) - 2 \cdot (3)\} \rightarrow \left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 2 & 1 & -2 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array}\right) \rightarrow \{(1) - (2)\} \rightarrow \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & -1 & -1 & 2 \\ 0 & 0 & 1 & 2 & 1 & -2 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array}\right)$$

Gaussian Elimination

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array}\right) \rightarrow \{(3) - (1)\} \rightarrow \left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array}\right) \rightarrow$$

$$\rightarrow \{(2) - 2 \cdot (3)\} \rightarrow \left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 2 & 1 & -2 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array}\right) \rightarrow \{(1) - (2)\} \rightarrow \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & -1 & -1 & 2 \\ 0 & 0 & 1 & 2 & 1 & -2 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array}\right) \rightarrow$$

$$\rightarrow \{\text{swap (2) and (3)}\} \rightarrow \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & -1 & -1 & 2 \\ 0 & 1 & 0 & -1 & 0 & 1 \\ 0 & 0 & 1 & 2 & 1 & -2 \end{array}\right)$$

Gaussian Elimination

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array}\right) \rightarrow \dots \rightarrow \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & -1 & -1 & 2 \\ 0 & 1 & 0 & -1 & 0 & 1 \\ 0 & 0 & 1 & 2 & 1 & -2 \end{array}\right)$$

$$A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 1 \\ 1 & 1 & 1 \end{pmatrix}, \quad A^{-1} = \begin{pmatrix} -1 & -1 & 2 \\ -1 & 0 & 1 \\ 2 & 1 & -2 \end{pmatrix}, \quad AA^{-1} = A^{-1}A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Rank



Column Space

- Consider a square matrix A .
- Its columns A^1, \dots, A^n can be seen as vectors.

Column Space

- Consider a square matrix A .
- Its columns A^1, \dots, A^n can be seen as vectors.
- $U = \text{span}\{A^1, \dots, A^n\}$ – **column space** of A .
 - All vectors that can be obtain by linearly combining columns of A .
 - \Leftrightarrow **image** of linear transformation A (= all the vectors we can get by applying A).

Rank

- Column space $U = \text{span}\{A^1, \dots, A^n\}$ is the image of linear transformation A .
- **Rank** of a matrix is the number of dimensions in its column space.

Rank

- Column space $U = \text{span}\{A^1, \dots, A^n\}$ is the image of linear transformation A .
- **Rank** of a matrix is the number of dimensions in its column space.
 - Full rank matrix: n columns, all linearly independent.
 - Lower-rank matrices: linearly dependent columns present.

Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$

- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$

- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 1$

- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$

- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 1$

- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 2$

- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 1$

- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 2$

- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \text{rank}(A) = 3$

Column vs Row Rank

- Column space of A = span of A 's columns.
Its dimensionality = (column) rank.

Column vs Row Rank

- Column space of A = span of A 's columns.
Its dimensionality = (column) rank.
- We can also define the row space of A as a span of A 's rows.
Its dimensionality = row rank.

Column vs Row Rank

- Column space of A = span of A 's columns.
Its dimensionality = (column) rank.
- We can also define the row space of A as a span of A 's rows.
Its dimensionality = row rank.
- Column rank vs. row rank?

Column vs Row Rank

- Column space of A = span of A 's columns.
Its dimensionality = (column) rank.
- We can also define the row space of A as a span of A 's rows.
Its dimensionality = row rank.
- Column rank vs. row rank?
- **Fundamental result: the column rank and the row rank are always equal.**
See [proofs](#).

Rank

- This is why there cannot be more than n linearly independent vectors in \mathbb{R}^n !

Rank

- This is why there cannot be more than n linearly independent vectors in \mathbb{R}^n !
- Imagine $x_1, x_2, \dots, x_m \in \mathbb{R}^n$:

Rank

- This is why there cannot be more than n linearly independent vectors in \mathbb{R}^n !
- Imagine $x_1, x_2, \dots, x_m \in \mathbb{R}^n$:

$$X = [x_1 \mid x_2 \mid \dots \mid x_n] = \begin{bmatrix} x_{11} & x_{12} & \dots & x_{1m} \\ x_{21} & x_{22} & \dots & x_{2m} \\ \dots & \dots & \dots & \dots \\ x_{n1} & x_{n2} & \dots & x_{nm} \end{bmatrix}$$

Rank

- This is why there cannot be more than n linearly independent vectors in \mathbb{R}^n !
- Imagine $x_1, x_2, \dots, x_m \in \mathbb{R}^n$:

$$X = [x_1 \mid x_2 \mid \dots \mid x_m] = \begin{bmatrix} x_{11} & x_{12} & \dots & x_{1m} \\ x_{21} & x_{22} & \dots & x_{2m} \\ \dots & \dots & \dots & \dots \\ x_{n1} & x_{n2} & \dots & x_{nm} \end{bmatrix}$$

$$\text{rank}(X) \leq \min\{n, m\}$$

Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 1$

- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 2$

- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \text{rank}(A) = 3$

Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 1 \Leftrightarrow \mathbb{R}^3 \text{ is mapped onto a line}$

- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 2$

- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \text{rank}(A) = 3$

Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$, $\text{rank}(A) = 1 \Leftrightarrow \mathbb{R}^3$ is mapped onto a line
- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$, $\text{rank}(A) = 2 \Leftrightarrow \mathbb{R}^3$ is mapped onto a plane
- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$, $\text{rank}(A) = 3$

Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$, $\text{rank}(A) = 1 \Leftrightarrow \mathbb{R}^3$ is mapped onto a line
- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$, $\text{rank}(A) = 2 \Leftrightarrow \mathbb{R}^3$ is mapped onto a plane
- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$, $\text{rank}(A) = 3 \Leftrightarrow \mathbb{R}^3$ is mapped on itself (isomorphism)

Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$, $\text{rank}(A) = 1 \Leftrightarrow \mathbb{R}^3$ is mapped onto a line
Infinitely many vectors are mapped into a zero vector.
- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$, $\text{rank}(A) = 2 \Leftrightarrow \mathbb{R}^3$ is mapped onto a plane
Infinitely many vectors are mapped into a zero vector.
- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$, $\text{rank}(A) = 3 \Leftrightarrow \mathbb{R}^3$ is mapped on itself (isomorphism)
Only a zero vector is mapped into a zero vector.

Null space

- A set of vectors that are mapped to $\mathbf{0}$ by a linear transformation A .

Null space

- A set of vectors that are mapped to $\mathbf{0}$ by a linear transformation A .
- Example: projection onto XY -plane:

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad A \begin{bmatrix} 0 \\ 0 \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \Rightarrow$$

$$\text{Null space: } \left\{ v \in \mathbb{R}^3 \mid v = \begin{bmatrix} 0 \\ 0 \\ z \end{bmatrix}, z \in \mathbb{R} \right\}$$

Systems of Linear Equations



What is a SLE?

$$\begin{cases} 2x_1 + 5x_2 + 3x_3 = -3 \\ 4x_1 + 0x_2 + 8x_3 = 0 \\ 1x_1 + 3x_2 + 0x_3 = 2 \end{cases}$$

Solutions to SLE



$$1. \quad \begin{cases} x + y = 1 \\ x + y = 2 \end{cases}$$

$$2. \quad \begin{cases} x + y = 1 \\ 2x + y = 2 \end{cases}$$

$$3. \quad \begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

Solutions to SLE



1.
$$\begin{cases} x + y = 1 \\ x + y = 2 \end{cases}$$

No solutions.

2.
$$\begin{cases} x + y = 1 \\ 2x + y = 2 \end{cases}$$

A single solution: $x = 1, y = 0$

3.
$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

Infinitely many solutions.

SLE: Matrix Notation

$$\begin{cases} 2x_1 + 5x_2 + 3x_3 = -3 \\ 4x_1 + 0x_2 + 8x_3 = 0 \\ 1x_1 + 3x_2 + 0x_3 = 2 \end{cases}$$

SLE: Matrix Notation

$$\begin{cases} 2x_1 + 5x_2 + 3x_3 = -3 \\ 4x_1 + 0x_2 + 8x_3 = 0 \\ 1x_1 + 3x_2 + 0x_3 = 2 \end{cases}$$

$$A = \begin{bmatrix} 2 & 5 & 3 \\ 4 & 0 & 8 \\ 1 & 3 & 0 \end{bmatrix}, \quad b = \begin{bmatrix} -3 \\ 0 \\ 2 \end{bmatrix}, \quad x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$Ax = b$$

SLE: Matrix Notation

$$\begin{cases} 2x_1 + 5x_2 + 3x_3 = -3 \\ 4x_1 + 0x_2 + 8x_3 = 0 \\ 1x_1 + 3x_2 + 0x_3 = 2 \end{cases}$$

$$A = \begin{bmatrix} 2 & 5 & 3 \\ 4 & 0 & 8 \\ 1 & 3 & 0 \end{bmatrix}, \quad b = \begin{bmatrix} -3 \\ 0 \\ 2 \end{bmatrix}, \quad x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$Ax = b$$

“Find vector(s) x that are mapped into b by transform A ”

Number of Solutions

- $Ax = b$: “Find vector(s) x that are mapped into b by transform A ”.

Number of Solutions

- $Ax = b$: “Find vector(s) x that are mapped into b by transform A ”.
- If $\det A \neq 0$, there exists A^{-1} : $x = A^{-1}b$.
So, there is a single solution.

Number of Solutions

- $Ax = b$: “Find vector(s) x that are mapped into b by transform A ”.
- If $\det A \neq 0$, there exists A^{-1} : $x = A^{-1}b$.
So, there is a single solution.
- If $\det A = 0$, it's not that simple:

Number of Solutions

- $Ax = b$: “Find vector(s) x that are mapped into b by transform A ”.
- If $\det A \neq 0$, there exists A^{-1} : $x = A^{-1}b$.
So, there is a single solution.
- If $\det A = 0$, it's not that simple:

A maps the original space into a lower dimensional subspace.

Number of Solutions

- $Ax = b$: “Find vector(s) x that are mapped into b by transform A ”.
- If $\det A \neq 0$, there exists A^{-1} : $x = A^{-1}b$.
So, there is a single solution.
- If $\det A = 0$, it's not that simple:

A maps the original space into a lower dimensional subspace.

If we are lucky, b is there, and we have infinitely many solutions.

If not, there're no solutions.

Solutions to SLE



1.
$$\begin{cases} x + y = 1 \\ x + y = 2 \end{cases}$$

No solutions.

2.
$$\begin{cases} x + y = 1 \\ 2x + y = 2 \end{cases}$$

A single solution: $x = 1, y = 0$

$$A = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix}, \det A \neq 0.$$

3.
$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

Infinitely many solutions.

Solutions to SLE



1.
$$\begin{cases} x + y = 1 \\ x + y = 2 \end{cases}$$

2.
$$\begin{cases} x + y = 1 \\ 2x + y = 2 \end{cases}$$

3.
$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

No solutions.

$A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$, $\det A = 0$. A maps \mathbb{R}^2 onto a line, $b = [1, 2]^T$ isn't there.

A single solution: $x = 1$, $y = 0$

$$A = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix}, \det A \neq 0.$$

Infinitely many solutions.

Solutions to SLE



1.
$$\begin{cases} x + y = 1 \\ x + y = 2 \end{cases}$$

2.
$$\begin{cases} x + y = 1 \\ 2x + y = 2 \end{cases}$$

3.
$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

No solutions.

$A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$, $\det A = 0$. A maps \mathbb{R}^2 onto a line, $b = [1, 2]^T$ isn't there.

A single solution: $x = 1$, $y = 0$

$$A = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix}, \det A \neq 0.$$

Infinitely many solutions.

$A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$, $\det A = 0$. A maps \mathbb{R}^2 onto a line, $b = [1, 2]^T$ is there.

Number of Solutions

- $Ax = b$: “Find vector(s) x that are mapped into b by transform A ”.
- If $\det A \neq 0$, there exists A^{-1} : $x = A^{-1}b$.
So, there is a single solution.
- If $\det A = 0$, it's not that simple:

A maps the original space into a lower dimensional subspace.

If we are lucky, b is there, and we have infinitely many solutions.

If not, there're no solutions.

How do we check that?

Number of Solutions

- $Ax = b$ – SLE.

- Consider matrix $(A|b) = \begin{bmatrix} a_{11} & \dots & a_{1n} & b_1 \\ \dots & \dots & \dots & \dots \\ a_{m1} & \dots & a_{mn} & b_n \end{bmatrix}$.

Number of Solutions

- $Ax = b$ – SLE.

- Consider matrix $(A|b) = \begin{bmatrix} a_{11} & \dots & a_{1n} & b_1 \\ \dots & \dots & \dots & \dots \\ a_{m1} & \dots & a_{mn} & b_n \end{bmatrix}$.

- $Ax = b$ has a unique solution $\Leftrightarrow \text{rank}(A|b) = \text{rank}(A) = n$.

Number of Solutions

- $Ax = b$ – SLE.

- Consider matrix $(A|b) = \begin{bmatrix} a_{11} & \dots & a_{1n} & b_1 \\ \dots & \dots & \dots & \dots \\ a_{m1} & \dots & a_{mn} & b_n \end{bmatrix}$.

- $Ax = b$ has a unique solution $\Leftrightarrow \text{rank}(A|b) = \text{rank}(A) = n$.
- $Ax = b$ has infinitely many solutions $\Leftrightarrow \text{rank}(A|b) = \text{rank}(A) < n$.

Number of Solutions

- $Ax = b$ – SLE.

- Consider matrix $(A|b) = \begin{bmatrix} a_{11} & \dots & a_{1n} & b_1 \\ \dots & \dots & \dots & \dots \\ a_{m1} & \dots & a_{mn} & b_n \end{bmatrix}$.

- $Ax = b$ has a unique solution $\Leftrightarrow \text{rank}(A|b) = \text{rank}(A) = n$.
- $Ax = b$ has infinitely many solutions $\Leftrightarrow \text{rank}(A|b) = \text{rank}(A) < n$.
- $Ax = b$ has no solutions $\Leftrightarrow \text{rank}(A|b) > \text{rank}(A)$.

Solutions to SLE



1.
$$\begin{cases} x + y = 1 \\ x + y = 2 \end{cases}$$

No solutions.

$$1 = \text{rank}(A) < \text{rank}(A|b) = 2$$

2.
$$\begin{cases} x + y = 1 \\ 2x + y = 2 \end{cases}$$

A single solution: $x = 1, y = 0$

$$\text{rank}(A) = \text{rank}(A|b) = 2$$

3.
$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

Infinitely many solutions.

$$\text{rank}(A) = \text{rank}(A|b) = 1 < 2$$

Gaussian Elimination



Gaussian elimination

- $Ax = b$

$$A = \begin{bmatrix} 1 & 3 & -2 \\ 3 & 5 & 6 \\ 2 & 4 & 3 \end{bmatrix}, \quad b = \begin{bmatrix} 5 \\ 7 \\ 8 \end{bmatrix}$$

Gaussian elimination

- $Ax = b$

$$A = \begin{bmatrix} 1 & 3 & -2 \\ 3 & 5 & 6 \\ 2 & 4 & 3 \end{bmatrix}, \quad b = \begin{bmatrix} 5 \\ 7 \\ 8 \end{bmatrix}$$

- Elementary row operations:

$$\begin{aligned} \left[\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 3 & 5 & 6 & 7 \\ 2 & 4 & 3 & 8 \end{array} \right] &\sim \left[\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & -4 & 12 & -8 \\ 2 & 4 & 3 & 8 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & -4 & 12 & -8 \\ 0 & -2 & 7 & -2 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 1 & -3 & 2 \\ 0 & -2 & 7 & -2 \end{array} \right] \\ &\sim \left[\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 1 & -3 & 2 \\ 0 & 0 & 1 & 2 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 1 & 0 & 8 \\ 0 & 0 & 1 & 2 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 3 & 0 & 9 \\ 0 & 1 & 0 & 8 \\ 0 & 0 & 1 & 2 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 0 & 0 & -15 \\ 0 & 1 & 0 & 8 \\ 0 & 0 & 1 & 2 \end{array} \right] \end{aligned}$$

Gaussian elimination

- $Ax = b$

$$A = \begin{bmatrix} 1 & 3 & -2 \\ 3 & 5 & 6 \\ 2 & 4 & 3 \end{bmatrix}, \quad b = \begin{bmatrix} 5 \\ 7 \\ 8 \end{bmatrix}$$

- Elementary row operations:

$$\begin{aligned} \left[\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 3 & 5 & 6 & 7 \\ 2 & 4 & 3 & 8 \end{array} \right] &\sim \left[\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & -4 & 12 & -8 \\ 2 & 4 & 3 & 8 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & -4 & 12 & -8 \\ 0 & -2 & 7 & -2 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 1 & -3 & 2 \\ 0 & -2 & 7 & -2 \end{array} \right] \\ &\sim \left[\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 1 & -3 & 2 \\ 0 & 0 & 1 & 2 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 1 & 0 & 8 \\ 0 & 0 & 1 & 2 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 3 & 0 & 9 \\ 0 & 1 & 0 & 8 \\ 0 & 0 & 1 & 2 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 0 & 0 & -15 \\ 0 & 1 & 0 & 8 \\ 0 & 0 & 1 & 2 \end{array} \right] \end{aligned}$$

Unique solution.

Gaussian Elimination

- $Ax = b$ – SLE.

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & -3 \\ 2 & 1 & 5 \end{bmatrix}, \quad b = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$$

Gaussian Elimination

- $Ax = b$ – SLE.

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & -3 \\ 2 & 1 & 5 \end{bmatrix}, \quad b = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$$

- Elementary row operations:

$$\left(\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 1 & -3 & 1 \\ 2 & 1 & 5 & 0 \end{array} \right) \rightarrow \dots \rightarrow \left(\begin{array}{ccc|c} 1 & 0 & 4 & 1 \\ 0 & 1 & -3 & 1 \\ 0 & 0 & 0 & -3 \end{array} \right)$$

Gaussian Elimination

- $Ax = b$ – SLE.

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & -3 \\ 2 & 1 & 5 \end{bmatrix}, \quad b = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$$

- Elementary row operations:

$$\left(\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 1 & -3 & 1 \\ 2 & 1 & 5 & 0 \end{array} \right) \rightarrow \dots \rightarrow \left(\begin{array}{ccc|c} 1 & 0 & 4 & 1 \\ 0 & 1 & -3 & 1 \\ 0 & 0 & 0 & -3 \end{array} \right)$$

No solutions.

Gaussian Elimination

- $Ax = b$ – SLE.

$$A = \begin{bmatrix} -3 & -5 & 36 \\ -1 & 0 & 7 \\ 1 & 1 & -10 \end{bmatrix}, \quad b = \begin{bmatrix} 10 \\ 5 \\ -4 \end{bmatrix}$$

-

Gaussian Elimination

- $Ax = b$ – SLE.

$$A = \begin{bmatrix} -3 & -5 & 36 \\ -1 & 0 & 7 \\ 1 & 1 & -10 \end{bmatrix}, \quad b = \begin{bmatrix} 10 \\ 5 \\ -4 \end{bmatrix}$$

- Elementary row operations:

$$\left(\begin{array}{ccc|c} -3 & -5 & 36 & 10 \\ -1 & 0 & 7 & 5 \\ 1 & 1 & -10 & -4 \end{array} \right) \rightarrow \dots \rightarrow \left(\begin{array}{ccc|c} 1 & 0 & -7 & -5 \\ 0 & 2 & -3 & -1 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

Gaussian Elimination

- $Ax = b$ – SLE.

$$A = \begin{bmatrix} -3 & -5 & 36 \\ -1 & 0 & 7 \\ 1 & 1 & -10 \end{bmatrix}, \quad b = \begin{bmatrix} 10 \\ 5 \\ -4 \end{bmatrix}$$

- Elementary row operations:

$$\left(\begin{array}{ccc|c} -3 & -5 & 36 & 10 \\ -1 & 0 & 7 & 5 \\ 1 & 1 & -10 & -4 \end{array} \right) \rightarrow \dots \rightarrow \left(\begin{array}{ccc|c} 1 & 0 & -7 & -5 \\ 0 & 2 & -3 & -1 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

Infinitely many solutions.

Homogeneous SLE



Homogeneous SLE

$$\begin{cases} 2x_1 + 5x_2 + 3x_3 = 0 \\ 4x_1 + 0x_2 + 8x_3 = 0 \\ 1x_1 + 3x_2 + 0x_3 = 0 \end{cases}$$

Homogeneous SLE

$$Ax = 0$$

Homogeneous SLE

$$Ax = 0$$

Solutions = null space of A .

Homogeneous SLE

$$Ax = 0$$

Solutions = null space of A .

$\text{rank } A = \# \text{variables} \rightarrow$ unique solution (0)

$\text{rank } A < \# \text{variables} \rightarrow$ infinitely many solutions.

Homogeneous SLE

- $Ax = 0$
- Let V be a set of solutions:

$$\forall v \in V \quad Av = 0$$

Homogeneous SLE

- $Ax = 0$
- Let V be a set of solutions:

$$\forall v \in V \quad Av = 0$$

- Note that
 - $0 \in V$ because $A \cdot 0 = 0$.

Homogeneous SLE

- $Ax = 0$
- Let V be a set of solutions:

$$\forall v \in V \quad Av = 0$$

- Note that
 - $0 \in V$ because $A \cdot 0 = 0$.
 - $\forall v \in V \rightarrow \lambda v \in V$
Indeed, $A(\lambda v) = \lambda Av = 0$

Homogeneous SLE

- $Ax = 0$
- Let V be a set of solutions:

$$\forall v \in V \quad Av = 0$$

- Note that
 - $0 \in V$ because $A \cdot 0 = 0$.
 - $\forall v \in V \rightarrow \lambda v \in V$
Indeed, $A(\lambda v) = \lambda Av = 0$
 - $\forall v_1, v_2 \in V \rightarrow (v_1 + v_2) \in V$.
Indeed, $A(v_1 + v_2) = Av_1 + Av_2 = 0$

Homogeneous SLE

- $Ax = 0$
- Let V be a set of solutions:

$$\forall v \in V \quad Av = 0$$

- Note that
 - $0 \in V$ because $A \cdot 0 = 0$.
 - $\forall v \in V \rightarrow \lambda v \in V$
Indeed, $A(\lambda v) = \lambda Av = 0$
 - $\forall v_1, v_2 \in V \rightarrow (v_1 + v_2) \in V$.
Indeed, $A(v_1 + v_2) = Av_1 + Av_2 = 0$

V is a linear subspace!

To sum up

- Matrices as linear transforms
- Examples of common transforms
- Inverse
- Determinant
- Rank
- Solutions to SLE