


Math Refresher for DS

Lecture 1



girafe
ai



Today



1. Course overview
2. Linear Algebra
 - Core objects
 - Vector spaces
3. A bit of Analytic Geometry
 - Orthogonal projections
 - Hyperplanes
 - Normals

About this course




Image source: Mathematics for Machine Learning, p. 14
(<https://mml-book.github.io/book/mml-book.pdf>)

About this course

Abstract white line art on a blue background, consisting of several rounded, irregular shapes that resemble stylized letters or geometric forms.

In this course:

1. Linear algebra
2. Calculus
3. (Basic) optimization

Prerequisites:

- basic knowledge of math;
- some Python.

About this course



Logistics

- Pre-recorded lectures
- Online practical sessions
 - Tuesdays & Fridays
19:00 Moscow time
- 5 graded assignments
- 2 exams
- Final grade:
 - 30% Linear Algebra exam
 - 30% Calculus & Optimization exam
 - 40% graded assignments

About me

Evgeniya Korneva

✉ evgeniakorneva@gmail.com

in [evgeniyako](#)

◦ PhD researcher

KU LEUVEN

◦ Lecturer



◦ DS content lead



◦ *(Soon)* Data Scientists



Linear Algebra: the Basics



Linear Algebra: Core Objects

- $\alpha \in \mathbb{R}$ - a scalar *Example: -2*

Linear Algebra: Core Objects

- $\alpha \in \mathbb{R}$ - a scalar *Example: -2*
- $x \in \mathbb{R}^n$ - a vector with n entries

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \in \mathbb{R}^3, \quad \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix} \in \mathbb{R}^5$$

Linear Algebra: Core Objects

- $\alpha \in \mathbb{R}$ - a scalar *Example: -2*
- $x \in \mathbb{R}^n$ - a vector with n entries

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \in \mathbb{R}^3, \quad \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix} \in \mathbb{R}^5$$

- $A \in \mathbb{R}^{m \times n}$ - a matrix with m rows and n columns

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}, \quad \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \in \mathbb{R}^{2 \times 3}$$

Linear Algebra: Core Objects

- $\alpha \in \mathbb{R}$ - a scalar *Example: -2*
- $x \in \mathbb{R}^n$ - a **vector** with n entries

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \in \mathbb{R}^3, \quad \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix} \in \mathbb{R}^5$$

- $A \in \mathbb{R}^{m \times n}$ - a matrix with m rows and n columns


$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}, \quad \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \in \mathbb{R}^{2 \times 3}$$

What are Vectors?

- Ordered sets of numbers: $x = [1, 2]$

What are Vectors?


- Ordered sets of numbers: $x = [1, 2]$
- A point with Cartesian coordinates




What are Vectors?

- Ordered sets of numbers: $x = [1, 2]$

- A point with Cartesian coordinates



- Direction + length



Vector Spaces



Vector Space: Definition

- A real-valued vector space $(V, +, \cdot)$ is a set of vectors V with two operations

$$(1) +: V \times V \rightarrow V, \quad (2) \cdot: \mathbb{R} \times V \rightarrow V$$

Vector Space: Definition

- A real-valued vector space $(V, +, \cdot)$ is a set of vectors V with two operations

$$(1) +: V \times V \rightarrow V, \quad (2) \cdot: \mathbb{R} \times V \rightarrow V$$

that satisfy the following properties (axioms):

	Property	Meaning
1.	Associativity of addition	$x + (y + z) = (x + y) + z$
2.	Commutativity of addition	$x + y = y + x$
3.	Identity element of addition	$\exists 0 \in V: \forall x \in V \quad 0 + x = x$
4.	Identity element of scalar multiplication	$\forall x \in V \quad 1 \cdot x = x$
5.	Inverse element of addition	$\forall x \in V \exists -x \in V: x + (-x) = 0$
6.	Compatibility of scalar multiplication	$\alpha(\beta x) = (\alpha\beta)x$
7.	Distributivity	$(\alpha + \beta)x = \alpha x + \beta x$
8.		$\alpha(x + y) = \alpha x + \alpha y$

Let's define vector operations!

Operations with Vectors

1. Sum of two vectors:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^n, \quad y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} \in \mathbb{R}^n, \quad x + y = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_2 \end{bmatrix} \in \mathbb{R}^n$$

Operations with Vectors

1. Sum of two vectors:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^n, \quad y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} \in \mathbb{R}^n, \quad x + y = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_2 \end{bmatrix} \in \mathbb{R}^n$$

2. Multiplying by a scalar:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^n, \quad \alpha \in \mathbb{R}, \quad \alpha x = \begin{bmatrix} \alpha x_1 \\ \alpha x_2 \\ \vdots \\ \alpha x_n \end{bmatrix} \in \mathbb{R}^n$$

Operations with Vectors: Example

$x, y \in \mathbb{R}^3$:

$$x = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

Operations with Vectors: Example

$$x, y \in \mathbb{R}^3:$$

$$x = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

Multiplying by a scalar:

$$10x = \begin{bmatrix} 10 \\ 0 \\ 10 \end{bmatrix}$$

Operations with Vectors: Example

$$x, y \in \mathbb{R}^3:$$

$$x = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

Multiplying by a scalar:

$$10x = \begin{bmatrix} 10 \\ 0 \\ 10 \end{bmatrix}$$

Sum:


$$x + y = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$$

Vector Operations: Geometrical Interpretation

Vectors: Geometrical Interpretation



$$\vec{a} = [2, 3]$$




Vectors: Geometrical Interpretation



$$\vec{a} = [2, 3]$$

$$2\vec{a} = [4, 6]$$




Vectors: Geometrical Interpretation



$$\vec{a} = [2, 3]$$

$$2\vec{a} = [4, 6]$$

$$-0.5\vec{a} = [-1, -1.5]$$




Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$




Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$

$$\vec{a} + \vec{b} = [6, 6]$$




Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$

$$\vec{a} + \vec{b} = [6, 6]$$




Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$

$$\vec{a} + \vec{b} = [6, 6]$$




Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$

$$\vec{a} + \vec{b} = [6, 6]$$



Back to Vector Spaces

Operations with Vectors

1. Sum of two vectors:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}, \quad x + y = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_2 \end{bmatrix} \in \mathbb{R}^n$$

2. Multiplying by a scalar:

satisfy axioms (1) – (8)
(check it yourself)

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad \alpha \in \mathbb{R}, \quad \alpha x = \begin{bmatrix} \alpha x_1 \\ \alpha x_2 \\ \vdots \\ \alpha x_n \end{bmatrix} \in \mathbb{R}^n$$

Vector Spaces

$(\mathbb{R}^n, +, \cdot), n \in \mathbb{N}$ - a vector space with operations

1. vector addition:

$$x + y = (x_1, x_2, \dots, x_n) + (y_1, y_2, \dots, y_n) = (x_1 + y_1, \dots, x_n + y_n)$$

2. multiplication by a scalar:

$$\alpha x = \alpha(x_1, x_2, \dots, x_n) = (\alpha x_1, \alpha x_2, \dots, \alpha x_n)$$

Vector Spaces: Another Example

Vector Spaces: Another Example

- \mathbb{P}^n - a set of polynomials of degree $\leq n$ with real coefficients
 - *Example:* $\mathbb{P}^3 = \{ax^3 + bx^2 + cx + d \mid a, b, c, d \in \mathbb{R}\}$

Vector Spaces: Another Example

- \mathbb{P}^n - a set of polynomials of degree $\leq n$ with real coefficients
 - Example: $\mathbb{P}^3 = \{ax^3 + bx^2 + cx + d \mid a, b, c, d \in \mathbb{R}\}$
- We can add up polynomials:

$$\begin{aligned}(a_n x^n + a_{n-1} x^{n-1} + \dots + a_0) + (b_n x^n + b_{n-1} x^{n-1} + \dots + b_0) &= \\ = (a_n + b_n) x^n + (a_{n-1} + b_{n-1}) x^{n-1} + \dots + (a_0 + b_0) &\in \mathbb{P}^n\end{aligned}$$

Vector Spaces: Another Example

- \mathbb{P}^n - a set of polynomials of degree $\leq n$ with real coefficients

- Example: $\mathbb{P}^3 = \{ax^3 + bx^2 + cx + d \mid a, b, c, d \in \mathbb{R}\}$

- We can add up polynomials:

$$\begin{aligned} (a_n x^n + a_{n-1} x^{n-1} + \dots + a_0) + (b_n x^n + b_{n-1} x^{n-1} + \dots + b_0) &= \\ = (a_n + b_n) x^n + (a_{n-1} + b_{n-1}) x^{n-1} + \dots + (a_0 + b_0) &\in \mathbb{P}^n \end{aligned}$$

- And we can multiply them by a real number:

$$\lambda(a_n x^n + a_{n-1} x^{n-1} + \dots + a_0) = \lambda a_n x^n + \lambda a_{n-1} x^{n-1} + \dots + \lambda a_0 \in \mathbb{P}^n$$

Vector Spaces: Another Example

- \mathbb{P}^n - a set of polynomials of degree $\leq n$ with real coefficients

- Example: $\mathbb{P}^3 = \{ax^3 + bx^2 + cx + d \mid a, b, c, d \in \mathbb{R}\}$

- We can add up polynomials:

$$\begin{aligned}(a_n x^n + a_{n-1} x^{n-1} + \dots + a_0) + (b_n x^n + b_{n-1} x^{n-1} + \dots + b_0) &= \\ &= (a_n + b_n) x^n + (a_{n-1} + b_{n-1}) x^{n-1} + \dots + (a_0 + b_0) \in \mathbb{P}^n\end{aligned}$$

- And we can multiply them by a real number:

$$\lambda(a_n x^n + a_{n-1} x^{n-1} + \dots + a_0) = \lambda a_n x^n + \lambda a_{n-1} x^{n-1} + \dots + \lambda a_0 \in \mathbb{P}^n$$

- These operations satisfy axioms (1) – (8)!

$\rightarrow (\mathbb{P}^n, +, \cdot)$ is also a vector space!

Inner Product



Inner Product

- Inner product is a function $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{R}$ that satisfies the following properties:
 - *Symmetric*: $\forall x, y \in V \quad \langle x, y \rangle = \langle y, x \rangle$
 - *Positive definite*: $\forall x \in V \setminus \{0\} \quad \langle x, x \rangle > 0$ and $\langle x, 0 \rangle = 0$.

Dot Product

- A particular type of inner product.

Dot Product

- A particular type of inner product.
- For $x = [x_1, \dots, x_n]$, $y = [y_1, \dots, y_n] \in \mathbb{R}^n$

$$(x, y) = x_1y_1 + x_2y_2 + \dots + x_ny_n$$

Dot Product

- A particular type of inner product.

- For $x = [x_1, \dots, x_n]$, $y = [y_1, \dots, y_n] \in \mathbb{R}^n$

$$(x, y) = x_1y_1 + x_2y_2 + \dots + x_ny_n$$

- Example:

$$x = [1, 2, 3, 4], \quad y = [-1, 0, 1, 2]$$

$$(x, y) = 1 \cdot (-1) + 2 \cdot 0 + 3 \cdot 1 + 4 \cdot 2 = -1 + 0 + 3 + 8 = 10$$

Euclidian Vector Space

- A vector space with inner product is called an *inner product space*.

Euclidian Vector Space

- A vector space with inner product is called an *inner product space*.
- $(\mathbb{R}^n, +, \cdot)$ with dot product (\cdot, \cdot) is called a *Euclidian vector space*.

Euclidian Vector Space

- A vector space with inner product is called an *inner product space*.
- $(\mathbb{R}^n, +, \cdot)$ with dot product (\cdot, \cdot) is called a *Euclidian vector space*.
- *Note: there're inner products different from dot product.*

Norms



Norm

- A **norm** on a vector space \mathbb{V} is a function $\|\cdot\|: \mathbb{V} \rightarrow \mathbb{R}$ which assigns each vector $x \in \mathbb{V}$ its *length* $\|x\| \in \mathbb{R}$.

Norm

- A **norm** on a vector space \mathbb{V} is a function $\|\cdot\|: \mathbb{V} \rightarrow \mathbb{R}$ which assigns each vector $x \in \mathbb{V}$ its *length* $\|x\| \in \mathbb{R}$.
- A norm must satisfy the following properties:

Norm


- A **norm** on a vector space \mathbb{V} is a function $\|\cdot\|: \mathbb{V} \rightarrow \mathbb{R}$ which assigns each vector $x \in \mathbb{V}$ its *length* $\|x\| \in \mathbb{R}$.
- A norm must satisfy the following properties:
 - *Absolutely homogeneous*: $\forall x \in \mathbb{V}, \forall \lambda \in \mathbb{R} \quad \|\lambda x\| = |\lambda| \|x\|$

Norm

- A **norm** on a vector space \mathbb{V} is a function $\|\cdot\|: \mathbb{V} \rightarrow \mathbb{R}$ which assigns each vector $x \in \mathbb{V}$ its *length* $\|x\| \in \mathbb{R}$.
- A norm must satisfy the following properties:
 - *Absolutely homogeneous*: $\forall x \in \mathbb{V}, \forall \lambda \in \mathbb{R} \quad \|\lambda x\| = |\lambda| \|x\|$
 - *Positive definite*: $\forall x \in \mathbb{V} \quad \|x\| \geq 0$ and $\|x\| = 0 \Leftrightarrow x = 0$

Norm

- A **norm** on a vector space \mathbb{V} is a function $\|\cdot\|: \mathbb{V} \rightarrow \mathbb{R}$ which assigns each vector $x \in \mathbb{V}$ its *length* $\|x\| \in \mathbb{R}$.
- A norm must satisfy the following properties:
 - *Absolutely homogeneous*: $\forall x \in \mathbb{V}, \forall \lambda \in \mathbb{R} \quad \|\lambda x\| = |\lambda| \|x\|$
 - *Positive definite*: $\forall x \in \mathbb{V} \quad \|x\| \geq 0$ and $\|x\| = 0 \Leftrightarrow x = 0$
 - *Triangle inequality*: $\forall x, y \in \mathbb{V} \quad \|x + y\| \leq \|x\| + \|y\|$



Examples of Norms

Manhattan Norm



- A norm for $x \in \mathbb{R}^n$:

$$\|x\|_1 = \sum_{i=1}^n |x_i|$$



Manhattan Norm

- A norm for $x \in \mathbb{R}^n$:

$$\|x\|_1 = \sum_{i=1}^n |x_i|$$

- Examples:

$$\|[1, 2, 3]\|_1 = 1 + 2 + 3 = 6$$

$$\|[1, 0]\|_1 = 1 + 0 = 1$$

$$\|[-1, 0]\|_1 = |-1| + 0 = 1$$



Manhattan Norm

- A norm for $x \in \mathbb{R}^n$:


$$\|x\|_1 = \sum_{i=1}^n |x_i|$$

- Examples:

$$\|[1, 2, 3]\|_1 = 1 + 2 + 3 = 5$$

$$\|[1, 0]\|_1 = 1 + 0 = 1$$

$$\|[-1, 0]\|_1 = |-1| + 0 = 1$$





Euclidian Norm

- A norm for $x \in \mathbb{R}^n$:

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- Computes *Euclidian distance* from the origin.



Euclidian Norm

- A norm for $x \in \mathbb{R}^n$:

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- Computes *Euclidian distance* from the origin.



Euclidian Norm

- A norm for $x \in \mathbb{R}^n$:

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- Computes *Euclidian distance* from the origin.
- Examples:

$$\|[1, 2, 3]\|_2 = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}$$

$$\|[1, 0]\|_2 = \sqrt{1^2 + 0} = 1$$

$$\|[-1, 0]\|_2 = \sqrt{(-1)^2 + 0} = 1$$



Euclidian Norm

- A norm for $x \in \mathbb{R}^n$:


$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- Computes *Euclidian distance* from the origin.
- Examples:

$$\|[1, 2, 3]\|_2 = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}$$

$$\|[1, 0]\|_2 = \sqrt{1^2 + 0} = 1$$

$$\|[-1, 0]\|_2 = \sqrt{(-1)^2 + 0} = 1$$



Other norms

- In general, for $x = [x_1, \dots, x_n] \in \mathbb{R}^n$ an ℓ_p -norm is defined as follows:

$$\|x\|_p = \sqrt[p]{\sum_{i=1}^n |x_i|^p}.$$

Other norms

- In general, for $x = [x_1, \dots, x_n] \in \mathbb{R}^n$ an ℓ_p -norm is defined as follows:

$$\|x\|_p = \sqrt[p]{\sum_{i=1}^n |x_i|^p}.$$

- ℓ_1 - Manhattan norm $\|\cdot\|_1$;
- ℓ_2 - Euclidian norm $\|\cdot\|$ (default);

Other norms

- In general, for $x = [x_1, \dots, x_n] \in \mathbb{R}^n$ an ℓ_p -norm is defined as follows:

$$\|x\|_p = \sqrt[p]{\sum_{i=1}^n |x_i|^p}.$$

- ℓ_1 - Manhattan norm $\|\cdot\|_1$;
- ℓ_2 - Euclidian norm $\|\cdot\|$ (default);
- ℓ_∞ : $\|x\|_\infty = \max_i |x_i|$

Example: $\|[1, 2, 3]\|_\infty = 3$, $\|[1, 0]\|_\infty = 1$, $\|[-1, 0.5]\|_\infty = 1$.

Inner Product and Norms

- Any inner product induces a norm:

$$\|x\| := \sqrt{\langle x, x \rangle}$$

Inner Product and Norms

- Any inner product induces a norm:

$$\|x\| := \sqrt{\langle x, x \rangle}$$

Check yourself that this is indeed a norm.

Inner Product and Norms

- Any inner product induces a norm:

$$\|x\| := \sqrt{\langle x, x \rangle}$$

Check yourself that this is indeed a norm.

- Example: dot product induces Euclidian norm:

$$\sqrt{(x, x)} = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2} = \|x\|_2$$

Inner Product and Norms

- Any inner product induces a norm:

$$\|x\| := \sqrt{\langle x, x \rangle}$$

Check yourself that this is indeed a norm.

- Example: dot product induces Euclidian norm:

$$\sqrt{(x, x)} = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2} = \|x\|_2$$

- (!) Not every norm is induced by an inner product.
Example: Manhattan norm.

Cauchy-Schwarz Inequality

- For an inner product vector space, the induced norm satisfies the inequality:

$$|\langle x, y \rangle| \leq \|x\| \cdot \|y\|$$

Cauchy-Schwarz Inequality

- For an inner product vector space, the induced norm satisfies the inequality:

$$|\langle x, y \rangle| \leq \|x\| \cdot \|y\|$$

- For dot product and Euclidian norm:

$$|(x, y)| \leq \|x\|_2 \cdot \|y\|_2$$

Distance between Vectors

- Distance between two vectors x and y is defined as

$$d(x, y) := \|x - y\| = \sqrt{\langle x - y, x - y \rangle}.$$

Distance between Vectors

- Distance between two vectors x and y is defined as

$$d(x, y) := \|x - y\| = \sqrt{\langle x - y, x - y \rangle}.$$

- For dot product and Euclidian norm, we get *Euclidian distance*:

$$\begin{aligned} d(x, y) &= \|x - y\|_2 = \sqrt{(x - y, x - y)} = \\ &= \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \dots + (x_n - y_n)^2}. \end{aligned}$$

Angles and Orthogonality




Angle between Two Vectors

- Inner product also captures the geometry of vector space by defining the angle between two vectors.
- Remember Cauchy-Schwarz inequality:

$$|(x, y)| \leq \|x\| \cdot \|y\|$$

$$-1 \leq \frac{(x, y)}{\|x\| \cdot \|y\|} \leq 1$$

$$\omega: \cos \omega = \frac{(x, y)}{\|x\| \cdot \|y\|} \text{ - angle between } x \text{ and } y.$$



Angle between Two Vectors: Example

- What is the angle ω between $x = [5, 0]$ and $y = [1, 1]$?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{5 \cdot 1 + 0 \cdot 1}{\sqrt{5^2 + 0^2} \cdot \sqrt{1^2 + 1^2}} = \arccos \frac{5}{5\sqrt{2}} = \arccos \frac{\sqrt{2}}{4} = \frac{\pi}{4}.$$

Angle between Two Vectors: Example

- What is the angle ω between $x = [5, 0]$ and $y = [1, 1]$?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{5 \cdot 1 + 0 \cdot 1}{\sqrt{5^2 + 0^2} \cdot \sqrt{1^2 + 1^2}} = \arccos \frac{5}{5\sqrt{2}} = \arccos \frac{\sqrt{2}}{4} = \frac{\pi}{4}.$$

- What is the angle ω between $x = [\sqrt{3}, 0, 1]$ and $y = [1, 0, 0]$?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{\sqrt{3}}{\sqrt{3 + 1} \cdot \sqrt{1}} = \arccos \frac{\sqrt{3}}{2} = \frac{\pi}{6}.$$

Angle between Two Vectors: Example

- What is the angle ω between $x = [5, 0]$ and $y = [1, 1]$?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{5 \cdot 1 + 0 \cdot 1}{\sqrt{5^2 + 0^2} \cdot \sqrt{1^2 + 1^2}} = \arccos \frac{5}{5\sqrt{2}} = \arccos \frac{\sqrt{2}}{4} = \frac{\pi}{4}.$$

- What is the angle ω between $x = [\sqrt{3}, 0, 1]$ and $y = [1, 0, 0]$?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{\sqrt{3}}{\sqrt{3 + 1} \cdot \sqrt{1}} = \arccos \frac{\sqrt{3}}{2} = \frac{\pi}{6}.$$

- What is the angle ω between $x = [1, 0, 0, 0, 1]$ and $y = [0, 1, 1, 0, 0]$?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{0}{\sqrt{2} \cdot \sqrt{2}} = \arccos 0 = \frac{\pi}{2}.$$

Orthogonality

- x and y are orthogonal if and only if $\langle x, y \rangle = 0$.

Orthogonality

- x and y are orthogonal if and only if $\langle x, y \rangle = 0$.
- Examples in \mathbb{R}^n with dot product:

$$x = [2, 3], \quad y = [1, 0], \quad (x, y) = 2 \neq 0 \rightarrow \\ x \text{ and } y \text{ are not orthogonal.}$$

Orthogonality

- x and y are orthogonal if and only if $\langle x, y \rangle = 0$.
- Examples in \mathbb{R}^n with dot product:

$$x = [2, 3], \quad y = [1, 0], \quad (x, y) = 2 \neq 0 \rightarrow \\ x \text{ and } y \text{ are not orthogonal.}$$

$$x = [1, 2, 3], \quad y = [-2, 1, 0], \quad (x, y) = -2 + 2 + 0 = 0 \rightarrow \\ x \text{ and } y \text{ are orthogonal.}$$

Orthogonality

- x and y are orthogonal if and only if $\langle x, y \rangle = 0$.
- Examples in \mathbb{R}^n with dot product:

$$x = [2, 3], \quad y = [1, 0], \quad (x, y) = 2 \neq 0 \rightarrow \\ x \text{ and } y \text{ are not orthogonal.}$$


$$x = [1, 2, 3], \quad y = [-2, 1, 0], \quad (x, y) = -2 + 2 + 0 = 0 \rightarrow \\ x \text{ and } y \text{ are orthogonal.}$$

$$x = [1, 0], \quad y = [0, 1], \quad (x, y) = 0, \quad \|x\| = \|y\| = 1 \rightarrow \\ x \text{ and } y \text{ are } \textit{orthonormal}.$$

Orthogonal Projection




- Suppose we have two vectors u and v .



Orthogonal Projection




- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .



Orthogonal Projection



- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?




Orthogonal Projection



- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?

If $0 \leq \omega \leq 90$


$$\begin{aligned}(u, v) &= \|u\| \|v\| \cos \omega = \|u\| \|v\| \frac{\|u'_v\|}{\|u\|} = \\ &= \|u'_v\| \|v\|\end{aligned}$$



Orthogonal Projection

- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?


If $90 \leq \omega \leq 180$



Orthogonal Projection

- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?


If $90 \leq \omega \leq 180$



Orthogonal Projection

- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?

If $90 \leq \omega \leq 180$




Orthogonal Projection

- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?

If $90 \leq \omega \leq 180$

$$\begin{aligned} -(u, v) &= \|u\| \|v\| \cos \omega = \|u\| \|v\| \frac{\|u'_v\|}{\|u\|} = \\ &= \|u'_v\| \|v\| \end{aligned}$$




Orthogonal Projection



- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?

$$|(\mathbf{u}, \mathbf{v})| = \|\mathbf{u}'_v\| \|\mathbf{v}\| \iff \|\mathbf{u}'_v\| = \frac{|(\mathbf{u}, \mathbf{v})|}{\|\mathbf{v}\|}$$




Orthogonal Projection



- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?


$$|(u, v)| = \|u'_v\| \|v\| \iff \|u'_v\| = \frac{|(u, v)|}{\|v\|}$$

$$u'_v = \frac{(u, v)}{(v, v)} v.$$



Orthogonal Projection: Example

- What's projection of $u = [1, 3, 2]$ on $z = [0, 0, 1]$?




Orthogonal Projection: Example

- What's projection of $u = [1, 3, 2]$ on $z = [0, 0, 1]$?

$$u'_z = \frac{(u, z)}{(z, z)} z = 2z = [0, 0, 2]$$

(Projection on the axis = drop other coordinates)




Orthogonal Projection: Example

- What's projection of $u = [1, 3, 2]$ on $z = [0, 0, 1]$?

$$u'_z = \frac{(u, z)}{(z, z)} z = 2z = [0, 0, 2]$$

(Projection on the axis = drop other coordinates)

- What's projection of $u = [1, 3, 2]$ on $v = [4, 1, 3]$?



Orthogonal Projection: Example


- What's projection of $u = [1, 3, 2]$ on $z = [0, 0, 1]$?

$$u'_z = \frac{(u, z)}{(z, z)} z = 2z = [0, 0, 2]$$

(Projection on the axis = drop other coordinates)

- What's projection of $u = [1, 3, 2]$ on $v = [4, 1, 3]$?

$$u'_v = \frac{(u, v)}{(v, v)} v = \frac{4 + 3 + 6}{16 + 1 + 9} v = \frac{1}{2} v = [2, 0.5, 1.5].$$



Hyperplanes

- A hyperplane is described by equation

$$w_1x_1 + w_2x_2 + \cdots + w_nx_n + b = 0$$

where at least one $w_i \neq 0$.

- A more compact notation:

$$(w, x) + b = 0, \quad w = (w_1, \dots, w_n)$$

Hyperplanes



Hyperplanes


- A hyperplane in \mathbb{R}^n is described by equation

$$w_1x_1 + w_2x_2 + \cdots + w_nx_n + b = 0$$

where at least one $w_i \neq 0$.

Hyperplanes

A hyperplane in \mathbb{R}^2 is a line



A hyperplane in \mathbb{R}^3 is a plane




Image source: <https://deepai.org/machine-learning-glossary-and-terms/hyperplane>

Hyperplanes

- A hyperplane in \mathbb{R}^n is described by equation

$$w_1x_1 + w_2x_2 + \cdots + w_nx_n + b = 0$$

where at least one $w_i \neq 0$.

- A more compact notation:

$$(w, x) + b = 0, \quad w = (w_1, \dots, w_n)$$




Normal to a Hyperplane

- Consider a hyperplane $(w, x) + b = 0$.
- Vector $w = (w_1, \dots, w_n)$ defines the hyperplane.




Normal to a Hyperplane

- Consider a hyperplane $(w, x) + b = 0$.
- Vector $w = (w_1, \dots, w_n)$ defines the hyperplane.
- w is a *normal vector* to this hyperplane: it's orthogonal to every vector on it.




ML Example: Linear Classifier

- Objects = 2D vectors
- Binary classification: classes +1 and -1.




ML Example: Linear Classifier

- Objects = 2D vectors
- Binary classification: classes +1 and -1.
- Linear classifier: separating hyperplane $(w, x) + b = 0$
 - objects “above” are class +1
 - objects “below” are class -1.




ML Example: Linear Classifier

- Objects = 2D vectors
- Binary classification: classes +1 and -1.
- Linear classifier: separating hyperplane $(w, x) + b = 0$
 - objects “above” are class +1
 - objects “below” are class -1.
- How can we formalize this?



ML Example: Linear Classifier

- Objects = 2D vectors
- Binary classification: classes +1 and -1.
- Linear classifier: separating hyperplane $(w, x) + b = 0$
 - objects “above” are class +1
 - objects “below” are class -1.
- How can we formalize this?
 - objects “above”: $(w, x) + b > 0$
 - objects “below”: $(w, x) + b < 0$



To sum up

- Vectors
 - Vector spaces
 - Inner products
 - Lengths
 - Distances
 - Angles
- Analytic Geometry
 - Projections
 - Hyperplanes
 - Normal vector

Next Time

- More on vector spaces.