




Math Basics for DS

Practical Session 8



Today

- Multivariate calculus

Partial Derivatives

- Derivative for univariate functions:

$$y = f(x), \quad \frac{dy}{dx} = f'(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

Partial Derivatives

- Derivative for univariate functions:

$$y = f(x), \quad \frac{dy}{dx} = f'(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

- Partial derivative for multivariate functions:

$$y = f(x_1, \dots, x_n), \quad \frac{\partial y}{\partial x_i} = f'_{x_i} = \lim_{\Delta x_i \rightarrow 0} \frac{f(x_1, \dots, x_i + \Delta x_i, \dots, x_n) - f(x_1, \dots, x_i, \dots, x_n)}{\Delta x_i}$$

*Compute derivative with respect to x_i
regarding all other variables as constants.*

Partial Derivatives - Example

- Let $f(x, y) = x^3 + x^2y^3 - 2y^2$.
- Find $f_x(2,1)$ and $f_y(2,1)$.

Partial Derivatives - Example

- Let $f(x, y) = x^3 + x^2y^3 - 2y^2$.
- Find $f_x(2,1)$ and $f_y(2,1)$.

$$f_x(x, y) = \frac{\partial f}{\partial x} =$$

$$f_y(x, y) = \frac{\partial f}{\partial y} =$$

Partial Derivatives - Example

- Let $f(x, y) = x^3 + x^2y^3 - 2y^2$.
- Find $f_x(2,1)$ and $f_y(2,1)$.

$$f_x(x, y) = \frac{\partial f}{\partial x} = 3x^2 + 2xy^3$$

$$f_y(x, y) = \frac{\partial f}{\partial y} =$$

Partial Derivatives - Example

- Let $f(x, y) = x^3 + x^2y^3 - 2y^2$.
- Find $f_x(2,1)$ and $f_y(2,1)$.

$$f_x(x, y) = \frac{\partial f}{\partial x} = 3x^2 + 2xy^3 \Big|_{(2,1)} =$$

$$f_y(x, y) = \frac{\partial f}{\partial y} =$$

Partial Derivatives - Example

- Let $f(x, y) = x^3 + x^2y^3 - 2y^2$.
- Find $f_x(2,1)$ and $f_y(2,1)$.

$$f_x(x, y) = \frac{\partial f}{\partial x} = 3x^2 + 2xy^3 \Big|_{(2,1)} = 12 + 4 = 16$$

$$f_y(x, y) = \frac{\partial f}{\partial y} =$$

Partial Derivatives - Example

- Let $f(x, y) = x^3 + x^2y^3 - 2y^2$.
- Find $f_x(2,1)$ and $f_y(2,1)$.

$$f_x(x, y) = \frac{\partial f}{\partial x} = 3x^2 + 2xy^3 \Big|_{(2,1)} = 12 + 4 = 16$$

$$f_y(x, y) = \frac{\partial f}{\partial y} = 3x^2y^2 - 4y$$

Partial Derivatives - Example

- Let $f(x, y) = x^3 + x^2y^3 - 2y^2$.
- Find $f_x(2,1)$ and $f_y(2,1)$.

$$f_x(x, y) = \frac{\partial f}{\partial x} = 3x^2 + 2xy^3 \Big|_{(2,1)} = 12 + 4 = 16$$

$$f_y(x, y) = \frac{\partial f}{\partial y} = 3x^2y^2 - 4y \Big|_{(2,1)} =$$

Partial Derivatives - Example

- Let $f(x, y) = x^3 + x^2y^3 - 2y^2$.
- Find $f_x(2,1)$ and $f_y(2,1)$.

$$f_x(x, y) = \frac{\partial f}{\partial x} = 3x^2 + 2xy^3 \Big|_{(2,1)} = 12 + 4 = 16$$

$$f_y(x, y) = \frac{\partial f}{\partial y} = 3x^2y^2 - 4y \Big|_{(2,1)} = 12 - 4 = 8$$

The Chain Rule

- For univariate functions:

$$\left(f(g(x))\right)' = f'(g(x)) \cdot g'(x) \quad \Leftrightarrow \quad \frac{df}{dx} = \frac{df}{dg} \cdot \frac{dg}{dx}$$

The Chain Rule

- For univariate functions:

$$\left(f(g(x))\right)' = f'(g(x)) \cdot g'(x) \iff \frac{df}{dx} = \frac{df}{dg} \cdot \frac{dg}{dx}$$

- For multivariate functions:

$$u(x_1, \dots, x_n), \quad x_i = x_i(t_1, \dots, t_m)$$

$$\frac{\partial u}{\partial t_k} = \frac{\partial u}{\partial x_1} \cdot \frac{\partial x_1}{\partial t_k} + \frac{\partial u}{\partial x_2} \cdot \frac{\partial x_2}{\partial t_k} + \dots + \frac{\partial u}{\partial x_n} \cdot \frac{\partial x_n}{\partial t_k}, \quad k = 1, \dots, m$$

The Chain Rule - Example

$$u = x^2y + 3xy^4, \quad x = \sin 2t, \quad y = \cos t$$

The Chain Rule - Example

$$u = x^2y + 3xy^4, \quad x = \sin 2t, \quad y = \cos t$$

$$\frac{\partial u}{\partial t} = \frac{\partial u}{\partial x} \cdot \frac{\partial x}{\partial t} + \frac{\partial u}{\partial y} \cdot \frac{\partial y}{\partial t}$$

The Chain Rule - Example

$$u = x^2y + 3xy^4, \quad x = \sin 2t, \quad y = \cos t$$

$$\frac{\partial u}{\partial t} = \frac{\partial u}{\partial x} \cdot \frac{\partial x}{\partial t} + \frac{\partial u}{\partial y} \cdot \frac{\partial y}{\partial t}$$

$$\frac{\partial u}{\partial t} =$$

The Chain Rule - Example

$$u = x^2y + 3xy^4, \quad x = \sin 2t, \quad y = \cos t$$

$$\frac{\partial u}{\partial t} = \frac{\partial u}{\partial x} \cdot \frac{\partial x}{\partial t} + \frac{\partial u}{\partial y} \cdot \frac{\partial y}{\partial t}$$

$$\frac{\partial u}{\partial t} = (2xy + 3y^4) \cdot 2 \cos 2t +$$

The Chain Rule - Example

$$u = x^2y + 3xy^4, \quad x = \sin 2t, \quad y = \cos t$$

$$\frac{\partial u}{\partial t} = \frac{\partial u}{\partial x} \cdot \frac{\partial x}{\partial t} + \frac{\partial u}{\partial y} \cdot \frac{\partial y}{\partial t}$$

$$\frac{\partial u}{\partial t} = (2xy + 3y^4) \cdot 2 \cos 2t + (x^2 + 12xy^3)(-\sin t)$$

The Chain Rule - Example

$$u = x^4 y + y^2 z^3$$

$$x = rse^t, \quad y = rs^2e^{-t}, \quad z = r^2s \sin t$$

The Chain Rule - Example

$$u = x^4 y + y^2 z^3$$

$$x = rse^t, \quad y = rs^2e^{-t}, \quad z = r^2s \sin t$$

$$\frac{\partial u}{\partial s} = \frac{\partial u}{\partial x} \cdot \frac{\partial x}{\partial s} + \frac{\partial u}{\partial y} \cdot \frac{\partial y}{\partial s} + \frac{\partial u}{\partial z} \cdot \frac{\partial z}{\partial s}$$

The Chain Rule - Example

$$u = x^4 y + y^2 z^3$$

$$x = rse^t, \quad y = rs^2e^{-t}, \quad z = r^2s \sin t$$

$$\frac{\partial u}{\partial s} = \frac{\partial u}{\partial x} \cdot \frac{\partial x}{\partial s} + \frac{\partial u}{\partial y} \cdot \frac{\partial y}{\partial s} + \frac{\partial u}{\partial z} \cdot \frac{\partial z}{\partial s}$$

$$\frac{\partial u}{\partial s} =$$

The Chain Rule - Example

$$u = x^4 y + y^2 z^3$$

$$x = rse^t, \quad y = rs^2e^{-t}, \quad z = r^2s \sin t$$

$$\frac{\partial u}{\partial s} = \frac{\partial u}{\partial x} \cdot \frac{\partial x}{\partial s} + \frac{\partial u}{\partial y} \cdot \frac{\partial y}{\partial s} + \frac{\partial u}{\partial z} \cdot \frac{\partial z}{\partial s}$$

$$\frac{\partial u}{\partial s} = 4x^3 y \cdot re^t + (x^4 + 2yz^3) \cdot 2rse^{-t} + 3y^2 z^2 \cdot r^2 \sin t$$

Gradient

- A vector of partial derivatives:

$$\nabla f(x_1, \dots, x_n) = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)$$

Gradient

- A vector of partial derivatives:

$$\nabla f(x_1, \dots, x_n) = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)$$

- Example:

$$f(x, y, z) = e^{xy} \ln z$$

Gradient

- A vector of partial derivatives:

$$\nabla f(x_1, \dots, x_n) = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)$$

- Example:

$$f(x, y, z) = e^{xy} \ln z$$

$$\nabla f(x, y, z) = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right) =$$

Gradient

- A vector of partial derivatives:

$$\nabla f(x_1, \dots, x_n) = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)$$

- Example:

$$f(x, y, z) = e^{xy} \ln z$$

$$\nabla f(x, y, z) = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right) = \left(ye^{xy} \ln z, \quad xe^{xy} \ln z, \quad \frac{e^{xy}}{z} \right).$$

Gradient

Gradient shows the direction of the maximal growth of the function.

Negative gradient = direction of the maximal descent of the function.

Higher Derivatives

- Consider $f(x, y)$.
- $f'_x(x, y)$, $f'_y(x, y)$ – also functions of two variables.
We can compute their partial derivatives!

Higher Derivatives

- Consider $f(x, y)$.
- $f'_x(x, y)$, $f'_y(x, y)$ – also functions of two variables.
We can compute their partial derivatives!

$$(f'_x)'_x = f''_{xx}(x, y) = \frac{\partial^2 f}{\partial x^2}, \quad (f'_x)'_y = f''_{xy}(x, y) = \frac{\partial^2 f}{\partial y \partial x}$$

$$(f'_y)'_y = f''_{yy}(x, y) = \frac{\partial^2 f}{\partial y^2}, \quad (f'_y)'_x = f''_{yx}(x, y) = \frac{\partial^2 f}{\partial x \partial y}$$

Higher Derivatives - Example

- Consider $f(x, y) = x^3 + x^2y^3 - 2y^2$.

$$f_x(x, y) = \frac{\partial f}{\partial x} = 3x^2 + 2xy^3, \quad f_y(x, y) = \frac{\partial f}{\partial y} = 3x^2y^2 - 4y$$

$$f_{xx}''(x, y) = \frac{\partial^2 f}{\partial x^2} = \quad , \quad f_{xy}''(x, y) = \frac{\partial^2 f}{\partial y \partial x} =$$

$$f_{yy}''(x, y) = \frac{\partial^2 f}{\partial y^2} = \quad f_{yx}''(x, y) = \frac{\partial^2 f}{\partial x \partial y} =$$

Higher Derivatives - Example

- Consider $f(x, y) = x^3 + x^2y^3 - 2y^2$.

$$f_x(x, y) = \frac{\partial f}{\partial x} = 3x^2 + 2xy^3, \quad f_y(x, y) = \frac{\partial f}{\partial y} = 3x^2y^2 - 4y$$

$$f_{xx}''(x, y) = \frac{\partial^2 f}{\partial x^2} = 6x + 2y^3, \quad f_{xy}''(x, y) = \frac{\partial^2 f}{\partial y \partial x} =$$

$$f_{yy}''(x, y) = \frac{\partial^2 f}{\partial y^2} = \quad f_{yx}''(x, y) = \frac{\partial^2 f}{\partial x \partial y} =$$

Higher Derivatives - Example

- Consider $f(x, y) = x^3 + x^2y^3 - 2y^2$.

$$f_x(x, y) = \frac{\partial f}{\partial x} = 3x^2 + 2xy^3, \quad f_y(x, y) = \frac{\partial f}{\partial y} = 3x^2y^2 - 4y$$

$$f_{xx}''(x, y) = \frac{\partial^2 f}{\partial x^2} = 6x + 2y^3, \quad f_{xy}''(x, y) = \frac{\partial^2 f}{\partial y \partial x} = 6xy^2$$

$$f_{yy}''(x, y) = \frac{\partial^2 f}{\partial y^2} = \quad f_{yx}''(x, y) = \frac{\partial^2 f}{\partial x \partial y} =$$

Higher Derivatives - Example

- Consider $f(x, y) = x^3 + x^2y^3 - 2y^2$.

$$f_x(x, y) = \frac{\partial f}{\partial x} = 3x^2 + 2xy^3, \quad f_y(x, y) = \frac{\partial f}{\partial y} = 3x^2y^2 - 4y$$

$$f_{xx}''(x, y) = \frac{\partial^2 f}{\partial x^2} = 6x + 2y^3, \quad f_{xy}''(x, y) = \frac{\partial^2 f}{\partial y \partial x} = 6xy^2$$

$$f_{yy}''(x, y) = \frac{\partial^2 f}{\partial y^2} = 6x^2y - 4, \quad f_{yx}''(x, y) = \frac{\partial^2 f}{\partial x \partial y} =$$

Higher Derivatives - Example

- Consider $f(x, y) = x^3 + x^2y^3 - 2y^2$.

$$f_x(x, y) = \frac{\partial f}{\partial x} = 3x^2 + 2xy^3, \quad f_y(x, y) = \frac{\partial f}{\partial y} = 3x^2y^2 - 4y$$

$$f_{xx}''(x, y) = \frac{\partial^2 f}{\partial x^2} = 6x + 2y^3, \quad f_{xy}''(x, y) = \frac{\partial^2 f}{\partial y \partial x} = 6xy^2$$

$$f_{yy}''(x, y) = \frac{\partial^2 f}{\partial y^2} = 6x^2y - 4, \quad f_{yx}''(x, y) = \frac{\partial^2 f}{\partial x \partial y} = 6xy^2$$

Hessian

- A matrix of second derivatives.
- $f(x_1, \dots, x_n)$

$$H = \begin{pmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_1 x_2} & \cdots & \frac{\partial^2 f}{\partial x_1 x_n} \\ \frac{\partial^2 f}{\partial x_2 x_1} & \frac{\partial^2 f}{\partial x_2^2} & \cdots & \frac{\partial^2 f}{\partial x_2 x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n x_1} & \frac{\partial^2 f}{\partial x_n x_2} & \cdots & \frac{\partial^2 f}{\partial x_n^2} \end{pmatrix}$$

Hessian - Example

- $f(x, y) = x^3 + x^2y^3 - 2y^2$

$$f''_{xx}(x, y) = \frac{\partial^2 f}{\partial x^2} = 6x + 2y^3, \quad f''_{xy}(x, y) = \frac{\partial^2 f}{\partial y \partial x} = 6xy^2$$

$$f''_{yy}(x, y) = \frac{\partial^2 f}{\partial y^2} = 6x^2y - 4, \quad f''_{yx}(x, y) = \frac{\partial^2 f}{\partial x \partial y} = 6xy^2$$

$$H = \begin{pmatrix} & \end{pmatrix}$$

Hessian - Example

- $f(x, y) = x^3 + x^2y^3 - 2y^2$

$$f''_{xx}(x, y) = \frac{\partial^2 f}{\partial x^2} = 6x + 2y^3, \quad f''_{xy}(x, y) = \frac{\partial^2 f}{\partial y \partial x} = 6xy^2$$

$$f''_{yy}(x, y) = \frac{\partial^2 f}{\partial y^2} = 6x^2y - 4, \quad f''_{yx}(x, y) = \frac{\partial^2 f}{\partial x \partial y} = 6xy^2$$

$$H = \begin{pmatrix} 6x + 2y^3 & 6xy^2 \\ 6xy^2 & 6x^2y - 4 \end{pmatrix}$$

Extrema

- Univariate case: a stationary point x_0 is a local **minimum** (**maximum**) if

$$f''(x_0) > 0 \text{ (} f''(x_0) < 0 \text{)}.$$

Extrema

- Univariate case: a stationary point x_0 is a local **minimum** (**maximum**) if

$$f''(x_0) > 0 \text{ (} f''(x_0) < 0 \text{)}.$$

- Multivariate case:

x_0 is a stationary point: $\nabla f(x_0) = 0$

H – Hessian.


Then, if H is a **positive-definite** matrix, then x_0 is a **local minimum**.

If H is a **negative-definite** matrix, then x_0 is a **local maximum**.

If $\det H = 0$, we need to check manually.

Otherwise, x_0 is a **saddle point**.

Saddle Points



Positive vs Negative Definite Matrices

- A matrix A is called positive-definite if

$$x^T A x > 0 \quad \forall x \neq 0$$

- A matrix A is called negative-definite if

$$x^T A x < 0 \quad \forall x \neq 0$$

Positive vs Negative Definite Matrices



- How to check if a matrix is positive (negative) definite?

Positive vs Negative Definite Matrices



- How to check if a matrix is positive (negative) definite?

Check principal minors D_k !

For positive definite:

$$D_1 > 0, \quad D_2 > 0, \quad \dots, D_n > 0$$

For negative definite:

$$D_1 < 0, \quad D_2 > 0, \quad D_3 < 0, \dots$$

$$H = \begin{pmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_2 x_1} & \dots & \frac{\partial^2 f}{\partial x_n x_1} \\ \frac{\partial^2 f}{\partial x_1 x_2} & \frac{\partial^2 f}{\partial x_2^2} & \dots & \frac{\partial^2 f}{\partial x_2 x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n x_1} & \frac{\partial^2 f}{\partial x_n x_2} & \dots & \frac{\partial^2 f}{\partial x_n^2} \end{pmatrix}$$

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) =$$

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) = (4x^3 - 4y, \quad 4y^3 - 4x)$$

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) = (4x^3 - 4y, \quad 4y^3 - 4x)$$

Stationary points: $\nabla f = 0 \iff$

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) = (4x^3 - 4y, \quad 4y^3 - 4x)$$

Stationary points: $\nabla f = 0 \iff$

$$\begin{aligned} 4x^3 - 4y &= 0 \\ 4y^3 - 4x &= 0 \end{aligned} \iff$$

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) = (4x^3 - 4y, \quad 4y^3 - 4x)$$

Stationary points: $\nabla f = 0 \iff$

$$\begin{array}{l} 4x^3 - 4y = 0 \\ 4y^3 - 4x = 0 \end{array} \iff \begin{array}{l} y = x^3 \\ x^9 - x = 0 \end{array} \iff$$

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) = (4x^3 - 4y, \quad 4y^3 - 4x)$$

Stationary points: $\nabla f = 0 \iff$

$$\begin{array}{l} 4x^3 - 4y = 0 \\ 4y^3 - 4x = 0 \end{array} \iff \begin{array}{l} y = x^3 \\ x^9 - x = 0 \end{array} \iff \begin{array}{l} y = x^3 \\ x(x^8 - 1) = 0 \end{array}$$

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) = (4x^3 - 4y, \quad 4y^3 - 4x)$$

Stationary points: $\nabla f = 0 \iff$

$$\begin{array}{lcl} 4x^3 - 4y = 0 & \iff & y = x^3 \\ 4y^3 - 4x = 0 & \iff & x^9 - x = 0 \iff x(x^8 - 1) = 0 \end{array}$$

Stationary points: $(0, 0), (-1, -1), (1, 1)$

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) = (4x^3 - 4y, \quad 4y^3 - 4x)$$

Stationary points: $(0, 0)$, $(-1, -1)$, $(1, 1)$

$$H = \begin{pmatrix} & \end{pmatrix}$$

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) = (4x^3 - 4y, \quad 4y^3 - 4x)$$

Stationary points: $(0, 0)$, $(-1, -1)$, $(1, 1)$

$$H = \begin{pmatrix} 12x^2 & -4 \\ -4 & 12y^2 \end{pmatrix}$$

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) = (4x^3 - 4y, \quad 4y^3 - 4x)$$

Stationary points: $(0, 0)$, $(-1, -1)$, $(1, 1)$

$$H = \begin{pmatrix} 12x^2 & -4 \\ -4 & 12y^2 \end{pmatrix}$$

$\det H = 144x^2y^2 - 16|_{(0;0)} < 0$, $(0, 0)$ – saddle point.

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) = (4x^3 - 4y, \quad 4y^3 - 4x)$$

Stationary points: $(0, 0)$, $(-1, -1)$, $(1, 1)$

$$H = \begin{pmatrix} 12x^2 & -4 \\ -4 & 12y^2 \end{pmatrix}$$

$$\det H = 144x^2y^2 - 16 \quad , \quad \det H_1 = 12x^2$$

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) = (4x^3 - 4y, \quad 4y^3 - 4x)$$

Stationary points: $(0, 0)$, $(-1, -1)$, $(1, 1)$

$$H = \begin{pmatrix} 12x^2 & -4 \\ -4 & 12y^2 \end{pmatrix}$$

$$\det H = 144x^2y^2 - 16 \Big|_{(1;1)} > 0, \quad \det H_1 = 12x^2 \Big|_{(1,1)} > 0$$

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) = (4x^3 - 4y, \quad 4y^3 - 4x)$$

Stationary points: $(0, 0)$, $(-1, -1)$, $(1, 1)$

$$H = \begin{pmatrix} 12x^2 & -4 \\ -4 & 12y^2 \end{pmatrix}$$

$$\det H = 144x^2y^2 - 16 \Big|_{(1;1)} > 0, \quad \det H_1 = 12x^2 \Big|_{(1,1)} > 0$$

$(1, 1)$ – local minimum.

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) = (4x^3 - 4y, \quad 4y^3 - 4x)$$

Stationary points: $(0, 0)$, $(-1, -1)$, $(1, 1)$

$$H = \begin{pmatrix} 12x^2 & -4 \\ -4 & 12y^2 \end{pmatrix}$$

$$\det H = 144x^2y^2 - 16 \quad , \quad \det H_1 = 12x^2$$

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$

$$\nabla f = (f'_x, f'_y) = (4x^3 - 4y, \quad 4y^3 - 4x)$$

Stationary points: $(0, 0)$, $(-1, -1)$, $(1, 1)$


$$H = \begin{pmatrix} 12x^2 & -4 \\ -4 & 12y^2 \end{pmatrix}$$

$$\det H = 144x^2y^2 - 16 \Big|_{(-1,-1)} > 0, \quad \det H_1 = 12x^2 \Big|_{(-1,-1)} > 0$$

$(-1, -1)$ – local minimum.

Extrema – Example

$$f(x, y) = x^4 + y^4 - 4xy + 1$$



Linear Regression the Other Way



Least Squares (Again)

- Remember the Least Squares problem:

$$\hat{y}_i = w_0 + w_1 x_1^i + \cdots + w_n x_n^i, \quad y_i = \hat{y}_i + e_i, \quad w = ?$$

Least Squares (Again)

- Remember the Least Squares problem:

$$\hat{y}_i = w_0 + w_1 x_1^i + \cdots + w_n x_n^i, \quad y_i = \hat{y}_i + e_i, \quad w = ?$$

- Before: solve by projecting y onto the $\text{col}(A)$:

$$\hat{w} = (X^T X)^{-1} X^T y$$

Least Squares (Again)

- Remember the Least Squares problem:

$$\hat{y}_i = w_0 + w_1 x_1^i + \dots + w_n x_n^i, \quad y_i = \hat{y}_i + e_i, \quad w = ?$$

- Before: solve by projecting y onto the $\text{col}(A)$:

$$\hat{w} = (X^T X)^{-1} X^T y$$

- We can also see this as an optimization problem:

$$\mathcal{L}(w_0, \dots, w_n) = \sum e_i^2 = \sum (y_i - \hat{y})^2 = \sum \left(y_i - (w_0 + w_1 x_1^i + \dots + w_n x_n^i) \right)^2 \rightarrow \min_{w_0, \dots, w_n}$$

Least Squares

- Let's consider a simple case: fitting a line to a set of points

$$(x_1, y_1), \quad (x_2, y_2), \quad \dots, \quad (x_n, y_n)$$

$$y \approx w_0 + w_1 x$$

We want to pick the weights w_0 , w_1
so that the sum of squares of the errors is the smallest possible:

$$\mathcal{L}(w_0, w_1) = \rightarrow \min_{w_0, w_1}$$

Least Squares

- Let's consider a simple case: fitting a line to a set of points

$$(x_1, y_1), \quad (x_2, y_2), \quad \dots, \quad (x_n, y_n)$$

$$y \approx w_0 + w_1 x$$

We want to pick the weights w_0 , w_1
so that the sum of squares of the errors is the smallest possible:

$$\mathcal{L}(w_0, w_1) = \sum_{i=1}^n (y_i - w_0 - w_1 x_i)^2 \rightarrow \min_{w_0, w_1}$$

Least Squares

$$\mathcal{L}(w_0, w_1) = \sum_{i=1}^n (y_i - w_0 - w_1 x_i)^2 \rightarrow \min_{w_0, w_1}$$

$$\frac{\partial \mathcal{L}}{\partial w_1} =$$

$$\frac{\partial \mathcal{L}}{\partial w_0} =$$

Least Squares

$$\mathcal{L}(w_0, w_1) = \sum_{i=1}^n (y_i - w_0 - w_1 x_i)^2 \rightarrow \min_{w_0, w_1}$$

$$\frac{\partial \mathcal{L}}{\partial w_1} = -2 \sum_{i=1}^n x_i \cdot (y_i - w_0 - w_1 x_i) = 0$$

$$\frac{\partial \mathcal{L}}{\partial w_0} = -2 \sum_{i=1}^n (y_i - w_0 - w_1 x_i) = 0$$