




Math Refresher for DS

Practical Session 3



Plan for Today



- A short quiz
- SLE with no solutions
- Practice in Python

Short Quiz Lectures 1 - 3

Solving Systems of Linear Equations

- $Ax = b$ – a system of linear equations (SLE).
- A – $m \times n$ matrix (= m equations, n variables).

Solving Systems of Linear Equations

- $Ax = b$ – a system of linear equations (SLE).
- A – $m \times n$ matrix (= m equations, n variables).
- How can we determine the number of solutions to it?

Solving Systems of Linear Equations

- $Ax = b$ – a system of linear equations (SLE).
- A – $m \times n$ matrix (= m equations, n variables).
- How can we determine the number of solutions to it?
 - $\text{rank}(A) = \text{rank}(A|b) = n \Leftrightarrow$ there is a unique solution $x = A^{-1}b$;
 - $\text{rank}(A) = \text{rank}(A|b) < n \Leftrightarrow$ there are infinitely many solutions;
 - $\text{rank}(A) < \text{rank}(A|b) \Leftrightarrow$ there are no solutions.

Solving Systems of Linear Equations

- $Ax = b$ – a system of linear equations (SLE).
- A – $m \times n$ matrix (= m equations, n variables).
- How can we determine the number of solutions to it?
 - $\text{rank}(A) = \text{rank}(A|b) = n \Leftrightarrow$ there is a unique solution $x = A^{-1}b$;
 - $\text{rank}(A) = \text{rank}(A|b) < n \Leftrightarrow$ there are infinitely many solutions;
 - $\text{rank}(A) < \text{rank}(A|b) \Leftrightarrow$ there are no solutions.
- Sometimes, we need to “solve” systems that don’t have a solution.

Solving Systems of Linear Equations

- $Ax = b$ – a system of linear equations (SLE).
- A – $m \times n$ matrix (= m equations, n variables).
- How can we determine the number of solutions to it?
 - $\text{rank}(A) = \text{rank}(A|b) = n \Leftrightarrow$ there is a unique solution $x = A^{-1}b$;
 - $\text{rank}(A) = \text{rank}(A|b) < n \Leftrightarrow$ there are infinitely many solutions;
 - $\text{rank}(A) < \text{rank}(A|b) \Leftrightarrow$ there are no solutions.
- Sometimes, we need to “solve” systems that don’t have a solution.

How to find a reasonable approximate solution?


Least Squares



Motivating Example

- Imagine that you have m observations:

$(x_1, y_1), \quad (x_2, y_2), \quad \dots, \quad (x_m, y_m)$




Motivating Example

- Imagine that you have m observations:

$$(x_1, y_1), \quad (x_2, y_2), \quad \dots, \quad (x_m, y_m)$$

- You want to draw a line through your observations:




Motivating Example

- Imagine that you have m observations:

$$(x_1, y_1), \quad (x_2, y_2), \quad \dots, \quad (x_m, y_m)$$

- You want to draw a line through your observations:

$$\begin{aligned} w_0 + w_1 x_1 &= y_1 \\ w_0 + w_1 x_2 &= y_2 \\ &\vdots \\ w_0 + w_1 x_m &= y_m \end{aligned} \Leftrightarrow$$




Motivating Example

- Imagine that you have m observations:

$$(x_1, y_1), \quad (x_2, y_2), \quad \dots, \quad (x_m, y_m)$$

- You want to draw a line through your observations:




$$\begin{array}{l} w_0 + w_1 x_1 = y_1 \\ w_0 + w_1 x_2 = y_2 \\ \vdots \\ w_0 + w_1 x_m = y_m \end{array} \Leftrightarrow Xw = y, \text{ where } X = \begin{pmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_m \end{pmatrix}, w = \begin{bmatrix} w_0 \\ w_1 \end{bmatrix}, y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix}$$

Motivating Example

- Imagine that you have m observations:

$$(x_1, y_1), \quad (x_2, y_2), \quad \dots, \quad (x_m, y_m)$$

- You want to draw a line through your observations:



$$\begin{array}{l} w_0 + w_1 x_1 = y_1 \\ w_0 + w_1 x_2 = y_2 \\ \vdots \\ w_0 + w_1 x_m = y_m \end{array} \Leftrightarrow Xw = y, \text{ where } X = \begin{pmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_m \end{pmatrix}, w = \begin{bmatrix} w_0 \\ w_1 \end{bmatrix}, y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix}$$


- Overdetermined system, no solutions.

Motivating Example

- Imagine that you have m observations:

$$(x_1, y_1), \quad (x_2, y_2), \quad \dots, \quad (x_m, y_m)$$

- You want to draw a line through your observations:



$$\begin{array}{l} w_0 + w_1 x_1 = y_1 \\ w_0 + w_1 x_2 = y_2 \\ \vdots \\ w_0 + w_1 x_m = y_m \end{array} \Leftrightarrow Xw = y, \text{ where } X = \begin{pmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_m \end{pmatrix}, w = \begin{bmatrix} w_0 \\ w_1 \end{bmatrix}, y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix}$$


- Overdetermined system, no solutions.
- What shall we do?

Motivating Example

- Imagine that you have m observations:

$$(x_1, y_1), \quad (x_2, y_2), \quad \dots, \quad (x_m, y_m)$$

- You want to draw a line through your observations:



$$\begin{array}{l} w_0 + w_1 x_1 = y_1 \\ w_0 + w_1 x_2 = y_2 \\ \vdots \\ w_0 + w_1 x_m = y_m \end{array} \Leftrightarrow Xw = y, \text{ where } X = \begin{pmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_m \end{pmatrix}, w = \begin{bmatrix} w_0 \\ w_1 \end{bmatrix}, y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix}$$


- Overdetermined system, no solutions.
- What shall we do?

Motivating Example

- Imagine that you have m observations:

$$(x_1, y_1), \quad (x_2, y_2), \quad \dots, \quad (x_m, y_m)$$

- You want to draw a line through your observations:



$$\begin{array}{l} w_0 + w_1 x_1 = y_1 \\ w_0 + w_1 x_2 = y_2 \\ \vdots \\ w_0 + w_1 x_m = y_m \end{array} \Leftrightarrow Xw = y, \text{ where } X = \begin{pmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_m \end{pmatrix}, w = \begin{bmatrix} w_0 \\ w_1 \end{bmatrix}, y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix}$$


- Overdetermined system, no solutions.
- What shall we do? Method of least squares: pick a line such that average squared error is minimized.

Motivating Example

- Imagine that you have m observations:

$$(x_1, y_1), \quad (x_2, y_2), \quad \dots, \quad (x_m, y_m)$$

- You want to draw a line through your observations:




$$\begin{array}{l} w_0 + w_1 x_1 = y_1 \\ w_0 + w_1 x_2 = y_2 \\ \vdots \\ w_0 + w_1 x_m = y_m \end{array} \Leftrightarrow Xw = y, \text{ where } X = \begin{pmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_m \end{pmatrix}, w = \begin{bmatrix} w_0 \\ w_1 \end{bmatrix}, y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix}$$

- Overdetermined system, no solutions.
- What shall we do? Method of least squares: pick a line such that average squared error is minimized.
- Let's look at it from the Linear Algebra perspective.

Method of Least Squares




- $Xw = y$ has no solutions $\Leftrightarrow y$ is not in the column space of X .



Method of Least Squares



- $Xw = y$ has no solutions $\Leftrightarrow y$ is not in the column space of X .
- Let's chose \hat{y} such that




Method of Least Squares



- $Xw = y$ has no solutions $\Leftrightarrow y$ is not in the column space of X .
- Let's chose \hat{y} such that


$$Xw = \hat{y} \text{ has a solution } w^* \\ \Leftrightarrow$$





Method of Least Squares


- $Xw = y$ has no solutions $\Leftrightarrow y$ is not in the column space of X .
- Let's chose \hat{y} such that
$$Xw = \hat{y} \text{ has a solution } w^*$$
$$\Leftrightarrow \hat{y} \text{ is in the column space of } X$$
and





Method of Least Squares


- $Xw = y$ has no solutions $\Leftrightarrow y$ is not in the column space of X .
- Let's chose \hat{y} such that
$$Xw = \hat{y} \text{ has a solution } w^*$$
$$\Leftrightarrow \hat{y} \text{ is in the column space of } X$$
and
$$\hat{y} \text{ is as close to } y \text{ as possible}$$
$$\Leftrightarrow$$





Method of Least Squares

- $Xw = y$ has no solutions $\Leftrightarrow y$ is not in the column space of X .
- Let's choose \hat{y} such that
 $Xw = \hat{y}$ has a solution w^*
 $\Leftrightarrow \hat{y}$ is in the column space of X
and
 \hat{y} is as close to y as possible
 $\Leftrightarrow \|y - \hat{y}\|$ is minimized





Method of Least Squares

- $Xw = y$ has no solutions $\Leftrightarrow y$ is not in the column space of X .
- Let's choose \hat{y} such that
$$Xw = \hat{y} \text{ has a solution } w^*$$
$$\Leftrightarrow \hat{y} \text{ is in the column space of } X$$
and
$$\hat{y} \text{ is as close to } y \text{ as possible}$$
$$\Leftrightarrow \|y - \hat{y}\| \text{ is minimized}$$
- What do \hat{y} and e look like?

