

One query over a ragged measured grid

1 Three fixed/free selection axes give 8 query shapes

Polymer

named set / all 12

Solvent

named set / all 990

Temperature

stored node(s) / all 28

Optional min_red / max_red → source-grain Hansen join + coverage


42 °C → request refusal: temperature_off_grid · nearest 40, 45 °C · success=false

2 Coverage is ragged

stored solubility pairs / 990

■ Hansen RED solvents: 380 / 990

9 polymers 990 

PU 78 

NYLON6 32 

PES 32 

3 Asset stamp ≠ engine disposition

332,640-cell Cartesian envelope

■ 253,456 stored  79,184 not measured

Engine: 252,474 evaluable · 982 unavailable

■ 248,378 raw-valid → served

5,078 asset-flagged split:

■ 4,096 exact_100_artifact → served: saturated

■ 982 nonpositive → measurement-rejected

4 Qualify first; page second

Base selection: LDPE × all 990 solvents × 140 °C · no filters (RED inactive)

Asset: 990 stored = 972 raw-valid + 18 exact_100_artifact

**990
evaluable**



cell predicates
0 exclusions



**990
qualified**



1 2 3 4 5 ... 985 later

offset=0 · returned=5 · has_more=true · next_offset=5

Row: identity · T · exact wt% · BP/margin · atmosphere {true/false/unknown} · GHS signal word
RED active (min_red=0): 380 cells joined · 610 unmatched · 786 source-record rows
Joined rows add RED/Ra/R0 + HSP source IDs; coverage reports 380 of 990 solvents