

Preventive Cloud Seeding for Wildfire Risk Reduction

A staged research framework for silver iodide ice nucleation, antecedent moisture, and human and environmental safety

Aaron Corbin Mazie | Philadelphia, Pennsylvania | 29 September 2026

Research concept paper. This manuscript proposes experiments and decision gates. It presents no original field data and does not establish that cloud seeding reduces wildfires.

Abstract

Wildfire mitigation often begins only after ignition, when heat, wind, and dry fuels sharply limit available response time. This paper proposes an earlier, conditional intervention: augment precipitation from naturally occurring, suitable clouds over selected fire-prone watersheds before anticipated high-risk periods. The initial physical pathway is glaciogenic seeding with silver iodide (AgI) particles in supercooled liquid or mixed-phase clouds, followed by ice growth, precipitation, and measurable changes in snowpack, soil water, or live and dead fuel moisture. The causal chain from seeding to lower fire risk remains unproven. It can fail because suitable clouds are absent, particles do not reach them, precipitation does not increase, water misses relevant fuels, moisture does not persist, or increased vegetation growth later adds burnable fuel. Particular attention is given to the dry diameter, composition, ice-active fraction, atmospheric transport, and human exposure of AgI-containing particles. Published experiments suggest that particle size affects ice nucleation, but no single diameter guarantees either effectiveness or safety. A phased program combines laboratory characterization, retrospective cloud and fuel analysis, prospective monitoring, and only then a permitted, randomized field evaluation. The primary near-term endpoints are precipitation and fuel moisture; reduced wildfire incidence or severity requires a substantially longer and larger evaluation. [1-7]

Keywords: cloud seeding; silver iodide; ice-nucleating particles; wildfire prevention; fuel moisture; environmental exposure; causal evaluation.

1 Problem and proposed mechanism

The intervention is preventive, not an attempt to extinguish an active wildfire. In a region with recurring fire exposure, a public agency and research team would identify the seasonal interval when additional water could plausibly improve snowpack, soil moisture, or fuel moisture. They would then determine how often suitable natural clouds actually occur before those high-risk intervals. Under a tested protocol, an approved contractor could seed only eligible cloud events. The hypothesis is that additional precipitation reaches specified target areas and remains relevant to fuels when fire risk rises. Weather modification cannot create cloud systems or pull water vapor into a dry region. [1,2]

AgI is a glaciogenic ice-nucleating material. Its proposed role is to enter supercooled liquid or mixed-phase cloud regions, initiate additional ice crystals, and permit those crystals to grow through vapor deposition, collection, and aggregation. Precipitation can arrive as snow or, after melting, rain. The phrase “marrying silver iodide with rainclouds” therefore describes physical transport into an eligible cloud and interaction with supercooled droplets; it does not imply a chemical union with water or the creation of a cloud. Warm clouds without suitable supercooled liquid require different hypotheses and are outside the initial AgI pathway. [1,3,4]

The strongest direct evidence of an AgI-induced precipitation chain comes from winter orographic settings. Radar and gauge studies documented attributable snowfall in selected cases. Such evidence cannot be assumed to transfer to a summer wildfire landscape, nor does a snowfall increment prove a change in burned area. Across studies reviewed by the U.S. Government Accountability Office, estimated additional precipitation ranged from zero to 20 percent, with substantial evaluation limitations. [2,4]

2 The preventive wildfire hypothesis

The causal sequence is: eligible clouds and transport → measured ice response → incremental precipitation at the intended watershed → retained snow water, soil moisture, or fuel moisture → changed fire behavior under comparable weather and ignition conditions. Each arrow is an independent test. Primary outcomes should be seasonal precipitation or snow water equivalent and fuel moisture at predeclared locations and time lags. Secondary outcomes should include soil moisture, streamflow, relative humidity, and fire-weather indices. Wildfire ignition, rate of spread, and burned area are longer-term outcomes requiring appropriate controls and enough seasons to separate seeding from weather, fuel management, suppression, and human ignition patterns.

The timing and vegetation type matter. In some forests, moisture shortly before a fire-prone period may dampen fuels; in some grass or shrub systems, wetter antecedent conditions can also increase vegetation production and later dry fuel. The study therefore must specify vegetation strata and measure both near-term moisture gains and later fuel loads. A policy of “more rain everywhere” is not a valid inference from the mechanism. [5,6]

3 Particle size, ice effectiveness, and exposure

The “seed” is an aerosol particle, usually much smaller than an ordinary cloud droplet. A dry particle diameter must be measured separately from its wet or ice-coated size in a cloud. A 2024 laboratory study of commercial AgI-containing flare aerosol at mixed-phase temperatures of 243-267 K found that particles 200 and 400 nanometers (nm) in diameter had ice-nucleating ability comparable to pure AgI; flare-generated particles at or below 90 nm were less ice active, apparently because their non-AgI components reduced the active AgI fraction. These are observations for particular products and test conditions, not a universal minimum diameter or a prescription for release. [3]

In a 2025 natural-cloud study, the tested flare had a measured dry modal diameter of about 120 nm under one laboratory airflow condition, and most generated particles were below 200 nm in a separate clear-sky characterization. Its particle size changed with airflow. Across 16 experiments at approximately -5 to -8 degrees C, median ice-nucleated fractions ranged from 0.07% to 1.63%. This reinforces why counting emitted particles alone is insufficient: the useful quantity is the size- and temperature-resolved ice-active fraction that actually enters an eligible cloud. [7]

Research comparison	Reason to measure	Decision use
Below 90 nm	Test the weaker ice activity observed for some mixed-composition flares; quantify ultrafine number and exposure.	Do not assume smaller is more efficient or safer.
About 100-200 nm	Covers the measured ~120 nm dry mode in one field system.	Compare cloud delivery and ice activation under relevant temperatures.
About 200-400 nm	Covers a range with comparable activity to pure AgI in one lab study.	Test efficiency per emitted AgI mass and deposition, not diameter alone.

These bands are **measurement strata, not an approved target specification**. A candidate particle distribution would be selected only after material composition, emission mass, ice-active fraction, plume dispersion, and exposure modeling are measured together. Submicron particles can be inhaled; larger particles can deposit differently and may be transported less efficiently. Size alone cannot establish a human-safe formulation. Product binders, combustion products, silver speciation, total dose, worker handling, ambient concentration, and water and soil deposition all belong in the safety assessment. Published assessments have found no significant health or environmental impact at previously studied operational levels, while wider or greater use remains uncertain. [1,2,8]

4 Research design and decision gates

Stage	Work	Advance only if
A Desk and site study	Map eligible cloud hours, winds, precipitation, watersheds, vegetation, fuels, community uses, and background silver.	A meaningful seasonal window overlaps the desired moisture window.
B Laboratory	Measure dry size distribution, composition, ice-active fraction across cloud temperature, aerosol persistence and exposure potential; compare products and no-seed controls.	A candidate has repeatable activity and a defensible exposure profile.
C Observational preparation	Install radar, gauges, snow and soil sensors, fuel-moisture transects, and sampling for air, water and soil; preregister analysis.	Transport and outcome measurements can distinguish treatment from natural variability.
D Conditional field study	After authorization, randomize comparable eligible events or clusters to seeding and no seeding; use independent operations and evaluation teams.	Prespecified precipitation, exposure and fuel outcomes justify a later fire-risk study.

Eligibility must be set before the forecast event: cloud type and depth, supercooled liquid water, temperature, wind trajectory, precipitation background, aviation conditions, and target watershed. The evaluator should remain blind to assignment where feasible. Compare treated with untreated eligible events, report absolute changes and confidence intervals, and analyze downwind areas. Mechanistic evidence from radar and cloud probes should support the statistical estimate. Do not choose only successful-looking seeded storms after the fact. The World Meteorological Organization recommends objective event criteria, randomization, and physical secondary analyses. [1]

Prespecified stop conditions include failed cloud delivery, no repeatable incremental precipitation, unfavorable fuel response, environmental concentrations exceeding independently established project thresholds, unacceptable downwind effects, or lack of partner authorization. The thresholds themselves must be set by qualified agencies before any release; this paper does not invent universal safe exposure limits.

5 Human and environmental safeguards

Before any outdoor use, obtain full formulation disclosure from the contractor, assess occupational handling and combustion byproducts, establish background air and water silver levels, model plume and deposition under adverse winds, and monitor workers and nearby communities. Sampling should cover drinking-water sources, aquatic habitats, snowpack, soils, and downwind areas, with independent laboratory analysis and public reporting. Compare measured concentrations with applicable exposure and water-quality criteria for the specific chemical forms. Evaluate cumulative use across seasons rather than treating each event as isolated. [1,2,8]

Weather modification activity in the United States is subject to federal reporting to NOAA; state and local requirements vary. NOAA reporting is not itself permission to operate. Site selection should involve relevant water and land managers, fire agencies, tribes, affected communities, and aviation authorities. A proposed treatment should also consider flood and erosion exposure, water allocation, ecological effects, and where precipitation might be displaced or increased downwind. [1,9]

6 Interpretation and limits

A successful study would first show a reproducible precipitation increment in the right place and season, and then demonstrate a durable moisture response in the relevant fuels. It would still not establish a lower wildfire count without an adequately powered, multi-season comparison. Because cloud opportunity may be sparse during severe drought, a null feasibility result is useful: it would rule out the approach for that site or season. Cloud seeding should be evaluated alongside fuel treatment, ignition prevention, defensible space, prescribed fire, forecasts, and conventional water management rather than being credited with outcomes those measures produce.

The scientific claim advanced here is narrow: *preventive glaciogenic seeding might, in carefully selected landscapes and weather windows, add enough precipitation to alter antecedent fuel moisture without unacceptable exposure*. It is a testable hypothesis. The paper does not claim an operational method for stopping wildfires.

References

- [1] World Meteorological Organization. Statement on Weather Modification (2025). <https://wmo.int/content/wmo-statement-weather-modification>
- [2] U.S. Government Accountability Office. Cloud Seeding Technology: Assessing Effectiveness and Other Challenges, GAO-25-107328 (2024). <https://www.gao.gov/products/gao-25-107328>
- [3] Chen et al. Critical Size of Silver Iodide Containing Glaciogenic Cloud Seeding Particles. Geophysical Research Letters (2024), doi:10.1029/2023GL106680. <https://doi.org/10.1029/2023GL106680>
- [4] Friedrich et al. Quantifying snowfall from orographic cloud seeding. Proceedings of the National Academy of Sciences (2020), doi:10.1073/pnas.1917204117. <https://doi.org/10.1073/pnas.1917204117>
- [5] Jensen et al. The sensitivity of US wildfire occurrence to pre-season soil moisture conditions across ecosystems. Environmental Research Letters (2018), doi:10.1088/1748-9326/aa9853. <https://doi.org/10.1088/1748-9326/aa9853>
- [6] Littell et al. Climate and wildfire area burned in western U.S. ecoprovinces, 1916-2003. Ecological Applications (2009), doi:10.1890/07-1183.1. <https://doi.org/10.1890/07-1183.1>
- [7] Miller et al. Quantified ice-nucleating ability of AgI-containing seeding particles in natural clouds. Atmospheric Chemistry and Physics 25, 5387-5407 (2025). <https://doi.org/10.5194/acp-25-5387-2025>
- [8] U.S. Environmental Protection Agency. Frequent Questions on Weather Modification and Cloud Seeding. <https://www.epa.gov/geoengineering/frequent-questions>
- [9] NOAA Central Library. Weather Modification Project Reports and reporting instructions. <https://library.noaa.gov/weather-climate/weather-modification-project-reports>