

# Intro to Proofs: HW05

Aamod Varma

September 24, 2024

## 9.4

**Proof.** If the polynomial is prime  $\forall n \in \mathbb{N}$  then  $n^2 + 17n + 17$  has only 1 and itself as its factors.

However let us take  $n = 17$  we have,  $17^2 + 17^2 + 17 = 17(17 + 17 + 1) = 17 \times 35$   
We see that the polynomial has factors 17 and 35 in this case (also note that 35 can be further factored). So this shows us that  $n^2 + 17n + 17$  is not prime  $\forall n \in \mathbb{N}$

So the statement is false.  $\square$

## 9.9

**Proof.** Let  $A = \{1, 2\}$  and  $B = \{2\}$ . With this we have  $A - B = \{1\}$ .

$$P(A) = \{\{1\}, \{2\}, \{1, 2\}, \phi\}$$

$$P(B) = \{\{2\}, \phi\}$$

$$P(A) - P(B) = \{\{1\}, \{1, 2\}\}$$

$$P(A - B) = \{\{1\}, \phi\}$$

It is easy to see that  $P(A) - P(B) \not\subseteq P(A - B)$

So the statement is false.  $\square$

## 9.23

**Proof.** We have  $x^3 < y^3$ . We can write this as,

$$x^3 - y^3 < 0$$

$$(x - y)(x^2 + xy + y^2) < 0$$

We know that  $x^2 + xy + y^2$  is always positive as we can write it as,  $(x + \frac{y}{2})^2 + \frac{3y^2}{4}$  so we can divide both sides by it.

$$(x - y) < 0 \Rightarrow x < y$$

$\square$

## 9.34

**Proof.** We use a counter example to disprove this statement.

Consider  $A = \{1, 2, 3\}$  and  $B = \{3, 4, 5\}$  where  $A \cup B = \{1, 2, 3, 4, 5\}$

We can take  $X = \{2, 3, 4\}$ . We see that  $X \subseteq A \cup B$ . However it is not true that either  $X \subseteq A$  as  $4 \notin A$  and not true that  $X \subseteq B$  as  $2 \notin B$

$\square$

## 10.2

**Proof.** First let us consider  $n = 1$ , we have,

$$1^2 = \frac{1(1+1)(2+1)}{6} = \frac{6}{6} = 1$$

Now let us assume the statement is true for an arbitrary  $n = k$ , we have,

$$1^2 + 2^2 + \dots + k^2 = \frac{k(k+1)(2k+1)}{6}$$

We need to show just using this result that the statement holds true for  $n = k + 1$  or that,

$$1^2 + \dots + k^2 + (k+1)^2 = \frac{(k+1)(k+2)(2(k+1)+1)}{6} = \frac{(k+1)(2k^2+7k+6)}{6}$$

We can replace the first  $k$  terms in the left with the formula above and we get,

$$\begin{aligned} \frac{k(k+1)(2k+1)}{6} + (k+1)^2 &= \frac{k(k+1)(2k+1) + 6(k+1)^2}{6} \\ \frac{(k+1)(k(2k+1) + 6(k+1))}{6} &= \frac{(k+1)(2k^2+7k+6)}{6} \end{aligned}$$

We see that this is the same formula for the sum of  $k + 1$  terms that we had to show.

Hence we showed that if the statement holds true for  $n = k$  then it will also hold true for  $n = k + 1$  and by induction we can say that the statement is true for every positive integer  $n$ . □

## 10.8

**Proof.** First let us consider the case when  $n = 1$ . We get,

$$\frac{1}{2!} = 1 - \frac{1}{(n+1)!}$$

The left hand side evaluates to  $\frac{1}{2!} = \frac{1}{2}$ . And the right hand side evaluates to,

$$1 - \frac{1}{2!} = 1 - \frac{1}{2} = \frac{1}{2}$$

So we have  $\frac{1}{2!} = 1 - \frac{1}{(1+1)!} = \frac{1}{2}$  so the equality holds for  $n = 1$ .

Now let us assume the equality holds for an arbitrary  $n = k$ , we get,

$$\frac{1}{2!} + \dots + \frac{k}{(k+1)!} = 1 - \frac{1}{(k+1)!}$$

We need to show that the following equality is true given this,

$$\frac{1}{2!} + \cdots + \frac{k}{(k+1)!} + \frac{k+1}{(k+2)!} = 1 - \frac{1}{(k+2)!}$$

Using the righthand side from the equality we assumed for  $n = k$  we have the lefthand side of what we want to prove as,

$$\begin{aligned} & 1 - \frac{1}{(k+1)!} + \frac{k+1}{(k+2)!} \\ &= 1 + \frac{k+1}{(k+2)!} - \frac{k+2}{(k+1)!} = 1 - \frac{1}{(k+2)!} \end{aligned}$$

Which is the righthand side of what we want to prove.

Hence by inductino we showed that the statement holds for all  $n \in N$

□

## 10.14

**Proof.** We need to show  $5|2^n a \Rightarrow 5|a, \forall n \in N$  If  $n = 1$  we see that

$$5|2^n a = 5|a \text{ as } 2^n = 1$$

For an arbitrary  $n = k$  we have,

$$5|2^k a \Rightarrow 5|a$$

We have to show that given this, the statemetn holds for  $n = k + 1$  or,

$$5|2^{k+1} a \Rightarrow 5|a$$

If the statement holds for  $k$  then we know that  $2^k a = 5m_0 \Rightarrow a = 5n_0$  for some  $m \in Z$

Taking the lefthand side of what we have to prove we see,

$$2^{k+1} a = 5m_1$$

First we start with  $2^k a = 5m_0$ . Multiplifying both sides by 2 we get,

$$2^{k+1} a = 10m_0 = 5(2m_0) = 5m_1$$

However notice that we have the same  $a$  that we already assumed is divisbile in the case where  $n = k$ . Hence we show that even in the case where  $n = k + 1$  a is still divisbile by 5.

□

## 10.17

**Proof.** Let us first check for  $n = 2$ . We have,

$$\overline{A_1 \cap A_2} = \overline{A_1} \cup \overline{A_2}$$

which is true by using demorgans law.

Or in other words if  $x \notin A_1 \cap A_2$  means that  $x \notin A_1$  or  $x \notin A_2$  which means that  $x \in \overline{A_1} \cup \overline{A_2}$

Let us now assume it is true for an arbitrary  $n = k$  we have,

$$\overline{A_1 \cap \dots \cap A_k} = \overline{A_1} \cup \dots \cup \overline{A_k}$$

we need to show that the following follows,

$$\overline{A_1 \cap \dots \cap A_k \cap A_{k+1}} = \overline{A_1} \cup \dots \cup \overline{A_k} \cup \overline{A_{k+1}}$$

Consider the intersectino of  $A_1 \cap \dots \cap A_k = A_0$ . Now we have because of demorgans law (we also just showed it above for the case of  $n = 2$ ),

$$\begin{aligned} &= \overline{A_0 \cap A_{k+1}} = \overline{A_0} \cup \overline{A_{k+1}} \\ &= \overline{A_1 \cap \dots \cap A_k} \cup \overline{A_{k+1}} \end{aligned}$$

Now from our assumption we can write this as,

$$= \overline{A_1} \cup \dots \cup \overline{A_k} \cup \overline{A_{k+1}}$$

Which is the right hand side for the case of  $n = k + 1$ .

Hence by induction we have shown that the satement is true for all  $n \geq 2$   $\square$

## 10.21

Let us take the case for  $n = 1$ . We have,

$$\frac{1}{1} \leq 2 - \frac{1}{1} = 2$$

Which is obviously true.

Now let us assume the case for  $n = k$ . We get,

$$\frac{1}{1} + \dots + \frac{1}{k^2} \leq 2 - \frac{1}{k}$$

We need to show that the case for  $n = k + 1$  follows from this, or,

$$\frac{1}{1} + \dots + \frac{1}{k^2} + \frac{1}{(k+1)^2} \leq 2 - \frac{1}{k+1}$$

Let us take our inequality for  $n = k$  and add the term  $\frac{1}{(k+1)^2}$  on both sides, we get,

$$\frac{1}{1} + \dots + \frac{1}{k^2} + \frac{1}{(k+1)^2} \leq 2 - \frac{1}{k} + \frac{1}{(k+1)^2}$$

Now let us rearrange the terms in the right in this inequality, we have,

$$2 - \left( \frac{1}{k} - \frac{1}{(k+1)^2} \right) = 2 - \left( \frac{k^2 + k + 1}{k(k+1)^2} \right)$$

It is obvious that,

$$2 - \left( \frac{k^2 + k + 1}{k(k+1)^2} \right) = 2 - \left( \frac{k^2 + k}{k(k+1)^2} \right) - \frac{1}{k(k+1)^2} \leq 2 - \left( \frac{k^2 + k}{k(k+1)^2} \right) = 2 - \frac{1}{(k+1)}$$

So we have shown from the case  $n = k$  that,

$$\frac{1}{1} + \dots + \frac{1}{(k+1)^2} \leq 2 - \frac{1}{(k+1)}$$

which is the case for  $n = k + 1$ . Hence by induction we can say that our inequality is true for all  $n \in \mathbb{N}$

## 10.30

**Proof.** Let us start with the case for  $n = 1$ . We have,

$$\begin{aligned} F_1 &= \frac{\left(\frac{1+\sqrt{5}}{2}\right)^1 - \left(\frac{1-\sqrt{5}}{2}\right)^1}{\sqrt{5}} \\ &= \frac{2\sqrt{5}}{\sqrt{5}} = 2 \end{aligned}$$

which is true.

Now let us assume the case for  $n = k$  this means that,  $F_{n-2} + F_{n-1} = F_n$  or that,

$$F_k = \frac{\left(\frac{1+\sqrt{5}}{2}\right)^{k-2} - \left(\frac{1-\sqrt{5}}{2}\right)^{k-2}}{\sqrt{5}} + \frac{\left(\frac{1+\sqrt{5}}{2}\right)^{k-1} - \left(\frac{1-\sqrt{5}}{2}\right)^{k-1}}{\sqrt{5}} = \frac{\left(\frac{1+\sqrt{5}}{2}\right)^k - \left(\frac{1-\sqrt{5}}{2}\right)^k}{\sqrt{5}}$$

Now we need to show that it holds for  $n = k + 1$  or that,

$$F_{k+1} = \frac{\left(\frac{1+\sqrt{5}}{2}\right)^{k-1} - \left(\frac{1-\sqrt{5}}{2}\right)^{k-1}}{\sqrt{5}} + \frac{\left(\frac{1+\sqrt{5}}{2}\right)^k - \left(\frac{1-\sqrt{5}}{2}\right)^k}{\sqrt{5}} = \frac{\left(\frac{1+\sqrt{5}}{2}\right)^{k+1} - \left(\frac{1-\sqrt{5}}{2}\right)^{k+1}}{\sqrt{5}}$$

We have  $F_{k+1} = F_k + F_{k-1}$ . But we have an expression for  $F_k$  from our assumption. Putting that in here we get,

$$\begin{aligned} &= \frac{\left(\frac{1+\sqrt{5}}{2}\right)^{k-2} - \left(\frac{1-\sqrt{5}}{2}\right)^{k-2}}{\sqrt{5}} + \frac{\left(\frac{1+\sqrt{5}}{2}\right)^{k-1} - \left(\frac{1-\sqrt{5}}{2}\right)^{k-1}}{\sqrt{5}} + \frac{\left(\frac{1+\sqrt{5}}{2}\right)^{k-1} - \left(\frac{1-\sqrt{5}}{2}\right)^{k-1}}{\sqrt{5}} \\ &= \frac{\left(\frac{1+\sqrt{5}}{2}\right)^{k-2} - \left(\frac{1-\sqrt{5}}{2}\right)^{k-2}}{\sqrt{5}} + \frac{2\left(\frac{1+\sqrt{5}}{2}\right)^{k-1} - 2\left(\frac{1-\sqrt{5}}{2}\right)^{k-1}}{\sqrt{5}} \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{\sqrt{5}} \left( \left( \frac{1+\sqrt{5}}{2} \right)^{k-2} - \left( \frac{1-\sqrt{5}}{2} \right)^{k-2} + 2 \left( \frac{1+\sqrt{5}}{2} \right)^{k-1} - 2 \left( \frac{1-\sqrt{5}}{2} \right)^{k-1} \right) \\
&= \frac{1}{\sqrt{5}} \left( \frac{\left( \frac{1+\sqrt{5}}{2} \right)^{k+1}}{\frac{(1+\sqrt{5})^3}{2^3}} + \frac{2 \left( \frac{1+\sqrt{5}}{2} \right)^{k+1}}{\frac{(1+\sqrt{5})^2}{2^2}} - \frac{\left( \frac{1-\sqrt{5}}{2} \right)^{k+1}}{\frac{(1-\sqrt{5})^3}{2^3}} - \frac{2 \left( \frac{1-\sqrt{5}}{2} \right)^{k+1}}{\frac{(1-\sqrt{5})^2}{2^2}} \right) \\
&= \frac{1}{\sqrt{5}} \left( \left( \frac{1+\sqrt{5}}{2} \right)^{k+1} \left( \frac{1}{\left( \frac{1+\sqrt{5}}{2} \right)^3} + \frac{2}{\left( \frac{1+\sqrt{5}}{2} \right)^2} \right) - \left( \frac{1-\sqrt{5}}{2} \right)^{k+1} \left( \frac{1}{\left( \frac{1-\sqrt{5}}{2} \right)^3} + \frac{2}{\left( \frac{1-\sqrt{5}}{2} \right)^2} \right) \right)
\end{aligned}$$

Simplifying this expression we have,

$$\begin{aligned}
&= \frac{1}{\sqrt{5}} \left( \left( \frac{1+\sqrt{5}}{2} \right)^{k+1} \left( \frac{8(2+\sqrt{5})}{(1+\sqrt{5})^3} \right) - \left( \frac{1-\sqrt{5}}{2} \right)^{k+1} \left( \frac{8(2-\sqrt{5})}{(1-\sqrt{5})^3} \right) \right) \\
&= \frac{1}{\sqrt{5}} \left( \left( \frac{1+\sqrt{5}}{2} \right)^{k+1} \binom{1}{1} - \left( \frac{1-\sqrt{5}}{2} \right)^{k+1} \binom{1}{1} \right) \\
&= \frac{\left( \frac{1+\sqrt{5}}{2} \right)^{k+1} - \left( \frac{1-\sqrt{5}}{2} \right)^{k+1}}{\sqrt{5}}
\end{aligned}$$

Which is what we had to show for the case of  $n = k + 1$

□

## 10.35

Let us take the case for  $n = 2$  and  $k = 1$ . For the statement to be true we have  $\binom{n}{k}$  is even.

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{2!}{1!(1!)} = \frac{2}{1} = 2 \text{ which is even}$$

Now let us perform induction on both  $n$  and  $k$  independently. First let us fix  $n$  and perform induction on  $k$ . Let  $n = 2m$  for some integer  $m \in \mathbb{N}$ . Assume the case for  $k = 2k' + 1$  is true. This means that,

$$\binom{2m}{2k' + 1} = \frac{(2m)!}{(2k' + 1)!(2m - 2k' - 1)!} \text{ is even}$$

Now we need to show that it is also true for the next odd integer,  $k' + 1$  or  $k = 2k' + 3$

$$\binom{2m}{2k' + 3} = \frac{(2m)!}{(2k' + 3)!(2m - 2k' - 3)!}$$

We can write

$$(2k' + 3)! = (2k' + 1)!(2k' + 2)(2k' + 3)$$

and

$$(2m - 2k' - 3)! = \frac{(2m - 2k' - 1)!}{(2m - 2k' - 1)(2m - 2k' - 2)}$$

So we get,

$$\binom{2m}{2k' + 3} = \frac{(2m)!}{(2k' + 1)!(2m - 2k' - 1)!} \frac{(2m - 2k' - 1)(2m - 2k' - 2)}{(2k' + 2)(2k' + 3)}$$