

Implementation of the Second-Order Nédélec Curl-Conforming Prismatic Element for Computational Electromagnetics

Adrián Amor Martín, Luis Emilio García Castillo,
Daniel García Doñoro

Universidad Carlos III de Madrid
XXX Symposium Nacional de la Unión Científica Internacional de Radio

4 de Septiembre de 2015

Table of contents

1. Introduction

- 1.1 Motivation
- 1.2 Goals

2. Triangular Prismatic Element

- 2.1 Curl-Conforming Elements
- 2.2 Space of Basis Functions
- 2.3 Degrees of Freedom
- 2.4 Other considerations

3. Numerical results

- 3.1 Method of Manufactured Solutions
- 3.2 Comparison

4. Conclusions

- 4.1 Conclusions

Table of contents

1. Introduction

- 1.1 Motivation
- 1.2 Goals

2. Triangular Prismatic Element

- 2.1 Curl-Conforming Elements
- 2.2 Space of Basis Functions
- 2.3 Degrees of Freedom
- 2.4 Other considerations

3. Numerical results

- 3.1 Method of Manufactured Solutions
- 3.2 Comparison

4. Conclusions

- 4.1 Conclusions

Motivation: Why triangular prisms?

- ▶ Planar structures and waveguides.
 - ▶ Unstructured meshing: too many tetrahedra.

Motivation: Why triangular prisms?

- ▶ Planar structures and waveguides.
 - ▶ Unstructured meshing: too many tetrahedra.
 - ▶ Structured meshing: very deformed tetrahedra.

Motivation: Why triangular prisms?

- ▶ Planar structures and waveguides.
 - ▶ Unstructured meshing: too many tetrahedra.
 - ▶ Structured meshing: very deformed tetrahedra.
 - ▶ Structured meshing with hexahedra.

Motivation: Why triangular prisms?

- ▶ Planar structures and waveguides.
 - ▶ Unstructured meshing: too many tetrahedra.
 - ▶ Structured meshing: very deformed tetrahedra.
 - ▶ Structured meshing with hexahedra.
 - ▶ Extrude from 2D meshings with triangles: triangular prismatic elements.

Motivation: Why triangular prisms?

- ▶ Planar structures and waveguides.
 - ▶ Unstructured meshing: too many tetrahedra.
 - ▶ Structured meshing: very deformed tetrahedra.
 - ▶ Structured meshing with hexahedra.
 - ▶ Extrude from 2D meshings with triangles: triangular prismatic elements.
- ▶ Triangular prismatic elements: a combination of hexahedra and tetrahedra.

Motivation: Why triangular prisms?

- ▶ Planar structures and waveguides.
 - ▶ Unstructured meshing: too many tetrahedra.
 - ▶ Structured meshing: very deformed tetrahedra.
 - ▶ Structured meshing with hexahedra.
 - ▶ Extrude from 2D meshings with triangles: triangular prismatic elements.
- ▶ Triangular prismatic elements: a combination of hexahedra and tetrahedra.
- ▶ Fewer unknowns.

Motivation: Why triangular prisms?

- ▶ Planar structures and waveguides.
 - ▶ Unstructured meshing: too many tetrahedra.
 - ▶ Structured meshing: very deformed tetrahedra.
 - ▶ Structured meshing with hexahedra.
 - ▶ Extrude from 2D meshings with triangles: triangular prismatic elements.
- ▶ Triangular prismatic elements: a combination of hexahedra and tetrahedra.
- ▶ Fewer unknowns.

Motivation: Why another triangular prism?

- ▶ Other approaches:
 - ▶ Volakis, 1997: tangential continuity.

Motivation: Why another triangular prism?

- ▶ Other approaches:
 - ▶ Volakis, 1997: tangential continuity.
 - ▶ Graglia, 1998.

Motivation: Why another triangular prism?

- ▶ Other approaches:
 - ▶ Volakis, 1997: tangential continuity.
 - ▶ Graglia, 1998.
 - ▶ Jiao, 2009: explicit orthogonalization.

Motivation: Why another triangular prism?

- ▶ Other approaches:
 - ▶ Volakis, 1997: tangential continuity.
 - ▶ Graglia, 1998.
 - ▶ Jiao, 2009: explicit orthogonalization.
- ▶ Systematic approach:

Motivation: Why another triangular prism?

- ▶ Other approaches:
 - ▶ Volakis, 1997: tangential continuity.
 - ▶ Graglia, 1998.
 - ▶ Jiao, 2009: explicit orthogonalization.
- ▶ Systematic approach:
 - ▶ Known space of functions.

Motivation: Why another triangular prism?

- ▶ Other approaches:
 - ▶ Volakis, 1997: tangential continuity.
 - ▶ Graglia, 1998.
 - ▶ Jiao, 2009: explicit orthogonalization.
- ▶ Systematic approach:
 - ▶ Known space of functions.
 - ▶ *A priori* definition of degrees of freedom as functionals.

Motivation: Why another triangular prism?

- ▶ Other approaches:
 - ▶ Volakis, 1997: tangential continuity.
 - ▶ Graglia, 1998.
 - ▶ Jiao, 2009: explicit orthogonalization.
- ▶ Systematic approach:
 - ▶ Known space of functions.
 - ▶ *A priori* definition of degrees of freedom as functionals.
 - ▶ Basis functions as dual basis with respect to those degrees of freedom.

Motivation: Why another triangular prism?

- ▶ Other approaches:
 - ▶ Volakis, 1997: tangential continuity.
 - ▶ Graglia, 1998.
 - ▶ Jiao, 2009: explicit orthogonalization.
- ▶ Systematic approach:
 - ▶ Known space of functions.
 - ▶ *A priori* definition of degrees of freedom as functionals.
 - ▶ Basis functions as dual basis with respect to those degrees of freedom.

Goals

- Definition of the element.
- Implementation of the element in HOFEM (*Higher Order Finite Element Method*).

Goals

- ▶ Definition of the element.
- ▶ Implementation of the element in HOFEM (*Higher Order Finite Element Method*).
- ▶ Verification of the element.

Goals

- ▶ Definition of the element.
- ▶ Implementation of the element in HOFEM (*Higher Order Finite Element Method*).
- ▶ Verification of the element.
- ▶ Comparison with tetrahedra.

Goals

- ▶ Definition of the element.
- ▶ Implementation of the element in HOFEM (*Higher Order Finite Element Method*).
- ▶ Verification of the element.
- ▶ Comparison with tetrahedra.

Table of contents

1. Introduction

- 1.1 Motivation
- 1.2 Goals

2. Triangular Prismatic Element

- 2.1 Curl-Conforming Elements
- 2.2 Space of Basis Functions
- 2.3 Degrees of Freedom
- 2.4 Other considerations

3. Numerical results

- 3.1 Method of Manufactured Solutions
- 3.2 Comparison

4. Conclusions

- 4.1 Conclusions

Mixed-Order Curl-Conforming Nédélec Elements

- ▶ Finite element:
 - ▶ Domain.

Mixed-Order Curl-Conforming Nédélec Elements

- ▶ Finite element:
 - ▶ Domain.
 - ▶ Space of basis functions.

Mixed-Order Curl-Conforming Nédélec Elements

- ▶ Finite element:
 - ▶ Domain.
 - ▶ Space of basis functions.
 - ▶ Degrees of freedom.

Mixed-Order Curl-Conforming Nédélec Elements


- ▶ Finite element:
 - ▶ Domain.
 - ▶ Space of basis functions.
 - ▶ Degrees of freedom.
- ▶ Domain.

Mixed-Order Curl-Conforming Nédélec Elements

- ▶ Finite element:
 - ▶ Domain.
 - ▶ Space of basis functions.
 - ▶ Degrees of freedom.
- ▶ Domain.

Mixed-Order Curl-Conforming Nédélec Elements

- ▶ Finite element:
 - ▶ Domain.
 - ▶ Space of basis functions.
 - ▶ Degrees of freedom.
- ▶ Domain.



Space of Basis Functions

- Nédélec space of functions \mathcal{R}^k .

$$\mathcal{R}^k = \{ \bar{u} \in (P_k)^n; \epsilon^k \bar{u} = 0 \}$$

$$\mathcal{R}^k \subset (\mathcal{P}_k)^n; (\mathcal{P}_{k-1})^n \subset \mathcal{R}^k$$

Space of Basis Functions

- Nédélec space of functions \mathcal{R}^k .

$$\mathcal{R}^k = \{ \bar{u} \in (P_k)^n; \epsilon^k \bar{u} = 0 \}$$

$$\mathcal{R}^k \subset (\mathcal{P}_k)^n; (\mathcal{P}_{k-1})^n \subset \mathcal{R}^k$$

- Other definition:

Space of Basis Functions

- Nédélec space of functions \mathcal{R}^k .

$$\mathcal{R}^k = \{\bar{u} \in (P_k)^n; \epsilon^k \bar{u} = 0\}$$

$$\mathcal{R}^k \subset (P_k)^n; (\mathcal{P}_{k-1})^n \subset \mathcal{R}^k$$

- Other definition:

$$\mathcal{R}^k = (\mathcal{P}_{k-1})^n \oplus \mathcal{S}^k$$

$$\bar{u} \in \mathcal{S}^k \Leftrightarrow \bar{r} \cdot \bar{u} = 0; \bar{r} = (x, y, z)$$

Space of Basis Functions

- Nédélec space of functions \mathcal{R}^k .

$$\mathcal{R}^k = \{\bar{u} \in (P_k)^n; \epsilon^k \bar{u} = 0\}$$

$$\mathcal{R}^k \subset (P_k)^n; (\mathcal{P}_{k-1})^n \subset \mathcal{R}^k$$

- Other definition:

$$\mathcal{R}^k = (\mathcal{P}_{k-1})^n \oplus \mathcal{S}^k$$

$$\bar{u} \in \mathcal{S}^k \Leftrightarrow \bar{r} \cdot \bar{u} = 0; \bar{r} = (x, y, z)$$

- Different approximation order in different directions.

Space of Basis Functions

- Nédélec space of functions \mathcal{R}^k .

$$\mathcal{R}^k = \{\bar{u} \in (P_k)^n; \epsilon^k \bar{u} = 0\}$$

$$\mathcal{R}^k \subset (P_k)^n; (\mathcal{P}_{k-1})^n \subset \mathcal{R}^k$$

- Other definition:

$$\mathcal{R}^k = (\mathcal{P}_{k-1})^n \oplus \mathcal{S}^k$$

$$\bar{u} \in \mathcal{S}^k \Leftrightarrow \bar{r} \cdot \bar{u} = 0; \bar{r} = (x, y, z)$$

- Different approximation order in different directions.

Space of Basis Functions

- Nédélec space of functions \mathcal{R}^k .

$$\mathcal{R}^k = \{\bar{\mathbf{u}} \in (P_k)^n; \epsilon^k \bar{\mathbf{u}} = 0\}$$


$$\mathcal{R}^k \subset (\mathcal{P}_k)^n; (\mathcal{P}_{k-1})^n \subset \mathcal{R}^k$$

- Other definition:

$$\mathcal{R}^k = (\mathcal{P}_{k-1})^n \oplus \mathcal{S}^k$$

$$\bar{\mathbf{u}} \in \mathcal{S}^k \Leftrightarrow \bar{\mathbf{r}} \cdot \bar{\mathbf{u}} = 0; \bar{\mathbf{r}} = (x, y, z)$$

- Different approximation order in different directions.



Space construction

- ▶ Tensor product between triangle and segment.

$$\mathcal{P}_k^{\text{prism}} = (\mathcal{R}^k(\hat{T}) \otimes \mathcal{P}_k(\hat{I})) \times (\mathcal{P}_k(\hat{T}) \otimes \mathcal{P}_{k-1}(\hat{I}))$$

- ▶ Space dimension.

Space construction

- Tensor product between triangle and segment.

$$\mathcal{P}_k^{\text{prism}} = (\mathcal{R}^k(\hat{T}) \otimes \mathcal{P}_k(\hat{I})) \times (\mathcal{P}_k(\hat{T}) \otimes \mathcal{P}_{k-1}(\hat{I}))$$

- Space dimension.

Dimensions	$\mathcal{R}^k(\hat{T})$	$\mathcal{P}_k(\hat{I})$	$\mathcal{P}_k(\hat{T})$	$\mathcal{P}_{k-1}(\hat{I})$	Total
k	$(k+2)k$	$k+1$	$\frac{(k+1)(k+2)}{2}$	k	—
$k=1$	3	2	3	1	9
$k=2$	8	3	6	2	36
$k=3$	15	4	10	3	90
$k=4$	24	5	15	4	180

Space construction

- Tensor product between triangle and segment.

$$\mathcal{P}_k^{\text{prism}} = (\mathcal{R}^k(\hat{T}) \otimes \mathcal{P}_k(\hat{I})) \times (\mathcal{P}_k(\hat{T}) \otimes \mathcal{P}_{k-1}(\hat{I}))$$

- Space dimension.

Dimensions	$\mathcal{R}^k(\hat{T})$	$\mathcal{P}_k(\hat{I})$	$\mathcal{P}_k(\hat{T})$	$\mathcal{P}_{k-1}(\hat{I})$	Total
k	$(k+2)k$	$k+1$	$\frac{(k+1)(k+2)}{2}$	k	—
$k=1$	3	2	3	1	9
$k=2$	8	3	6	2	36
$k=3$	15	4	10	3	90
$k=4$	24	5	15	4	180

Space definition

- First order.

$$\mathcal{P}_1^{\text{prism}} \equiv \mathbf{N}_i (i = 1, \dots, 9) = \left\{ \begin{array}{l} a_1^{(i)} + a_2^{(i)} z + C^{(i)} y + D^{(i)} yz \\ b_1^{(i)} + b_2^{(i)} z - C^{(i)} x - D^{(i)} xz \\ c_1^{(i)} + c_2^{(i)} x + c_3^{(i)} y \end{array} \right\}$$

- Second order.

Space definition

- First order.

$$\mathcal{P}_1^{\text{prism}} \equiv \mathbf{N}_i (i = 1, \dots, 9) = \left\{ \begin{array}{l} a_1^{(i)} + a_2^{(i)} z + C^{(i)} y + D^{(i)} yz \\ b_1^{(i)} + b_2^{(i)} z - C^{(i)} x - D^{(i)} xz \\ c_1^{(i)} + c_2^{(i)} x + c_3^{(i)} y \end{array} \right\}$$

- Second order.

$$\mathcal{P}_2^{\text{prism}} \equiv \mathbf{N}_i (i = 1, \dots, 36) = \left\{ \begin{array}{l} a_1^{(i)} + a_2^{(i)} x + a_3^{(i)} y + a_4^{(i)} z + a_5^{(i)} xz + a_6^{(i)} yz + a_7^{(i)} z^2 + a_8^{(i)} xz^2 + \dots \\ \dots + a_9^{(i)} yz^2 + C^{(i)} y^2 + D^{(i)} xy + E^{(i)} y^2 z + F^{(i)} xyz + G^{(i)} y^2 z^2 + H^{(i)} xyz^2 \\ b_1^{(i)} + b_2^{(i)} x + b_3^{(i)} y + b_4^{(i)} z + b_5^{(i)} xz + b_6^{(i)} yz + b_7^{(i)} z^2 + b_8^{(i)} xz^2 + \dots \\ \dots + b_9^{(i)} yz^2 - C^{(i)} xy - D^{(i)} x^2 - E^{(i)} xyz - F^{(i)} x^2 z - G^{(i)} xyz^2 - H^{(i)} x^2 z^2 \\ c_1^{(i)} + c_2^{(i)} x + c_3^{(i)} y + c_4^{(i)} x^2 + c_5^{(i)} y^2 + c_6^{(i)} xy + c_7^{(i)} z + c_8^{(i)} xz + \dots \\ \dots + c_9^{(i)} yz + c_{10}^{(i)} x^2 z + c_{11}^{(i)} y^2 z + c_{12}^{(i)} xyz \end{array} \right\}$$

Space definition

- First order.

$$\mathcal{P}_1^{\text{prism}} \equiv \mathbf{N}_i (i = 1, \dots, 9) = \left\{ \begin{array}{l} a_1^{(i)} + a_2^{(i)} z + C^{(i)} y + D^{(i)} yz \\ b_1^{(i)} + b_2^{(i)} z - C^{(i)} x - D^{(i)} xz \\ c_1^{(i)} + c_2^{(i)} x + c_3^{(i)} y \end{array} \right\}$$

- Second order.

$$\mathcal{P}_2^{\text{prism}} \equiv \mathbf{N}_i (i = 1, \dots, 36) = \left\{ \begin{array}{l} a_1^{(i)} + a_2^{(i)} x + a_3^{(i)} y + a_4^{(i)} z + a_5^{(i)} xz + a_6^{(i)} yz + a_7^{(i)} z^2 + a_8^{(i)} xz^2 + \dots \\ \dots + a_9^{(i)} yz^2 + C^{(i)} y^2 + D^{(i)} xy + E^{(i)} y^2 z + F^{(i)} xyz + G^{(i)} y^2 z^2 + H^{(i)} xyz^2 \\ b_1^{(i)} + b_2^{(i)} x + b_3^{(i)} y + b_4^{(i)} z + b_5^{(i)} xz + b_6^{(i)} yz + b_7^{(i)} z^2 + b_8^{(i)} xz^2 + \dots \\ \dots + b_9^{(i)} yz^2 - C^{(i)} xy - D^{(i)} x^2 - E^{(i)} xyz - F^{(i)} x^2 z - G^{(i)} xyz^2 - H^{(i)} x^2 z^2 \\ c_1^{(i)} + c_2^{(i)} x + c_3^{(i)} y + c_4^{(i)} x^2 + c_5^{(i)} y^2 + c_6^{(i)} xy + c_7^{(i)} z + c_8^{(i)} xz + \dots \\ \dots + c_9^{(i)} yz + c_{10}^{(i)} x^2 z + c_{11}^{(i)} y^2 z + c_{12}^{(i)} xyz \end{array} \right\}$$

Definition of the degrees of freedom

- Edges.

$$g(\mathbf{u}) = \int_e (\mathbf{u} \cdot \hat{\tau}) q \, dl, \forall q \in P_1(e)$$

Definition of the degrees of freedom

- ▶ Edges.

$$g(\mathbf{u}) = \int_e (\mathbf{u} \cdot \hat{\boldsymbol{\tau}}) q \, dl, \forall q \in P_1(e)$$

- ▶ Triangular faces.

Definition of the degrees of freedom

- ▶ Edges.

$$g(\mathbf{u}) = \int_e (\mathbf{u} \cdot \hat{\boldsymbol{\tau}}) q \, dl, \forall q \in P_1(e)$$

- ▶ Triangular faces.

$$g(\mathbf{u}) = \int_{f_t} (\mathbf{u} \times \hat{\mathbf{n}}) \cdot \mathbf{q} \, ds, \forall \mathbf{q} \in \mathbf{P}_0(f_t)$$

Definition of the degrees of freedom

- ▶ Edges.

$$g(\mathbf{u}) = \int_e (\mathbf{u} \cdot \hat{\boldsymbol{\tau}}) q \, dl, \forall q \in P_1(e)$$

- ▶ Triangular faces.

$$g(\mathbf{u}) = \int_{f_t} (\mathbf{u} \times \hat{\mathbf{n}}) \cdot \mathbf{q} \, ds, \forall \mathbf{q} \in \mathbf{P}_0(f_t)$$

- ▶ Quadrangular faces.

Definition of the degrees of freedom

- ▶ Edges.

$$g(\mathbf{u}) = \int_e (\mathbf{u} \cdot \hat{\boldsymbol{\tau}}) q \, dl, \forall q \in P_1(e)$$

- ▶ Triangular faces.

$$g(\mathbf{u}) = \int_{f_t} (\mathbf{u} \times \hat{\mathbf{n}}) \cdot \mathbf{q} \, ds, \forall \mathbf{q} \in \mathbf{P}_0(f_t)$$

- ▶ Quadrangular faces.

$$g(\mathbf{u}) = \int_{f_q} (\hat{\mathbf{n}} \times \mathbf{u}) \cdot \mathbf{q} \, ds, \forall \mathbf{q} = (q_1, q_2); q_1 \in \mathcal{Q}_{0,1}; q_2 \in \mathcal{Q}_{1,0}$$

Definition of the degrees of freedom

- ▶ Edges.

$$g(\mathbf{u}) = \int_e (\mathbf{u} \cdot \hat{\boldsymbol{\tau}}) q \, dl, \forall q \in P_1(e)$$

- ▶ Triangular faces.

$$g(\mathbf{u}) = \int_{f_t} (\mathbf{u} \times \hat{\mathbf{n}}) \cdot \mathbf{q} \, ds, \forall \mathbf{q} \in \mathbf{P}_0(f_t)$$

- ▶ Quadrangular faces.

$$g(\mathbf{u}) = \int_{f_q} (\hat{\mathbf{n}} \times \mathbf{u}) \cdot \mathbf{q} \, ds, \forall \mathbf{q} = (q_1, q_2); q_1 \in \mathcal{Q}_{0,1}; q_2 \in \mathcal{Q}_{1,0}$$

- ▶ Volume.

Definition of the degrees of freedom

- ▶ Edges.

$$g(\mathbf{u}) = \int_e (\mathbf{u} \cdot \hat{\boldsymbol{\tau}}) q \, dl, \forall q \in P_1(e)$$

- ▶ Triangular faces.

$$g(\mathbf{u}) = \int_{f_t} (\mathbf{u} \times \hat{\mathbf{n}}) \cdot \mathbf{q} \, ds, \forall \mathbf{q} \in \mathbf{P}_0(f_t)$$

- ▶ Quadrangular faces.

$$g(\mathbf{u}) = \int_{f_q} (\hat{\mathbf{n}} \times \mathbf{u}) \cdot \mathbf{q} \, ds, \forall \mathbf{q} = (q_1, q_2); q_1 \in \mathcal{Q}_{0,1}; q_2 \in \mathcal{Q}_{1,0}$$

- ▶ Volume.

$$g(\mathbf{u}) = \int_v \mathbf{u} \cdot \mathbf{q} \, dV, \forall \mathbf{q} \in \mathbf{P}_0(f_t)$$

Definition of the degrees of freedom

- ▶ Edges.

$$g(\mathbf{u}) = \int_e (\mathbf{u} \cdot \hat{\boldsymbol{\tau}}) q \, dl, \forall q \in P_1(e)$$

- ▶ Triangular faces.

$$g(\mathbf{u}) = \int_{f_t} (\mathbf{u} \times \hat{\mathbf{n}}) \cdot \mathbf{q} \, ds, \forall \mathbf{q} \in \mathbf{P}_0(f_t)$$


- ▶ Quadrangular faces.

$$g(\mathbf{u}) = \int_{f_q} (\hat{\mathbf{n}} \times \mathbf{u}) \cdot \mathbf{q} \, ds, \forall \mathbf{q} = (q_1, q_2); q_1 \in \mathcal{Q}_{0,1}; q_2 \in \mathcal{Q}_{1,0}$$

- ▶ Volume.

$$g(\mathbf{u}) = \int_v \mathbf{u} \cdot \mathbf{q} \, dV, \forall \mathbf{q} \in \mathbf{P}_0(f_t)$$

Master element



Other considerations (i)

- Discretization: choice of \mathbf{q} .

Dual basis

$$g_i(\mathbf{N}_j) = \delta_{ij}$$

Other considerations (i)

- Discretization: choice of \mathbf{q} .

Dual basis

$$g_i(\mathbf{N}_j) = \delta_{ij}$$

- a_1, a_2, \dots as unknowns.

Other considerations (i)

- Discretization: choice of \mathbf{q} .

Dual basis

$$g_i(\mathbf{N}_j) = \delta_{ij}$$

- a_1, a_2, \dots as unknowns.

a_1	a_2	a_3	a_4	a_5	a_6	a_7	a_8	a_9	...
4	-6	-12	-16	24	48	12	-18	-36	...
-2	6	2	8	-24	-8	-6	18	6	...
0	0	2	0	0	-8	0	0	6	...
0	0	4	0	0	-16	0	0	12	...
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮

Other considerations (i)

- Discretization: choice of \mathbf{q} .

Dual basis

$$g_i(\mathbf{N}_j) = \delta_{ij}$$

- a_1, a_2, \dots as unknowns.

a_1	a_2	a_3	a_4	a_5	a_6	a_7	a_8	a_9	...
4	-6	-12	-16	24	48	12	-18	-36	...
-2	6	2	8	-24	-8	-6	18	6	...
0	0	2	0	0	-8	0	0	6	...
0	0	4	0	0	-16	0	0	12	...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\ddots

Other considerations (ii)

- ▶ Local definition of $\hat{\tau}$, $\hat{\mathbf{n}}$, \mathbf{q} .
- ▶ Use of a master element:

$$\mathbf{u} = [\mathcal{J}]^{-1} \hat{\mathbf{u}}$$

Other considerations (ii)


- ▶ Local definition of $\hat{\tau}$, \hat{n} , \mathbf{q} .
- ▶ Use of a master element:

$$\mathbf{u} = [\mathcal{J}]^{-1} \hat{\mathbf{u}}$$

Other considerations (ii)

- Local definition of $\hat{\tau}$, \hat{n} , \mathbf{q} .
- Use of a master element:

$$\mathbf{u} = [J]^{-1} \hat{\mathbf{u}}$$



Other considerations (& iii)

- ▶ Two versions:

- ▶ VC.

Other considerations (& iii)

► Two versions:


- vc .
- vq .

Other considerations (& iii)

- ▶ Two versions:
 - ▶ vc .
 - ▶ vq .
- ▶ Triangular face.

Other considerations (& iii)


- ▶ Two versions:
 - ▶ vc.
 - ▶ vq.
- ▶ Triangular face.




- ▶ Quadrangular face.

Other considerations (& iii)


- ▶ Two versions:
 - ▶ *vc*.
 - ▶ *vq*.
- ▶ Triangular face.
- ▶ Quadrangular face.



Basis functions on triangular faces




N_3




N_{19}

Basis functions on quadrangular faces



N_{13}



N_{24}

Table of contents

1. Introduction

- 1.1 Motivation
- 1.2 Goals

2. Triangular Prismatic Element

- 2.1 Curl-Conforming Elements
- 2.2 Space of Basis Functions
- 2.3 Degrees of Freedom
- 2.4 Other considerations

3. Numerical results


- 3.1 Method of Manufactured Solutions
- 3.2 Comparison

4. Conclusions


- 4.1 Conclusions

Method of Manufactured Solutions


- ▶ $\nabla \times \frac{1}{f_r} \nabla \times \mathbf{u} - k_0^2 \mathbf{g}_r \mathbf{u} = \Psi$.
- ▶ HOFEM: Monomials ($xyz^2, -xz^2, xyz$).



Analytical solution



Code solution




Error

Method of Manufactured Solutions


- ▶ $\nabla \times \frac{1}{f_r} \nabla \times \mathbf{u} - k_0^2 \mathbf{g}_r \mathbf{u} = \Psi$.
- ▶ HOFEM: Monomials $(xyz^2, -xz^2, xyz)$.
- ▶ HOFEM: Planewave.

Comparison between different elements and angles of incidence, ϕ -polarization




Comparison (i)

- Structure to be simulated.



Comparison (i)

- Structure to be simulated.



Comparison (i)

- Structure to be simulated.
- Comparison.

Element	Elements	Unknowns	Simulation time (s)
Tetrahedra	22797	157240	2637.972
Prism	5220	83090	1061.215

Comparison (i)

- ▶ Structure to be simulated.
- ▶ Comparison.

Element	Elements	Unknowns	Simulation time (s)
Tetrahedra	22797	157240	2637.972
Prism	5220	83090	1061.215


Comparison (& ii)

- Structure to be simulated.



Comparison (& ii)

- Structure to be simulated.



Comparison (& ii)

- Structure to be simulated.
- Comparison.

—	Tetrahedra	Triangular prism	Finer mesh
Number of elements	14852	5444	138943
Number of unknowns	101828	91468	2054524
Simulation time (s)	107.894	80.746	799.843

Comparison (& ii)

- ▶ Structure to be simulated.
- ▶ Comparison.

—	Tetrahedra	Triangular prism	Finer mesh
Number of elements	14852	5444	138943
Number of unknowns	101828	91468	2054524
Simulation time (s)	107.894	80.746	799.843

Table of contents

1. Introduction

- 1.1 Motivation
- 1.2 Goals

2. Triangular Prismatic Element

- 2.1 Curl-Conforming Elements
- 2.2 Space of Basis Functions
- 2.3 Degrees of Freedom
- 2.4 Other considerations

3. Numerical results

- 3.1 Method of Manufactured Solutions
- 3.2 Comparison

4. Conclusions

- 4.1 Conclusions

Conclusions

- ▶ Systematic approach for designing higher-order basis functions.
- ▶ Mathematical verification of the element.

Conclusions

- ▶ Systematic approach for designing higher-order basis functions.
- ▶ Mathematical verification of the element.
- ▶ Implementation of the element in HOFEM.

Conclusions

- ▶ Systematic approach for designing higher-order basis functions.
- ▶ Mathematical verification of the element.
- ▶ Implementation of the element in HOFEM.
- ▶ Comparison between triangular prismatic elements and tetrahedra.

Conclusions


- ▶ Systematic approach for designing higher-order basis functions.
- ▶ Mathematical verification of the element.
- ▶ Implementation of the element in HOFEM.
- ▶ Comparison between triangular prismatic elements and tetrahedra.
- ▶ Acknowledgements: TEC2010-18175/TCM and TEC2013-47753-C3-2.

Conclusions

- ▶ Systematic approach for designing higher-order basis functions.
- ▶ Mathematical verification of the element.
- ▶ Implementation of the element in HOFEM.
- ▶ Comparison between triangular prismatic elements and tetrahedra.
- ▶ Acknowledgements: TEC2010-18175/TCM and TEC2013-47753-C3-2.

Dudas y preguntas

Implementation of the Second-Order Nédélec Curl-Conforming Prismatic Element for Computational Electromagnetics



Adrián Amor Martín, aamor@tsc.uc3m.es.

Universidad Carlos III de Madrid.

Grupo de Radiofrecuencia, Electromagnetismo, Microondas y Antenas.