

David Morin - Cycloidal Pendulum

Alexander Ivanov

Jan 2022

Exercise 6.29

The standard pendulum frequency of $\sqrt{g/\ell}$ holds only for small oscillations. The frequency becomes smaller as the amplitude grows. It turns out that if you want to build a pendulum whose frequency is independent of the amplitude, you should hang it from the cusp of a cycloid of a certain size, as shown in Fig. 6.30. As the string wraps partially around the cycloid, the effect is to decrease the length of string in the air, which in turn increases the frequency back up to a constant value. In more detail: A cycloid is the path taken by a point on the rim of a rolling wheel. The upside-down cycloid in Fig. 6.30 can be parameterized by $(x, y) = R(\theta - \sin \theta, -1 + \cos \theta)$, where $\theta = 0$ corresponds to the cusp. Consider a pendulum of length $4R$ hanging from the cusp, and let α be the angle the string makes with the vertical, as shown.




Fig. 6.30

(a)

In terms of α , find the value of the parameter θ associated with the point where the string leaves the cycloid.

Solution:

From the question, we have $(dx, dy) = Rd\theta(1 - \cos \theta, -\sin \theta)$.

Note that the part of the string leaving the cycloid is tangent to the cycloid at the point where it leaves. So, by basic trigonometry and our previous observation,

$$\tan \alpha = \left| \frac{dx}{dy} \right| = \frac{1 - \cos \theta}{\sin \theta} = \tan \frac{\theta}{2} \Rightarrow \theta = 2\alpha$$

(b)

In terms of α , find the length of the string touching the cycloid.

Solution:

By arc length formula, the length of the string is $\int ds$, where ds is an infinitesimal piece of string touching the cycloid. We have that $ds^2 = dx^2 + dy^2 = R^2 d\theta^2 ((1 - \cos \theta)^2 + (-\sin \theta)^2) = 2R^2 d\theta^2 (1 - \cos \theta) = 4R^2 d\theta^2 (\frac{1 - \cos \theta}{2}) = 4R^2 d\theta^2 \sin^2 \frac{\theta}{2} \Rightarrow ds = 2R d\theta \sin \frac{\theta}{2} = 4R d\alpha \sin \alpha$.

And therefore,

$$\int ds = \int_0^\alpha 4R \sin \alpha d\alpha = -4R \cos \alpha + 4R \cos 0 = 4R(1 - \cos \alpha)$$

(c)

In terms of α , find the Lagrangian.

Solution:

In general, the Lagrangian is:

$$\mathcal{L} = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2) - mgy$$

Note that the length of string in the air for given α is

$$4R - 4R(1 - \cos \alpha) = 4R \cos \alpha$$

And so, the position of the mass relative to the contact point is,

$$4R \cos \alpha (\sin \alpha, -\cos \alpha) = R(2 \sin 2\alpha, -2 - 2 \cos 2\alpha)$$

Also, since $\theta = 2\alpha$ at the contact point, the position of the contact point is,

$$R(2\alpha - \sin 2\alpha, -1 + \cos 2\alpha)$$

Thus, the position of the mass is

$$(x, y) = R(2\alpha - \sin 2\alpha, -1 + \cos 2\alpha) + R(2 \sin 2\alpha, -2 - 2 \cos 2\alpha) = R(2\alpha + \sin 2\alpha, -3 - \cos 2\alpha)$$

Then, $(\dot{x}, \dot{y}) = 2R\dot{\alpha}(1 + \cos 2\alpha, \sin 2\alpha)$ and so $\dot{x}^2 + \dot{y}^2 = 8R^2\dot{\alpha}^2(1 + \cos 2\alpha)$.

Finally, we have

$$\mathcal{L} = 4mR^2\dot{\alpha}^2(1 + \cos 2\alpha) + mgR(3 + \cos 2\alpha)$$

(d)

Show that the quantity $\sin \alpha$ undergoes simple harmonic motion with frequency $\sqrt{g/4R}$, independent of the amplitude.

Solution:

By the Euler-Lagrange Equation,

$$\frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{\alpha}} \right) = \frac{\partial \mathcal{L}}{\partial \alpha}$$

And so,

$$\begin{aligned} \frac{d}{dt} (8mR^2\dot{\alpha}(1 + \cos 2\alpha)) &= -8mR^2\dot{\alpha}^2 \sin 2\alpha - 2mgR \sin 2\alpha \\ 4R\ddot{\alpha}(1 + \cos 2\alpha) - 8R\dot{\alpha}^2 \sin 2\alpha &= -4R\dot{\alpha}^2 \sin 2\alpha - g \sin 2\alpha \\ \ddot{\alpha}(1 + \cos 2\alpha) - \dot{\alpha}^2 \sin 2\alpha &= -(g/4R) \sin 2\alpha \\ \ddot{\alpha}(2 \cos^2 \alpha) - \dot{\alpha}^2(2 \sin \alpha \cos \alpha) &= -(g/4R) 2 \sin \alpha \cos \alpha \\ \ddot{\alpha} \cos \alpha - \dot{\alpha}^2 \sin \alpha &= -(g/4R) \sin \alpha \\ \frac{d^2}{dt^2}(\sin \alpha) &= -(g/4R) \sin \alpha \\ \sin \alpha &= A \cos \left(\sqrt{\frac{g}{4R}} t + \phi \right) \end{aligned}$$

Thus, $\sin \alpha$ undergoes SHM with frequency $\sqrt{g/4R}$, independent of A .

(e)

In place of parts (c) and (d), solve the problem again by using $F = ma$. This actually gives a much quicker solution.

Solution:

We have tangential speed about the contact point: $v = (4R \cos \alpha) \dot{\alpha}$.

So,

$$F = ma \Rightarrow -mg \sin \alpha = m \frac{d}{dt}(4R \cos \alpha \dot{\alpha}) \Rightarrow -\frac{g}{4R} \sin \alpha = \frac{d^2}{dt^2}(\sin \alpha)$$

as desired.