

# LOCATING PARKS WITH A LEXICOGRAPHIC MULTIOBJECTIVE OPTIMIZATION METHOD

Under the tutelage of Shahriari

---

Alan Peral

October 27, 2017

Pomona College

## BACKGROUND INFO

---

What does this mean?


- Parks have benefits!

- Parks have benefits!
- Bogotá's master plan

- Parks have benefits!
- Bogotá's master plan
- Parks department

- Parks have benefits!
- Bogotá's master plan
- Parks department
- Competing objectives

ESSENTIALLY






# A BASIC EXAMPLE

---

## EXAMPLE



This is our  $5 \times 5$  grid city

## EXAMPLE

1	2	3	4	5
6	7	8	9	10
11	12	13	14	15
16	17	18	19	20
21	22	23	24	25


Maybe call it Pawnee.

## EXAMPLE

1	2	3	4	5
6	7	8	9	10
11	12	13	14	15
16	17	18	19	20
21	22	23	24	25


But definitely not.

## EXAMPLE




- Candidate parcels

## EXAMPLE




- Candidate parcels
- Every lot same size

## EXAMPLE



- Candidate parcels
- Every lot same size
- Service area


## EXAMPLE



Which candidate parcel will maximize geographical coverage?



## EXAMPLE



Which candidate parcel will maximize geographical coverage?

Candidate 12

$$\begin{aligned} \max f_1 &= \sum_{j \in \mathcal{J}} z_j \\ \text{subject to } z_j &\leq \sum_{i \in \mathcal{W}_j} y_i, j \in \mathcal{J} \\ |\mathcal{W}_j| z_j &\geq \sum_{i \in \mathcal{W}_j} y_i, j \in \mathcal{J} \\ z_j &\in \{0, 1\}, j \in \mathcal{J} \\ y_i &\in \{0, 1\}, i \in \mathcal{I} \end{aligned}$$

$$\max f_1 = \sum_{j \in \mathcal{J}} z_j$$

- $\mathcal{J}$  is the set of all blocks that could benefit from a new park

## EXAMPLE

1	2	3	4	5
6	7	8	9	10
11	12	13	14	15
16	17	18	19	20
21	22	23	24	25

$$\mathcal{J} = \{2, 4, 6, 7, 8, \dots, 22, 23, 24, 25\}$$

$$\max f_1 = \sum_{j \in \mathcal{J}} z_j$$

subject to  $z_j \in \{0, 1\}, j \in \mathcal{J}$

- $\mathcal{J}$  is the set of all blocks that could benefit from a new park
- $z_j$  is a binary decision variable, takes value of 1 if block  $j \in \mathcal{J}$  is covered by at least one park, 0 otherwise

## EXAMPLE

1	2	3	4	5
6	7	8	9	10
11	12	13	14	15
16	17	18	19	20
21	22	23	24	25

$$\mathcal{J} = \{2, 4, 6, 7, 8, \dots, 22, 23, 24, 25\}$$

If we chose candidate 21, then  $z_7 = 0$  and  $z_{16} = z_{17} = z_{22} = 1$

$$\begin{aligned} \max f_1 &= \sum_{j \in \mathcal{J}} z_j \\ \text{subject to } z_j &\leq \sum_{i \in \mathcal{W}_j} y_i, j \in \mathcal{J} \\ z_j &\in \{0, 1\}, j \in \mathcal{J} \\ y_i &\in \{0, 1\}, i \in \mathcal{I} \end{aligned}$$

- $\mathcal{J}$  is the set of all blocks that could benefit from a new park
- $z_j$  is a binary decision variable, takes value of 1 if block  $j \in \mathcal{J}$  is covered by at least one park, 0 otherwise
- $\mathcal{I}$  is the set of candidate parcels
- candidate parcel  $i$  belongs to the set  $\mathcal{W}_j$  if block  $j$  is serviced by parcel  $i$
- $y_i$  a binary decision variable, takes value 1 if candidate  $i$  is selected to become a park

## EXAMPLE

1	2	3	4	5
6	7	8	9	10
11	12	13	14	15
16	17	18	19	20
21	22	23	24	25

$$\mathcal{J} = \{2, 4, 6, 7, 8, \dots, 22, 23, 24, 25\}$$

$$\mathcal{I} = \{1, 3, 12, 15, 19, 21\}$$



## EXAMPLE

1	2	3	4	5
6	7	8	9	10
11	12	13	14	15
16	17	18	19	20
21	22	23	24	25

$$\mathcal{J} = \{2, 4, 6, 7, 8, \dots, 22, 23, 24, 25\}$$

$$\mathcal{I} = \{1, 3, 12, 15, 19, 21\}$$

$$\mathcal{W}_7 = \{1, 3, 12\}$$

## EXAMPLE

1	2	3	4	5
6	7	8	9	10
11	12	13	14	15
16	17	18	19	20
21	22	23	24	25

$$\mathcal{J} = \{2, 4, 6, 7, 8, \dots, 22, 23, 24, 25\}$$

$$\mathcal{I} = \{1, 3, 12, 15, 19, 21\}$$


$$\mathcal{W}_7 = \{1, 3, 12\}$$

If we chose block 21, then  $y_{21} = 1$

$$\begin{aligned} \max f_1 &= \sum_{j \in \mathcal{J}} z_j \\ \text{subject to } z_j &\leq \sum_{i \in \mathcal{W}_j} y_i, j \in \mathcal{J} \\ |\mathcal{W}_j| z_j &\geq \sum_{i \in \mathcal{W}_j} y_i, j \in \mathcal{J} \\ z_j &\in \{0, 1\}, j \in \mathcal{J} \\ y_i &\in \{0, 1\}, i \in \mathcal{I} \end{aligned}$$

- Guarantees that if block  $j$  is covered, then at least one candidate parcel covering it has been selected as a park
- If block  $j$  is not covered, then none of the candidate parcels serving it should be selected

## EXAMPLE



If we chose block 21, then the first constraint becomes

$$z_{16} \leq \sum_{i \in \mathcal{W}_{16}} y_i \Rightarrow z_{16} \leq y_{12} + y_{21} \Rightarrow 1 \leq 0 + 1$$

## EXAMPLE

1	2	3	4	5
6	7	8	9	10
11	12	13	14	15
16	17	18	19	20
21	22	23	24	25

If we chose block 21, then the second constraint becomes

$$|\mathcal{W}_4| z_4 \geq \sum_{i \in \mathcal{W}_4} y_i \Rightarrow (1 \times 0) \geq y_3 \Rightarrow 0 \geq 0$$

# COMPLICATIONS

---

- Maximize number of beneficiaries?

- Maximize number of beneficiaries?
- Maximize accessibility?



- Maximize number of beneficiaries?
- Maximize accessibility?
- Maximize connectivity to existing facilities?

- Maximize number of beneficiaries?
- Maximize accessibility?
- Maximize connectivity to existing facilities?
- Place parks closer to positive facilities (like schools) and further from negative facilities (like waste treatment plants)?

- Maximize number of beneficiaries?
- Maximize accessibility?
- Maximize connectivity to existing facilities?
- Place parks closer to positive facilities (like schools) and further from negative facilities (like waste treatment plants)?
- Minimize cost?

What if we want to do all these things?

What if we want to do all these things?

Need a way to maximize or minimize multiple objective functions at a time.

- Now what if we had 100 blocks? 1000 blocks? More?

- Now what if we had 100 blocks? 1000 blocks? More?
- What if the parcels weren't all the same size?

- Now what if we had 100 blocks? 1000 blocks? More?
- What if the parcels weren't all the same size?
- What if the parcels weren't all the same shape?



- Now what if we had 100 blocks? 1000 blocks? More?
- What if the parcels weren't all the same size?
- What if the parcels weren't all the same shape?
- How do you measure these things?

# CONCLUSION

---

Locating parks is a complicated problem.

QUESTIONS?

