

A novel and selective oral PI3K α/δ inhibitor, TQ-B3525, in patients with relapsed and/or refractory follicular lymphoma: A phase II, single-arm, open-label study.

Huaqing Wang, Aibin Liang, Jifeng Feng, YANYAN LIU, Zhengzi Qian, Da Gao, Xuehong Ran, Hui Zhou, Lihong Liu, Binghua Wang, Meiyun Fang, Hebing Zhou, Zhenqian Huang, Shi Tao, Zhuowen Chen, Liping Su, Hang Su, Yu Yang, Xiaobao Xie, Zhiming Li; Department of Medical Oncology, The Institute of Translational Medicine, Tianjin Union Medical Center of Nankai University, Tianjin, China; Department of Hematology, Tongji Hospital of Tongji University, Shanghai, China; Department of Hematology, Jiangsu Cancer Hospital, The Affiliated Cancer Hospital of Nanjing Medical University, Nanjing, China; Department of Medical Oncology, Henan Cancer Hospital, The Affiliated Cancer Hospital of Zhengzhou University, Hengzhou, China; Department of Medical Oncology, Tianjin Medical University Cancer Institute and Hospital, Tianjin, China; Department of Hematology, The Affiliated Hospital of Inner Mongolia Medical College, Hohhot, China; Department of Hematology, Weifang People's Hospital, The First Affiliated Hospital of Weifang Medical University, Weifang, China; Department of Lymphoma & Hematology, Hunan Cancer Hospital, Changsha, China; The Fourth Hospital of Hebei Medical University, Shijiazhuang, China; Department of Lymphoma, Weihai Central Hospital, Weihai, China; Department of Hematology and Rheumatology, The Affiliated Zhongshan Hospital of Dalian University, Dalian, China; Department of Hematology, Beijing Luhe Hospital, Beijing, China; Department of Hematology, The First Affiliated Hospital of Guangzhou Medical University, Guangzhou, China; Department of Hematology, The First Affiliated Hospital of Hainan Medical College, Haikou, China; Department of Hematology, The First People's Hospital of Foshan, Foshan, China; Department of Hematology, Shanxi Cancer Hospital, Taiyuan, China; Department of Lymphoma, Senior Department of Hematology, The Fifth Medical Center of Chinese PLA General Hospital, Beijing, China; Department of Lymphoma and Head and Neck Cancer, Fujian Cancer Hospital, Fuzhou, China; Department of Hematology, The First People's Hospital of Changzhou, The Third Affiliated Hospital of Soochow University, Changzhou, China; Department of Medical Oncology, State Key Laboratory of Oncology in South China, Collaborative Innovation Center for Cancer Medicine, Guangdong Provincial Clinical Research Center for Cancer, Sun Yat-sen University Cancer Center, Guangzhou, China

Background: TQ-B3525 is a novel and selective oral PI3K α/δ inhibitor. In an earlier Phase I trial, TQ-B3525 achieved outstanding efficacy in subjects with refractory/relapsed follicular lymphoma (R/R FL) (2020 ASCO Abstract #8058). Here, we report the result from a single-arm, open-label, phase II registration study evaluating the safety and efficacy of TQ-B3525 in R/R FL patients. **Methods:** This phase II study included exploratory stage 1 and confirmatory stage 2. Patients with R/R FL after ≥ 2 lines therapies received oral 20 mg TQ-B3525 once daily in a 28-day cycle until disease progression or intolerable toxicity. Primary endpoint was independent review committee (IRC)-assessed objective response rate (ORR). Secondary endpoints were ORR by investigator assessment; the IRC- and investigator-assessed disease control rate (DCR), time to response (TTR), duration of response (DOR), progression-free survival (PFS), overall survival (OS), and safety. **Results:** Based on results (ORR, 88.0%; DOR, 11.8 months; PFS, 12.0 months) in 25 patients at stage 1, second stage study was initiated and included 82 patients for efficacy/safety analysis. Patients received a median of 3 prior lines, with 56.1% refractory to previous therapies; 73.2% experienced POD24 at baseline. At stage 2, ORR was 86.6% (71/82; 95% CI, 77.3%-93.1%), with 28 (34.2%) complete responses. Seven (8.5%) had stable disease for DCR of 95.1%. Median TTR was 1.8 months. Among 71 responders, median DOR was not reached; 18-month DOR rate was 51.6%. At median follow-up of 13.3 months, median PFS was 18.5 (95% CI, 10.2-not estimable) months; estimated 24-month OS rate was 86.1%. Response rates and survival data were consistent across all subgroups. Grade 3 or higher treatment-related adverse events occurred in 63 (76.8%) patients, with neutrophil count decreased (22.0%), hyperglycemia (19.5%), and diarrhea (13.4%) being common. **Conclusions:** TQ-B3525 exhibited favorable efficacy and manageable safety profiles, supporting its potential as a valuable treatment modality for heavily pretreated Chinese R/R/ FL patients. Clinical trial information: NCT04324879. Research Sponsor: Chia Tai Tianqing Pharmaceutical Group Co., Ltd. (Nanjing, China); National Natural Science Foundation of China; 81872902, 82073917, and 82070206; National Natural Science Foundation of Guangdong Province; 2023A1515011525; the Lymphoma Research Fund of China Anti-Cancer Association; the Sun Yat-sen University Cancer Center Clinical Research 308 Program; 2014-fxy-106 and 2016-fxy-079; Tianjin Key Medical Discipline (Specialty) Construction Project; TJYXZDXK-053B.

Summary of TQ-B3525 efficacy in the ITT at stage 2.

Efficacy	IRC-Assessed (n=82)	Investigator-Assessed (n=82)
Median DOR [#] , months (95% CI)	NR (9.2-NE)	14.8 (9.2-20.4)
12-month DOR rate (%; 95% CI)	60.2% (44.3%-72.9%)	51.5% (36.0%-65.0%)
18-month DOR rate (%; 95% CI)	51.6% (30.6%-69.2%)	41.0% (23.5%-57.7%)
Median PFS, months (95% CI)	18.5 (10.2-NE)	18.4 (11.0-22.0)
12-month PFS rate (%; 95% CI)	58.3% (44.0%-70.1%)	52.8% (38.8%-65.0%)
18-month PFS rate (%; 95% CI)	58.3% (44.0%-70.1%)	52.8% (38.8%-65.0%)
Median OS, months (95% CI)	Not reached	
12-month OS rate (%; 95% CI)	91.8% (82.5%-96.3%)	
24-month OS rate (%; 95% CI)	86.1% (72.3%-93.3%)	

Data were presented as n (%) or n (%; 95% confidence interval). NE, not estimable # DOR was assessed in 71 patients achieving responses.