

**Assignment 2**  
**Operation Research**  
**BTech Sem IV (20-21 Batch)**

Q1. Describe graphically what the simplex method does step by step to solve the following problem. (4.1.6)

$$\begin{aligned} &\text{Maximize } Z = 2x_1 + 3x_2, \\ &\text{subject to} \\ &\quad -3x_1 + x_2 \leq 1 \\ &\quad 4x_1 + 2x_2 \leq 20 \\ &\quad 4x_1 + x_2 \leq 10 \\ &\quad -x_1 + 2x_2 \leq 5 \\ &\text{and} \\ &\quad x_1 \geq 0, x_2 \geq 0. \end{aligned}$$

Q2. Describe graphically what the simplex method does step by step to solve the following problem. (4.1.7)

$$\begin{aligned} &\text{Minimize } Z = 5x_1 + 7x_2, \\ &\text{subject to} \\ &\quad 2x_1 + 3x_2 \geq 147 \\ &\quad 3x_1 + 4x_2 \geq 210 \\ &\quad x_1 - x_2 \geq 63 \\ &\text{and} \\ &\quad x_1 \geq 0, x_2 \geq 0. \end{aligned}$$

Q3. Consider the following problem. (4.2.2)

$$\begin{aligned} &\text{Maximize } Z = x_1 + 2x_2, \\ &\text{subject to} \\ &\quad x_1 + 3x_2 \leq 8 \\ &\quad x_1 + x_2 \leq 4 \\ &\text{and} \\ &\quad x_1 \geq 0, x_2 \geq 0. \end{aligned}$$

- (a)** Introduce slack variables in order to write the functional constraints in augmented form.
- (b)** For each CPF solution, identify the corresponding BF solution by calculating the values of the slack variables. For each BF solution, use the values of the variables to identify the nonbasic variables and the basic variables.
- (c)** For each BF solution, demonstrate (by plugging in the solution) that, after the nonbasic variables are set equal to zero, this BF solution also is the simultaneous solution of the system of equations obtained in part (a).
- (d)** Repeat part (b) for the corner-point infeasible solutions and the corresponding basic infeasible solutions.
- (e)** Repeat part (c) for the basic infeasible solutions.

Q4. Consider the problem in Q3. . (4.3.3)

- a) Work through the simplex method (in algebraic form) step by step to solve the model
- b) Verify the optimal solution you obtained by using a software package based on the simplex method.

## 4.3-3.

$$\begin{aligned}
 \text{(a) maximize} \quad & Z = x_1 + 2x_2 \\
 \text{subject to} \quad & x_1 + 3x_2 + x_3 = 8 \\
 & x_1 + x_2 + x_4 = 4 \\
 & x_1, x_2, x_3, x_4 \geq 0
 \end{aligned}$$

Initialization:  $x_1 = x_2 = 0 \Rightarrow x_3 = 8, x_4 = 4, z = x_1 + 2x_2 = 0$ , is not optimal since the improvement rates are positive. Since it offers a rate of improvement of 2, choose to increase  $x_2$ , which becomes the entering basic variable for Iteration 1. Given  $x_1 = 0$ , the highest possible increase in  $x_2$  is found by looking at:

$$x_3 = 8 - 3x_2 \geq 0 \Rightarrow x_2 \leq 8/3$$

$$x_4 = 4 - x_2 \geq 0 \Rightarrow x_2 \leq 4$$

The minimum of these two bounds is  $8/3$ , so  $x_2$  can be raised to  $8/3$  and  $x_3 = 0$  leaves the basis. Using Gaussian elimination, we obtain:

$$\begin{aligned}
 Z &= \frac{1}{3}x_1 - \frac{2}{3}x_3 + \frac{16}{3} \\
 \frac{1}{3}x_1 + x_2 + \frac{1}{3}x_3 &= \frac{8}{3} \\
 \frac{2}{3}x_1 - \frac{1}{3}x_3 + x_4 &= \frac{4}{3} \\
 x_1, x_2, x_3, x_4 &\geq 0
 \end{aligned}$$

Again  $(0, \frac{8}{3}, 0, \frac{4}{3})$  is not optimal since the rate of improvement for  $x_1$  is  $\frac{1}{3} > 0$  and  $x_1$  can be increased to 2. Consequently,  $x_4$  becomes 0. By Gaussian elimination:

$$\begin{aligned}
 Z &= -\frac{1}{2}x_3 - \frac{1}{2}x_4 + 6 \\
 x_2 + \frac{1}{2}x_3 + \frac{1}{2}x_4 &= 2 \\
 x_1 - \frac{1}{2}x_3 + \frac{3}{2}x_4 &= 2 \\
 x_1, x_2, x_3, x_4 &\geq 0
 \end{aligned}$$

The current solution is optimal, since increasing  $x_3$  or  $x_4$  would decrease the objective value. Hence  $x^* = (2, 2, 0, 0)$ ,  $Z^* = 6$ .

[illegible]

5. Work through the simplex method step by step (in tabular form) to solve the following problem. (4.4-7)

$$\text{Maximize } Z = 2x_1 - x_2 + x_3,$$

subject to,

$$3x_1 + x_2 + x_3 \leq 6$$

$$x_1 - x_2 + 2x_3 \leq 1$$

$$x_1 + x_2 - x_3 \leq 2$$

and

$$x_1 \geq 0, x_3 \geq 0, x_3 \geq 0.$$

Iteration n	Basic Var.		Equation	Z	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$	RHS	Minimum Ratio Test	
1	Z		(0)	1	-2	1	-1	0	0	0	0		
Solution	$x_4$		(1)	0	3	1	1	1	0	0	6	2	
(0 0 0 2 1 2)	$x_5$		(2)	0	1	-1	2	0	1	0	1	1	Min
Z = 0	$x_6$		(3)	0	1	1	-1	0	0	1	2	2	
	EV	$x_1$								LV	$x_5$		
2			(0)	1	0	-1	3	0	2	0	2		
Solution	$x_4$		(1)	0	0	4	-5	1	-3	0	3	3/4	
(1 0 0 3 0 1)	$x_1$		(2)	0	1	-1	2	0	1	0	1	-1	New Equation
	$x_6$		(3)	0	0	2	-3	0	-1	1	1	1/2	Min
	EV	$x_2$								LV	$x_6$		
3			(0)	1	0	0	1.5	0	1.5	0.5	2.5		
Solution	$x_4$		(1)	0	0	0	1	1	-1	-2	1		
(1.5 0.5 0 1 0 0)	$x_1$		(2)	0	1	0	0.5	0	0.5	0.5	1.5		
Z = 2.5	$x_2$		(3)	0	0	1	-1.5	0	-0.5	0.5	0.5		New Equation

6. Consider the following problem. (4.4-6.)

Maximize  $Z = 3x_1 + 5x_2 + 6x_3,$

subject to

$$2x_1 + x_2 + x_3 \leq 4$$

$$x_1 + 2x_2 + x_3 \leq 4$$

$$x_1 + x_2 + 2x_3 \leq 4$$

$$x_1 + x_2 + x_3 \leq 3$$

and

$$x_1 \geq 0, x_2 \geq 0, x_3 \geq 0$$

- Work through the simplex method step by step in algebraic form.
- Work through the simplex method in tabular form.
- Use a computer package based on the simplex method to solve the problem.

[illegible]

7. Consider the following statements about linear programming and the simplex method. Label each statement as true or false, and then justify your answer. (4.5-1)

(a) In a particular iteration of the simplex method, if there is a tie for which variable should be the leaving basic variable, then the next BF solution must have at least one basic variable equal to zero.

(b) If there is no leaving basic variable at some iteration, then the problem has no feasible solutions.

(c) If at least one of the basic variables has a coefficient of zero in row 0 of the final tableau, then the problem has multiple optimal solutions.

(d) If the problem has multiple optimal solutions, then the problem must have a bounded feasible region.

8. Consider the following problem. 4.6-17

Maximize  $Z = 4x_1 + 5x_2 + 3x_3$ ,  
subject to

$$\begin{aligned}x_1 + x_2 + 2x_3 &\geq 20 \\15x_1 + 6x_2 - 5x_3 &\leq 50 \\x_1 + 3x_2 + 5x_3 &\leq 30\end{aligned}$$

and

$$x_1 \geq 0, x_2 \geq 0, x_3 \geq 0$$

Work through the simplex method step by step to demonstrate that this problem does not possess any feasible solutions.

9. Label each of the following statements as true or false, and then justify your answer. 4.6-11.

(a) When a linear programming model has an equality constraint, an artificial variable is introduced into this constraint in order to start the simplex method with an obvious initial basic solution that is feasible for the original model.

(b) When an artificial problem is created by introducing artificial variables and using the Big  $M$  method, if all artificial variables in an optimal solution for the artificial problem are equal to zero, then the real problem has no feasible solutions.

(c) The two-phase method is commonly used in practice because it usually requires fewer iterations to reach an optimal solution than the Big  $M$  method does.

10. Consider the following problem. 4.6-5.

Maximize  $Z = 5x_1 + 4x_2$ ,  
subject to

$$3x_1 + 2x_2 \leq 6$$

$$2x_1 - x_2 \geq 6$$

and

$$x_1 \geq 0, x_2 \geq 0.$$

(a) Demonstrate graphically that this problem has no feasible solutions.

(b) Use a computer package based on the simplex method to determine that the problem has no feasible solutions.

(c) Using the Big  $M$  method, work through the simplex method step by step to demonstrate that the problem has no feasible solutions.

11. Consider the following problem. 4.6-9.

Minimize  $Z = 3x_1 + 2x_2 + 4x_3$

subject to

$$2x_1 + x_2 + 3x_3 = 60$$

$$3x_1 + 3x_2 + 5x_3 \geq 120$$

and

$$x_1 \geq 0, x_2 \geq 0, x_3 \geq 0$$

(a) Using the Big  $M$  method, work through the simplex method step by step to solve the problem.

(b) Use a software package based on the simplex method to solve the problem.

12. Consider the following problem 4.6-10.

Minimize  $Z = 3x_1 + 2x_2 + 7x_3$ ,

subject to

$$-x_1 + x_2 = 10$$

$$2x_1 + x_2 + x_3 \geq 10$$

and

$$x_1 \geq 0, x_2 \geq 0, x_3 \geq 0$$

(a) Using the Big  $M$  method, work through the simplex method step by step to solve the problem

(b) Use a software package based on the simplex method to solve the problem.



13. Consider the following problem. 4.7-4.

Maximize  $Z = x_1 - 7x_2 + 3x_3$ ,  
subject to

$$2x_1 + x_2 - x_3 \leq 4 \text{ (resource 1)}$$

$$4x_1 - 3x_2 \leq 2 \text{ (resource 2)}$$

$$-3x_1 + 2x_2 + x_3 \leq 3 \text{ (resource 3)}$$

and

$$x_1 \geq 0, x_2 \geq 0, x_3 \geq 0.$$

(a) Work through the simplex method step by step to solve the problem.

(b) Identify the shadow prices for the three resources and describe their significance.

(c) Use a software package based on the simplex method to solve the problem and then to generate sensitivity information.

Use this information to identify the shadow price for each resource, the allowable range for each objective function coefficient, and the allowable range for each right hand side.

14. Consider the following problem. 4.7-6.

Maximize  $Z = 5x_1 + 4x_2 - x_3 + 3x_4$ ,  
subject to

$$3x_1 + 2x_2 - 3x_3 + x_4 \leq 24 \text{ (resource 1)}$$

$$3x_1 + 3x_2 + x_3 + 3x_4 \leq 36 \text{ (resource 2)}$$

and

$$x_1 \geq 0, x_2 \geq 0, x_3 \geq 0, x_4 \geq 0$$

(a) Work through the simplex method step by step to solve the problem.

(b) Identify the shadow prices for the two resources and describe their significance.

(c) Use a software package based on the simplex method to solve the problem and then to generate sensitivity information. Use this information to identify the shadow price for each resource, the allowable range for each objective function coefficient, and the allowable range for each right-hand side.