(8.1) 20 Question. with 4 marks each. Ø20 A200 Ø Too JAN ON Q38 p = peus of that A wins the game n=5, p=3 (+1,61,21, H1, SID) 028 Q19,20,21,22,23 035,36,37 240 03 (030 wrongly, he gets 48-8-40 marks. If he gets more that 12 correct answers, he gets more than 40 marks. 35 nx 10+4 x14 bood feggs. A= 4 m=6, b=10=0.1 at students solved 12 Quest consectly R & Ques Then And PCG) P(2 (72) = P(12) + -- -+ P(20) P = perobability of getting a consect Ans = } m -18, p-1 P= P(\$ or 5) - 1+ 2 -2 -1, m-6
P= P(\$ or 5) - 1+ 2 -2 -2, m-6
P= P(\$ or 5) - 1+ 2 -2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P(\$ or 5) - 1+ 2 -3, m-6
P= P( 7 11 3 P(5 even)= 2P(5 add) = 2P(15 even) P = getting a 9 = 4 = 1, \$ (2) X(n, b) Y(m, b) Then X+Y (m+n, b) N=3, N=3000 P=1/2-- 11 1.0 - 52-9 (-032