

(a) PP-DocLayoutV3 detection

have n dis-
at least 2,
of general-
 $P(x)$, so
omputation

$R''(x)$.

t the great-
but $P(x) =$
ce (since O

and the other of which sums to $xy / [(1 - y)(1 - y^2)]$
Therefore

Formula conf 0.93

$$S(x,y) = \frac{x^3y}{(1-x)(1-y)} - \frac{xy^3}{(1-x)(1-x^2y)} - \frac{xy^3}{(1-y)(1-xy^2)}$$
$$= \frac{xy(1+x+y+xy-x^2y^2)}{(1-x^2y)(1-xy^2)}$$

and the desired limit is

(c) band chunks: fraction fusion + trimming

B0

B1

$$S(x,y) = \frac{xy}{(1-x)(1-y)} - \frac{x^3y}{(1-x)(1-x^2y)}$$

isolated chunks in B0
trimmed (speck + detached numerator)

B2


$$= \frac{xy^3}{(1-y)(1-xy^2)}$$

B3

$$= \frac{xy(1+x+y+xy-x^2y^2)}{(1-x^2y)(1-xy^2)}$$

num / bar / den
re-fused into B3

(b) Otsu ink projection, row-band splitting



(d) Texo decode, compiled

with geometry layer — per-line CDM 0.90 / 0.77 / 0.97
 xy

$$S(x,y) = \frac{x^3y}{(1-x)(1-y)} - \frac{xy^3}{(1-x)(1-x^2y)} - \frac{xy^3}{(1-y)(1-xy^2)}$$
$$= \frac{xy(1+x+y+xy-x^2y^2)}{(1-x^2y)(1-xy^2)}$$

naive whole-box decode (geometry layer off)

$$\langle 1 - \chi \downarrow 1 - \chi V^- \rangle$$
$$x\mathcal{V}(1+x+v+xv-x^-v^-)$$
$$1-x^2v \ln 1-xv^2$$