


INTRODUCTION TO MACHINE LEARNING COMPSCI 4ML3

LECTURE 3

HASSAN ASHTIANI

ORDINARY LEAST SQUARES (D-DIMENSIONS)

- ASSUME $x \in \mathbb{R}^d$, $y \in \mathbb{R}$
- INSTEAD OF A LINE, WE NEED TO FIT A HYPERPLANE!
- WHY ARE THE LINES VERTICAL?
 - ANY DIFFERENT IF WE MINIMIZE THE DISTANCE TO THE HYPERPLANE?



MATRIX FORM OLS

$$\bullet \Delta = \begin{pmatrix} \Delta_1 \\ \dots \\ \Delta_n \end{pmatrix} = \begin{pmatrix} x_1^1 & \dots & x_d^1 \\ \vdots & \ddots & \vdots \\ x_1^n & \dots & x_d^n \end{pmatrix} \begin{pmatrix} w_1 \\ \dots \\ w_d \end{pmatrix} - \begin{pmatrix} y^1 \\ \dots \\ y^n \end{pmatrix}$$

$$\min_{W \in \mathbb{R}^{d \times 1}} \sum_{i=1}^n (\Delta_i)^2 = \min_{W \in \mathbb{R}^{d \times 1}} \langle \Delta, \Delta \rangle = \min_{W \in \mathbb{R}^{d \times 1}} \|\Delta\|_2^2$$

$$\min_{W \in \mathbb{R}^{d \times 1}} \|XW - Y\|_2^2$$

TAKING THE “DERIVATIVE”

- REAL-VALUED FUNCTION OF A VECTOR

GRADIENT:

- VECTOR-VALUED FUNCTION OF A VECTOR

JACOBIAN:

MATRIX/VECTOR CALCULUS

- $u, v \in R^n$
- $g(u) = u^T v$
- $\nabla u(g) =$

MATRIX/VECTOR CALCULUS

- $A \in R^{m \times n}, u \in R^n$
- $g(u) = Au$
- $\nabla u(g) =$

MATRIX/VECTOR CALCULUS

- $A \in R^{m \times n}, u \in R^n$
- $g(u) = u^T A u$
- $\nabla u(g) =$

SOLVING OLS

$$f(W) = \min_{W \in \mathbb{R}^{d \times 1}} \|XW - Y\|_2^2. \quad \text{WHAT IS } \nabla f?$$



SOLVING OLS

$$W^{LS} = (X^T X)^{-1} X^T Y$$

- DEGENERATE CASE WHEN $X^T X$ IS NOT INVERTIBLE?

BIAS/INTERCEPT TERM

- WE ARE MISSING THE BIAS TERM (w_0)

$$\min_{w_0, w_1, \dots, w_d \in \mathbb{R}} \sum_{i=1}^n (w_1 x_1^i + \dots + w_d x_d^i + w_0 - y^i)^2$$

- MATRIX FORM WITH THE BIAS TERM?

$$\min_{W \in \mathbb{R}^{d \times 1}, w_0 \in \mathbb{R}} \left\| XW + \begin{pmatrix} w_0 \\ w_0 \\ \dots \\ w_0 \end{pmatrix} - Y \right\|_2^2$$

EXAMPLE

BIAS/INTERCEPT TERM

- ADD A NEW AUXILIARY DIMENSION TO THE DATA

- $X'_{n \times (d+1)} = \begin{pmatrix} x_1^1 & \dots & x_d^1 & 1 \\ \vdots & \ddots & \vdots & 1 \\ x_1^n & \dots & x_d^n & 1 \end{pmatrix}, W'_{(d+1) \times 1} = \begin{pmatrix} w_1 \\ \vdots \\ w_d \\ w_0 \end{pmatrix}$

- SOLVE OLS: $\min_{W' \in \mathbb{R}^{(d+1) \times 1}} \|X'W' - Y\|_2^2$

- w_0 WILL BE THE BIAS TERM!

SOME EXAMPLES

- OLS NOTEBOOK