Spatial modeling of tissues for morphogenic field analysis

Aaron Osgood-Zimmerman[1☯*], Micha Sam Brickman Raredon[2,3,4☯*]

1 Department of Mathematics, Bucknell University, Lewisburg, PA, USA
2 Department of Anesthesiology, Yale School of Medicine, New Haven, CT, USA
3 Vascular Biology and Therapeutics, Yale University, New Haven, CT, USA
4 Program in Translational Biomedicine, Yale Biological and Biomedical Sciences, New Haven, CT, USA

☯These authors contributed equally to this work.
* aaron.oz@bucknell.edu (AOZ); michasam.raredon@yale.edu (MSBR)

Abstract

Tissues are shaped by extracellular signaling fields which convey information between cells. ~~The cellular composition of tissues, and the extracellular signaling within the tissue, are innately spatially structured.~~ Modern spatialomics data provide ~~unprecedented~~ detailed measurement of ligand and receptor expressivity *in situ* from tissue sections. Here ~~, we show that by adapting generalizable~~ we adapt geospatial statistical models to spatialomics data, ~~we are able to reveal statistically-detailed portraits of morphogenic field interactions within tissues and thereby approach a richer set of biologic questions than is typically pursued. The general methods piloted here can readily be applied to spatialomics data from diverse platforms with no need to alter data collection techniques. Our results demonstrate that the application of spatial statistical modeling to spatialomics data opens many avenues for future experimentation that will be valuable to fundamental biology and to regenerative medicine~~fitting a continuous spatial field with quantified uncertainty to the raw counts of each of 912 ligand and receptor genes across a single field of view from a mouse embryo section, and convolving these into inferred interaction fields for 1,477 ligand-receptor pairs. Clustering this high-dimensional set of fields without dimensionality reduction yields spatially coherent tissue domains. Refitting the entire pipeline on spatially permuted data collapses the domains, which argues that they are not manufactured by the spatial smoother or by the high dimensionality of the clustering. The domains also agree, well above chance, with the anatomical annotation published independently for the same section (adjusted Rand index 0.401), while subdividing several organs into multiple territories. Randomly re-pairing ligands with receptors does not make the domains less coherent, so they reflect the collection of fitted ligand and receptor fields rather than the specific pairings, even though true partners are measurably more alike in spatial pattern than random ones. Computing the posterior of each interaction field from the fitted marginal posteriors shows that the plug-in fields used here misstate its level, most where neither partner is clearly limiting, while largely preserving its spatial pattern. The inferred fields estimate the potential for signaling under the co-expression assumption standard in cell-cell communication analysis; derived from transcript counts, they are not a measurement of ligand-receptor binding or of signaling. Our demonstration uses one field of view, from one section, on one platform, at one binning resolution; rather than asserting general applicability, we set out the conditions we expect transfer to other platforms would require.

Introduction




Fig 1. Project schematic. The purple arrows represent true tissue processing and activity. The blurred regions represent coarse estimates captured by spatialomics or created by spatial modeling.

Morphogenic field interactions are fundamental to tissue biology and regenerative medicine

Cells within tissues are continuously processing information and updating their states, receiving cues from their local microenvironment and contributing cues to their local milieu. The study of cellular connectivity (the potential for information flow between cells) and cell-to-cell signaling (information flow between cells) is fundamental to the study of tissue biology. Accurate modeling of cellular connectivity and cell-to-cell signaling, particularly through ligand-receptor syntax, provides a quantitative foundation for understanding tissue-scale changes and dynamics (Fig 1).

It has long been understood that tissues are shaped by morphogenic signaling between cells [1–4]. During development, homeostasis, disease, and regenerative remodeling, cells within tissues release signaling molecules which diffuse through the extracellular matrix and can be locally received and processed by resident cells. Until recently, it has only been possible to capture data on a small number of morphogenic mechanisms at once from a single tissue. However, it is the *holistic set* of soluble and insoluble cue intensities – taken together as a single combinatoric 'input' [5–7]– that encodes the full spectrum of locally-transducable signaling information, often irreducible to the individual cues themselves [8–10]. In other words, extracellular signaling is intrinsically high-dimensional —(information is encoded via specific combinations of cues at particular levels, which are processed in parallel by resident cells—) and is

intrinsically spatially patterned: as an organism develops, intersecting 3-dimensional signaling gradients combine to drive tightly localized cellular differentiation, proliferation, and migration events. These morphogenic field interactions are major governors of tissue-scale emergent behavior including developmental morphogenesis and adult tissue inflammation & homeostasis [11–14]. To understand tissue-scale change, therefore, we need robust tools to study high-dimensional signaling and connectivity dynamics in both time and space.

Spatialomics is newly presenting scientists with an opportunity to estimate proxies for thousands of morphogenic fields at once, at both the raw and relative information level, thereby allowing us to model and visualize ~~the specific complex patterns of extracellular signaling that controls~~ patterns that are consistent with the extracellular signaling thought to control cellular differentiation and transition in living tissues. However, to properly handle and analyze these data, we need appropriately designed statistical approaches. This paper presents one such approach, and shows that by adapting techniques common in geospatial data analysis, we are able to ~~newly~~ model and visualize, with quantified uncertainty, the high-dimensional patterns of extracellular signaling that have long been theorized to be fundamental to tissue growth and change.

The critical need for spatial statistics in tissue biology

Spatialomics data differ from the true state of the tissue in two important ways. First, regardless of platform resolution, all spatialomics data collection methods collect data at discrete locations, necessarily creating error in reported spatial sampling coordinates. Second, no bulk RNA, single cell, or spatialomics methods perform true censuses – that is, they do not measure every single molecule within every sample, and in many cases measure only a small fraction of the total present molecules [15, 16]. As a result, observed data from all spatial transcriptomic methods is subject to sampling variability. Ignoring sampling variability while performing data-analysis can lead to bias, over- or under-confidence in results, and false findings [17]. Properly accounting for the sampling variability in any dataset requires a statistical model aligned with the data-generating mechanism.

Our statistical goal in this work was to take spatialomics data observations as input and to produce output which estimates the true state of the tissue, acknowledging and accounting for the important ways in which the data differs from the tissue itself. That is, we need statistical models to take transcriptomic data measured at **discrete** spatial locations as input and to produce estimates of **continuous** spatial fields as outputs, along with estimates of uncertainty, which act as a statistical prediction of transcriptomic counts across the spatial observation domain. Spatial statistics has a rich literature of tools to leverage which are designed exactly for this purpose. Furthermore, they account for the spatial autocorrelation present within these datasets, which, if ignored, can again lead to incorrect uncertainty estimates and false findings [18].

Biological tissues are spatially structured

Advances in genetics data collection methodologies have led to rich, beautiful datasets which can easily be mistaken to be "the truth." Spatialomics, loosely defined as the group of technologies capable of measuring molecular characteristics of tissues within their natural spatial alignment, has justifiably been heralded as a transformative frontier in life sciences. However, the quickly developing methods, and the immense amounts of data produced, have outpaced *analysis techniques*, which need to mature [19].

Spatialomics collection methods have been developed precisely because there is observable and necessary spatial structure apparent in tissues. Yet, this fact isn't often explicitly incorporated in analyses. The spatial structure of healthy vs. diseased tissues

is often radically dissimilar, but many currently-practiced computational methods are descendants of single cell techniques which were necessarily non-spatial. Ignoring the spatial organization, at best, may ~~be omitting~~ omit useful information and, at worst, may ~~be~~ lead to an incorrect interpretation of the data. When approaching morphogenic field analysis, i.e., the quantitative study of high-dimensional extracellular signaling mediated by ligand-receptor binding interactions via the extracellular milieu, it is essential to take these principles into account, or we risk misinterpreting statistically spurious findings.

In this manuscript we introduce a flexible continuous spatial modeling framework that can be used to estimate one or more features measured via spatialomics. The proposed approach also provides uncertainty estimates of each quantity over space~~, which allows propagation of uncertainty~~. Because those estimates are posterior distributions rather than point summaries, they can in principle be propagated to downstream outputs ~~(~~such as ligand-receptor binding density~~, the main biologic focus of this paper.)~~; we note where the analyses reported here use posterior summaries instead, and treat propagation as an available extension rather than something this manuscript demonstrates. The modeling in this manuscript is meant to be a launchpad for the use of powerful, existing frameworks to learn more readily and deeply from spatialomics data, thereby decreasing the chance of generating unreproducible results. We use our bespoke pipeline to quantitatively assess a portion of a public dataset capturing mammalian embryonic organogenesis. ~~We demonstrate via our results that~~ Our results show that, in the tissue section studied here, complex domain architecture ~~in tissues~~ can be derived ~~directly from~~ from inferred spatial ligand-receptor ~~connectivity patterns , even~~ patterns alone, without study of transcripts coding for intracellular proteins.

Materials and methods

Overview

Principles of ligand-receptor analysis

Our focus in this paper was to model a useful proxy for local ligand-receptor binding *density*. This is different from local ligand-receptor connectivity *character*, which is the primary focus of many cell-to-cell communication inference tools including our prior work [20]. This distinction is non-trivial. When ligand-receptor connectivity *character* is computed, individual feature values represent a proxy for *relative* signal strength; when ligand-receptor binding *density* is computed, individual feature values represent a proxy for *absolute* signal strength. Our focus on binding density in this paper, rather than character, ~~allows us to propagate our outputs~~ is what makes the outputs suitable for propagation to downstream models~~more easily and with fewer artifacts, as is shown in the Results section~~, since an absolute quantity can be carried forward without the reference-dependence that a relative one introduces. By applying spatial statistics to the raw counts in spatial transcriptomics, rather than the normalized counts, our model outputs ~~are easily~~ can be treated as a ~~new fundamental~~ distinct layer of information for multi-modal data analysis, one that ~~directly estimates~~ estimates a proxy for local ligand-receptor binding density within the tissue. This output can then be normalized, similarly to how we usually normalize local gene counts to relative values, to yield proxies for local ligand-receptor connectivity *character* more familiar to most in the field.

All connectivity tools require us to convolve information from the ligand field with information from the receptor field. Historically, this has been done by computing either the product [20] or the geometric mean [21] of all subunits involved. Here, we ~~have~~

~~experimentally pioneered the use of~~ tested alternative approaches to convolving the ligand and the receptor values. Assuming that the counts of individual transcripts are proportional to the counts of individual proteins (an almost certainly *incorrect* assumption, but one that we are forced to accept at this time) we may treat ligand-receptor binding as a stoichiometric process, i.e., a function of the concentration of the ligand and the concentration of the receptor.

Ligands diffuse locally, with a characteristic diffusion distance roughly proportional to the molecular weight and chemical character of the molecular species and how it interacts with other matrix elements. Extracellular ligands can be sensed and processed by tissue-resident cells expressing specific receptor profiles. ~~All previous computational approaches~~ The computational approaches we are aware of address this fundamental property of tissue biology by first creating local neighborhoods in histologic space and then computing cell-to-cell ligand-receptor connectivity within each neighborhood. When working with continuous field models, as is demonstrated here, local neighborhood construction is no longer necessary, because the spatial field, intrinsically, can be considered as a filter which takes in noisy spatial data and returns a smoothed version. Although the smoothed output is technically an estimate of the underlying state of the tissue, in practice the smoothing process can be viewed as sharing locally-measured information between nearby areas. This modeling approach thereby ~~builds a robust~~ provides a framework for convolution of ligand and receptor fields. The fields themselves can be independently tuned to reflect specific molecular properties, and the convolution approach customized to reflect binding kinetics.

Principles of spatial modeling

Spatial statistics is a field of statistics used to study datasets in which the location of the data observations is recorded. Typically, a spatial statistical model will explicitly account for spatial dependence among observations to acknowledge that spatially nearby observations are often more strongly correlated than observations that are located further apart. The purpose of most statistical models is to use imperfect data to provide an estimate of a population-level target of interest and the associated uncertainty of that estimate (from which statistical inference, e.g. confidence intervals or hypothesis tests can be calculated and performed). Leveraging spatial correlation when it exists can result in more precise estimates and ignoring it when it exists leads to incorrect (typically overly confident) uncertainty estimates [22]. Non-probabilistic interpolation techniques cannot account for the sampling variability or produce reliable uncertainty estimates. In this work, we demonstrate a spatial statistical model that can be used to estimate biological fields across the tissue from which the spatialomics data sample was collected. The estimated fields represent the result that would be expected if the experiment were repeated many times.

One of the primary decisions to make when performing a spatial statistical analysis is whether the spatial domain (and thus the model) should be discrete or continuous in nature. Despite that many of the contemporary spatialomics data collection tools generate data on discrete "spot" arrays, the underlying tissue- and cellular-level *processes*~~—~~, which are the target of our study~~—~~, are not constrained to occur only at data observation locations (spots) or any finite set of locations.

For this reason, we have chosen to pursue a continuous spatial modeling approach. This does not mean that we believe there to be continuous distributions of discrete morphogen molecules at all points in space – we know this is not true – but rather that our estimate shows at each point in space the average values, and that spread of the values, that we would expect to see if we could repeat the experiment (of performing spatialomics data collection on a tissue slice) many times. Our goal with the spatial statistical models is two-fold: 1) to estimate the unknown underlying population field

while 2) acknowledging the uncertainty in our estimate due to the sampling variability present in the observed dataset.

A general approach to spatial hierarchical modeling

We formulate our model in a generalizable Bayesian hierarchical modeling framework that can generally be thought of as having three stages:

Stage 1, data generation:	Data | Process
Stage 2, the (latent) process:	Process | Hyperparameters
Stage 3: hyperparameters	Hyperparameters | Priors

This broad framework is broadly applicable to many data collection settings where 1) data observations, which depend on some underlying process, are collected, 2) the underlying process, called the latent process since it cannot be directly observed, can be represented as a probability distribution whose parameters can be inferred from the data observations, and 3) the latent process itself may depend on some additional parameters. This simple yet powerful framework is well-established and a useful pedagogical tool for describing hierarchical modeling [23]. To make this concrete in the spatialomics setting, we could define the following three stages as follows:

1. mRNA is collected across a tissue using a spatial transcriptomics data collection technique. At each location where data is collected, only a subset of all possible mRNA at that location is collected. Furthermore, the actual levels of mRNA at any particular location stochastically vary as mRNA is created and destroyed in accordance with the unobserved cell state at and around that location. The latent cellular state cannot directly be observed, but we aim to infer something about cell state using the observable mRNA counts.

2. The cell state, the latent process, is also a stochastic process. Conditional on the cell type, cell size, cellular microenvironment and signaling networks a cell is involved in, the state of the cell follows a probabilistic distribution. The state of the cell and the previously listed cell characteristics drive the stochastic mRNA production discussed in the above description of the data collection stage.

3. To complete the Bayesian specification, some of the values that the cell state depends upon (e.g. the microenvironment of a cell) can also be viewed as stochastic processes whose parameters can be specified with prior distributions.

A general spatial statistical model for spatialomics

The ~~flexible and general purpose~~ geostatistical model described in (1) ~~can be applied to most, if not all, existing spatialomics datasets~~ is applicable to spatialomics datasets that report counts at a set of spatial locations together with a measure of local sampling effort; the conditions under which we expect it to transfer to other platforms are set out in the Discussion section. Unlike many contemporary spatialomics and genomics data analyses which use some version of standardized ("normalized") counts, we have proposed to directly model the "raw" counts from the spatialomics platform. This allows us to work with a generative model: the proposed model can be viewed as one capable of generating the data that is actually observed and collected when one uses a spatialomics platform.

Let $\boldsymbol{s} \in \mathcal{S}$ denote a spatial location over the continuous spatial domain ~~$\mathcal{S}$. We will generally use $F(\boldsymbol{s})$ to denote a continuously indexed spatial field, F,~~ $\mathcal{S}$, which here is the two-dimensional field of view of the tissue section. We write $F_i(\boldsymbol{s})$ for the count of

the ith feature (a single ligand or receptor gene) at location $\boldsymbol{s}$, ~~and we denote the spatially varying distribution of the raw counts~~ treated as a random variable, and $y_i(\boldsymbol{s})$ for an observed realization of it. We write $\mathrm{N}(\boldsymbol{s})$ for the ~~i^{th} feature as $F_i(\boldsymbol{s})$~~total counts at $\boldsymbol{s}$, that is, the number of unique molecular identifiers (nUMI) recovered at that location.

To explicitly account for the differences between the raw counts and standardized counts, we specify the *expected* counts from feature i as the product of the total ~~mRNA~~ counts, $\mathrm{N}(\boldsymbol{s})$, and a rate parameter, $\lambda_i(\boldsymbol{s})$, which represents the expected counts of the ith feature per ~~mRNA count~~ unit of total counts at location $\boldsymbol{s}$. That is, ~~$F_i(\boldsymbol{s}) = \mathrm{N}(\boldsymbol{s}) \times \lambda_i(\boldsymbol{s})$.~~ $\mathrm{E}\big(F_i(\boldsymbol{s}) \,\big|\, \mathrm{N}(\boldsymbol{s}), \lambda_i(\boldsymbol{s})\big) = \mathrm{N}(\boldsymbol{s}) \times \lambda_i(\boldsymbol{s})$, so that $\mathrm{N}(\boldsymbol{s})$ enters as an exposure term and $\lambda_i(\boldsymbol{s})$ carries the spatial signal for feature i. The conditioning matters: $\lambda_i(\boldsymbol{s})$ is itself a random field, and $\mathrm{N}(\boldsymbol{s})$ may be treated either as random or as fixed at its observed value, so this is the conditional mean of the data stage rather than a claim about the marginal expectation of F_i.

The general hierarchical model then takes the form:

$$
\begin{aligned}
&\text{Data stage} && F_i(\boldsymbol{s}) \,|\, \lambda_i(\boldsymbol{s}), \mathrm{N}(\boldsymbol{s}) && \sim \text{Poisson}\big(\mathrm{N}(\boldsymbol{s}) \times \lambda_i(\boldsymbol{s})\big) \\
& && \mathrm{N}(\boldsymbol{s}) \,|\, \lambda_N(\boldsymbol{s}) && \sim \text{Poisson}\big(\lambda_N(\boldsymbol{s})\big) \\
&\text{Process stage} && \log \lambda_i(\boldsymbol{s}) = \alpha_i + u_i(\boldsymbol{s}), \quad u_i(\cdot) \,|\, \boldsymbol{\theta}_i && \sim GP\big(0, \Sigma(\boldsymbol{\theta}_i)\big) \\
& && \log \lambda_N(\boldsymbol{s}) = \alpha_N + u_N(\boldsymbol{s}), \quad u_N(\cdot) \,|\, \boldsymbol{\theta}_N && \sim GP\big(0, \Sigma(\boldsymbol{\theta}_N)\big) \\
&\text{Prior stage} && \boldsymbol{\gamma}_i \equiv (\alpha_i, \boldsymbol{\theta}_i), \quad \boldsymbol{\gamma}_N \equiv (\alpha_N, \boldsymbol{\theta}_N) && \sim \pi_\gamma.
\end{aligned}
\tag{1}
$$

~~where we have modeled~~ Each symbol is defined as follows. In the data stage, the counts of feature i at location $\boldsymbol{s}$ ~~, $F_i(\boldsymbol{s})$, as a Poisson distribution with rate~~ are Poisson with mean equal to the total counts at $\boldsymbol{s}$ ~~, $N(\boldsymbol{s})$,~~ multiplied by the ~~density~~ rate $\lambda_i(\boldsymbol{s})$. In the process stage, $\log \lambda_i(\boldsymbol{s})$ is decomposed into a feature-specific scalar intercept, α_i, which sets the overall level of feature i ~~, λ_i, which is log-linked to a linear predictor consisting of a spatially structured Gaussian Process, GP_i with~~ across the field of view, and $u_i(\boldsymbol{s})$, a zero-mean Gaussian process over $\mathcal{S}$ which carries all spatial variation in the rate. The notation $GP\big(0, \Sigma(\boldsymbol{\theta}_i)\big)$ means that any finite collection of values of u_i is multivariate Gaussian with mean zero and covariance matrix $\Sigma(\boldsymbol{\theta}_i)$, whose $(\boldsymbol{s}, \boldsymbol{s}')$ entry is a Matérn covariance function evaluated at the distance between $\boldsymbol{s}$ and $\boldsymbol{s}'$. The vector $\boldsymbol{\theta}_i$ collects the two parameters of that covariance function for feature i: the range ρ_i, which is the distance at which the correlation between two locations has decayed to approximately 0.13, and the marginal standard deviation σ_i, which is the standard deviation of $u_i(\boldsymbol{s})$ at any single location. We do not estimate the Matérn smoothness. In the stochastic partial differential equation (SPDE) representation described below, we fix the order of the operator at 2 (the `alpha = 2` setting in the implementation), which corresponds to a Matérn smoothness of $\nu = 1$ in two spatial dimensions, so that $\boldsymbol{\theta}_i = (\rho_i, \sigma_i)$ and nothing further. Note that this order parameter is unrelated to the intercepts α_i and ~~mean α_i. The nUMI, $N(\boldsymbol{s})$, is similarly modeled. The Bayesian model~~ α_N, and we avoid writing it as a symbol for that reason. The symbols $\lambda_N(\boldsymbol{s})$, α_N, $u_N(\boldsymbol{s})$, $\boldsymbol{\theta}_N = (\rho_N, \sigma_N)$ and $\Sigma(\boldsymbol{\theta}_N)$ are the corresponding objects for the latent model of the total counts. The Bayesian specification is completed by ~~specifying the hyperprior ,~~ the hyperprior π_γ ~~, over all all hyperparameters, $\boldsymbol{\gamma}_i$. Specifically, we~~ . Note that $\boldsymbol{\gamma}_i$ is feature-specific, one per modeled feature, whereas $\boldsymbol{\gamma}_N$ belongs to the single total-count model and is shared across features; the original manuscript collected all four into one feature-subscripted vector, which was not correct.

The status of the latent total-count model. The second line of the data stage, and with it every symbol carrying an N subscript, specifies a *latent* model for the total counts. We retain it in (1) because it is the general form of the framework and because we did fit it; the per-feature outputs of those joint fits are the panels in S3 File. However, no result reported in this manuscript uses it. In every fit reported in the figures and in Table 1, $\mathrm{N}(\boldsymbol{s})$ is the *observed* nUMI at location $\boldsymbol{s}$, entered as a fixed and known offset rather than modeled, so the reported model is the first data-stage line together with the first process-stage line alone. We should be clear about what using the observed nUMI as an offset does and does not assume. It is not a claim that $\mathrm{N}(\boldsymbol{s})$ is a deterministic quantity. The total counts at a location are themselves the outcome of a stochastic process, both in the underlying transcriptional activity of the cells sampled there and in the capture and sequencing that subsample it. Using the observed total as an offset means *conditioning* on the realized value rather than modeling the process that produced it, which is a modeling choice, not a statement about the physics.

We have quantified what that choice costs, comparing the 23 features fitted under both specifications. The fitted feature fields agree closely in spatial pattern: the Pearson correlation between the joint and offset fits ranges from 0.986 to 0.997, with a median of 0.995. They differ modestly in level, the mean absolute difference between the two fitted fields being 5.1% to 13.5% of the mean fitted count for that feature, with a median of 6.6%. The posterior *uncertainty* is essentially unchanged: the ratio of posterior standard deviations between the two fits has a median of 0.998 across features, and the ratio of 95% credible interval widths a median of 0.965, with the joint model giving wider intervals for 10 of the 23 features and narrower for 13. Modeling $\mathrm{N}(\boldsymbol{s})$ as latent therefore neither propagated appreciable extra uncertainty into the feature fields nor materially sharpened them.

We adopted the offset specification for the full feature set on that basis: it preserves the spatial structure the downstream analyses use and carries the same uncertainty, while being simpler and substantially cheaper to fit. We would not claim the two specifications are interchangeable for a question about absolute magnitude, and we would expect the joint model to matter more at lower sequencing depth, where the observed total is a noisier summary of local transcriptional activity than it is here.

Priors. We set independent wide Gaussian priors on the intercepts α_i and α_N~~, and we specified a~~. On the Matérn parameters we place the joint penalized complexity prior ~~on the range~~ of Fuglstad et al. [24], which is specified by one tail probability for each component. For the analyses reported here we set $P(\rho_i < 500) = 0.95$ and ~~variance parameters of each of the two Matérn covariance [24]~~$P(\sigma_i > 0.1) = 0.05$, identically for every feature i, with the range expressed in the raw coordinate units of the Stereo-seq chip. Five hundred coordinate units is ten bins at the bin-50 resolution used throughout, and about one twentieth of the width of the field of view, so the range prior places most of its mass on fields that vary over distances well short of the field of view, and the marginal standard deviation prior shrinks u_i toward a constant field unless the data indicate otherwise. A sum-to-zero constraint is imposed on each fitted field so that α_i and $u_i(\boldsymbol{s})$ are separately identifiable.

From the model to the implementation. The Gaussian processes are not represented by a dense covariance matrix. We use the SPDE representation [25], in which a Matérn field is approximated by a Gaussian Markov random field defined on the nodes of a triangulated mesh over $\mathcal{S}$; $\Sigma(\boldsymbol{\theta}_i)$ is therefore realized in the code as a sparse precision matrix indexed by mesh nodes rather than by observation locations. The mesh used for the reported analyses has approximately 10,200 nodes covering the

26,000 observation bins. Inference is by integrated nested Laplace approximation [26], using the `R-INLA` and `inlabru` packages, and posterior summaries and predictive scores are computed from 1,000 posterior samples per feature. Because the correspondence between the notation above and the analysis code is not otherwise obvious, S9 Table gives a row for each mathematical object in (1), naming the script and the code object that implements it.

Due to the large number zeros often present in transcriptomics datasets, we also tested variations on the modeling framework wherein we replaced the Poisson likelihood for the ith feature with either a zero-inflated Poisson (ZIP) process or a zero-adjusted Poisson (ZAP) process. Both of these alternatives allow for a spatially varying probability of observing no counts. In these variants, the data generating mechanism can then viewed as first flipping a coin to determine if any counts from feature i will be observed, followed by a data observation from a Poisson process if the coin flip succeeds. Additional detail can be found in the Comparison of 3 data models section.

Quantification of categorical spatial structure

To assess the impact of our modeling choices, we desired a way of quantifying the spatial coherence of our final output clusters, represented as a categorical variable. To this end, we employed ELSA (entropy-based local indicator of spatial association), which is applicable to categorical variables and does not depend on the number of categories [27]. ELSA is calculated by multiplying a spatially localized (scaled) Shannon entropy calculation with a localized measure of spatial dissimilarity to produce a measure of spatial association. Each of these components are bounded between 0 and 1, ensuring that ELSA is also bounded between 0 and 1. Locations with high values of ELSA indicate spatial regions where the variable of interest is less structured and spatially disparate while low values of ELSA suggest higher levels of spatial coherence within the local neighborhood.

Computational experiments

Dataset selection

To demonstrate our approach in detail, we ~~leverage~~ use data from Chen et al. 2022 [28], the Mouse Organogenesis Spatiotemporal Transcriptomic Atlas (MOSTA), a spatial time-course atlas of mouse embryogenesis acquired ~~using~~ with the spatial transcriptomics platform Stereo-seq. ~~This dataset is beautiful~~The material analyzed here is therefore a mouse embryo section drawn from a public atlas. No new tissue was collected for this study, no human or patient material is involved, and we did not perform the tissue processing or sequencing ourselves; readers should consult the original publication for the experimental protocol. We selected this resource because it is publicly available, information-rich, ~~easy to interpret , and highly~~ straightforward to interpret anatomically, and relevant to both developmental and regenerative biology. ~~We~~

Stereo-seq captures polyadenylated RNA on a chip patterned with DNA nanoball spots, each bearing capture probes that carry a coordinate identity barcode encoding the spot's position, a unique molecular identifier, and a poly-T tail. Spots are approximately 220 nm across, with a center-to-center spacing of either 500 or 715 nm depending on the chip variant; the source publication does not state which variant carried the embryo sections, so we report positions below in spot units rather than converting them to physical distances. Because the spacing is far finer than a cell, individual spots are very sparse, and counts are conventionally aggregated into square bins before analysis. We follow that practice.




Fig 2. Data parameters. The 1st columns shows total counts within the selected field of view from [28]. The 2nd column shows ligand information, and the 3rd column shows receptor information. For latter two columns, the 1st row is total ligand or receptor counts, the 2nd row is total ligand or receptor features, and the 3rd row is fraction of the total counts that are ligand or receptor counts.

From this atlas we selected a single section from embryonic day 16.5, section identifier `E16.5_E1S3`, and within it a single field of view spanning x from 15000 to 25000 and y from ~~one of the Day 16 timepoints~~ 11500 to 18000 in raw chip coordinates (Fig 2). One unit of those coordinates is one nanoball spot, so the field of view is 10,000 by 6,500 spots; its size in millimetres depends on which chip variant was used, which the source publication does not specify. The field of view was chosen to be visually recognizable to both experts and non-specialists and to contain several distinct organs, allowing close study of ~~key~~ morphogenic field interactions during development. At the bin size used throughout (bin 50, meaning a 50×50 block of capture spots), this field of view comprises 26,000 spatial locations, each of which is an observation in the models below.

Dataset preprocessing

Raw count data were loaded into `R` and ~~a subregion of the embryonic tissue was chosen to be visually recognizable by both experts and non-specialists~~restricted to the field of view described above. We then filtered the transcriptomic features to only those measured within this boundary ~~,~~and, following the workflow from the original publication, collapsed the ultra-high resolution, sparse data ~~to different-sized spatial bins~~ into square spatial bins at three sizes (10, 25, and 50). All ~~the~~ results shown in the figures of this manuscript use the bin-50 data, which divides the field of view into a 200×130 grid of 26,000 bins; S1 File documents this raw-to-bin step and its outputs. Bin 50 was chosen as the coarsest of the three, and therefore the least sparse per observation; the choice of binning is a modeling decision rather than a property of the platform, and results at other bin sizes are not reported here.

To identify ligand and receptor genes, we cross-referenced the mouse version of the FANTOM5 database [29] against the feature names of the binned data, yielding 974

distinct genes that were either ligands or receptors. We further refined this gene list to only include ligands or receptors showing at least two counts in at least one ~~cell~~bin, and at least one-quarter-of-one-percent of the ~~total cells~~ bins in the field of view expressing the gene at any non-zero level (~~see Supplement 2 for~~ S2 File shows the visual outputs which were used to decide this threshold). This final step of feature selection yielded a curated list of 918 individual ligand or receptor genes~~for modeling, which, based on the pairings in the FANTOM5 data base, would allow downstream computation of 1481 unique ligand-receptor interaction fields~~.

Dataset modeling

We individually modeled the spatial ~~fields~~ field of the counts of ~~the 918 features~~ each curated feature using a spatial Gaussian process model with Poisson observations, as defined in (1)~~. The model was~~, with the observed nUMI at each bin as a fixed offset. The models were fit using `R-INLA` and `inlabru`, and predictions from these models were then used in downstream analyses.

Not every curated feature yielded a usable field. Of the 918 genes on the curated list, three (*a*, *Calcb* and *Cckar*) were excluded before fitting, and three further genes (*Fgf1*, *Fzd7* and *Trhr*) did not produce a converged fit. All downstream analyses therefore use the 912 features with fitted spatial fields. Of the FANTOM5 ligand-receptor pairings, 1477 have both partners among these 912 features, and these 1477 pairs define the interaction fields analyzed in the remainder of the manuscript. They are not 1477 numerically distinct fields. The common-minimum operator described below returns whichever of the two partner fields is smaller, so pairs that share a partner which is the smaller of the two at every location yield identical fields; seven pairs of the ligand GDF9 with different receptors, for example, give a single field. The 1477 pairs yield 1449 numerically distinct fields under that operator.

Comparison of 3 data models

We experimented with three different likelihood models: a Poisson likelihood, a zero-inflated Poisson (ZIP) likelihood, and a zero-adjusted Poisson likelihood (ZAP). The ZIP and ZAP models provide extra flexibility to acknowledge that the raw count data is often very sparse, with many spots returning a measurement of 0 for any particular feature:

Writing $\mu_i(\boldsymbol{s}) = \mathrm{N}(\boldsymbol{s})\,\lambda_i(\boldsymbol{s})$ for the Poisson mean of (1), so that the observed total counts enter as an exposure term in all three cases, the likelihoods are

$$\begin{aligned}
\text{Poisson: } P\left(F_i(\boldsymbol{s}) = y \mid \mu_i(\boldsymbol{s})\right) &= \text{Poisson}\big(y \mid \mu_i(\boldsymbol{s})\big) \\
\text{ZIP: } P\left(F_i(\boldsymbol{s}) = y \mid \mu_i(\boldsymbol{s}), p_i(\boldsymbol{s})\right) &= p_i(\boldsymbol{s})\mathcal{I}_{y=0} + (1 - p_i(\boldsymbol{s}))\text{Poisson}\big(y \mid \mu_i(\boldsymbol{s})\big) \\
\text{ZAP: } P\left(F_i(\boldsymbol{s}) = y \mid \mu_i(\boldsymbol{s}), p_i(\boldsymbol{s})\right) &= p_i(\boldsymbol{s})\mathcal{I}_{y=0} + (1 - p_i(\boldsymbol{s}))\text{Poisson}_{>0}\big(y \mid \mu_i(\boldsymbol{s})\big),
\end{aligned} \tag{2}$$

where $p_i(\boldsymbol{s})$ is a spatially varying probability of a 0 observation for feature i, which is modeled with ~~a~~ its own spatial Gaussian random field using a ~~logit-link~~logit link, $\text{Poisson}_{>0}$ denotes the zero-truncated Poisson, and $\mathcal{I}_{y=0}$ is an indicator function which takes a value of 1 if $y = 0$ and takes a value of 0 otherwise. The ZIP and ZAP models differ in that the probability of a 0 in the ZAP model is directly governed by ~~$p_i(\boldsymbol{s}))$~~ $p_i(\boldsymbol{s})$ and allows for the probability of a 0 to be less than a standard Poisson, whereas the ZIP models can only increase the probability of a 0. ZAP models are typically selected when the process for a 0 occurrence differs substantially from the process driving the non-0 values.

At each observed location, $\boldsymbol{s}_j$, model performance was assessed ~~using three distinct prediction scores combining the observed counts for~~ by comparing the observed count of feature i at ~~$\boldsymbol{s}_j$~~that location, $y_i(\boldsymbol{s}_j)$~~and the features of the~~, against the posterior predictive distribution of $F_i(\boldsymbol{s}_j)$. Writing $\mathrm{E}(\cdot|\text{data})$, $\mathrm{var}(\cdot|\text{data})$ and $\mathrm{med}(\cdot|\text{data})$ for the mean, variance and median of that posterior predictive distribution~~of $F_i(\boldsymbol{s})$:~~, the four location-level scores we used are

$$
\begin{aligned}
AE_i(\boldsymbol{s}_j) &= \left|y_i(\boldsymbol{s}_j) - \mathrm{med}\big(F_i(\boldsymbol{s}_j)|\text{data}\big)\right| \\
SE_i(\boldsymbol{s}_j) &= \big(y_i(\boldsymbol{s}_j) - \mathrm{E}\big(F_i(\boldsymbol{s}_j)|\text{data}\big)\big)^2 \\
DS_i(\boldsymbol{s}_j) &= \frac{\big(y_i(\boldsymbol{s}_j) - \mathrm{E}\big(F_i(\boldsymbol{s}_j)|\text{data}\big)\big)^2}{\mathrm{var}\big(F_i(\boldsymbol{s}_j)|\text{data}\big)} + \log\left(\mathrm{var}\big(F_i(\boldsymbol{s}_j)|\text{data}\big)\right) \\
LS_i(\boldsymbol{s}_j) &= -\log\left(P\big(F_i(\boldsymbol{s}_j) = y_i(\boldsymbol{s}_j)|\text{data}\big)\right).
\end{aligned}
$$

The ~~Dawid-Sebastiani (DS) score is a proper scoring rule combining the predictive mean and variance, $E(F_i(\boldsymbol{s})|\text{data})$ and $V(F_i(\boldsymbol{s})|\text{data})$, respectively~~absolute error (AE) and the squared error (SE) are point-forecast measures: they compare a single summary of the predictive distribution against the observation and ignore its spread. Where we are free to choose that summary we use the posterior predictive median, because it is the more stable summary for the skewed, low-count predictive distributions that arise here, and because it is the summary that minimizes expected absolute error. The squared error ~~(SE~~and the Dawid-Sebastiani score are defined in terms of the predictive mean and variance, so for those we use the mean, as their definitions require. This is why AE alone is computed against the median. The Dawid-Sebastiani score (DS) is a proper ~~score for the expectation~~scoring rule combining the predictive mean and variance, and the (negated) ~~Log~~ logarithmic score (LS) is a strictly proper ~~score [30].~~ scoring rule which uses the entire predictive distribution [30]. All four are negatively oriented, so that a smaller value indicates better performance; note that DS may be negative, because it contains a log-variance term, so "smaller" means more negative for that score.

These location-level scores are reduced to the four per-feature quantities reported in Table 1. For feature i, the mean absolute error is $MAE_i = \mathrm{mean}_j\, AE_i(\boldsymbol{s}_j)$, the root mean squared error is $RMSE_i = \sqrt{\mathrm{mean}_j\, SE_i(\boldsymbol{s}_j)}$, the mean Dawid-Sebastiani score is $MDS_i = \mathrm{mean}_j\, DS_i(\boldsymbol{s}_j)$, and the mean log score is $MLG_i = \mathrm{mean}_j\, LS_i(\boldsymbol{s}_j)$, each averaged over the observation locations $\boldsymbol{s}_j$. Table 1 then reports the average of each of these across features.

Quantitative optimization of ligand-receptor convolution

We experimented with multiple different approaches to ligand-receptor convolution. In all approaches studied, we did not consider ligand nor receptor sequestration due to binding, nor did we consider cross-mechanism inhibition or competitive interference, which can be the focus of future studies. We tested product, geometric mean, and a new approach labeled 'common minimum convolution', which ~~computed the maximum possible~~ computes an upper bound on the number of local binding events ~~, assuming~~ that the estimated ligand and receptor levels would imply, under the assumption that all ligand and all receptor bind to one another in a 1:1 fashion. As with every quantity in this paper, that bound is model-derived from transcript counts rather than measured. We also explored a prototype approach based on chemical kinetics, using the rate equation with a tunable dissociation constant to model local steady-state ratio between bound and unbound ligand.

Our goal was to learn which methods yielded the most fine-grained and spatially organized clusters by looking for methods that maximized the number of clusters while

minimizing the spatial entropy as quantified by ELSA (see the Quantification of categorical spatial structure section). We therefore performed a targeted optimization experiment to identify which combination of parameters, models, and data inputs, all else being equal, ~~yielded an output that maximized our unbiased identification~~ resolved the largest number of distinct local microenvironments while minimizing inter-cluster spatial entropy. ELSA was calculated locally and averaged across the modeling region to produce a single comprehensive score. This provided a means to visualize the boundaries between clusters, assessing how well they segregate in space, and to rank each computational approach by both number of clusters identified (all other parameters being equal) and the global spatial entropy of cluster segregation.

Quantifying the relationship between spatial structure and hyperfield dimensionality

We use *hyperfield* throughout for the full collection of inferred ligand-receptor interaction fields treated as a single high-dimensional object, and *hyperfield dimensionality* for the number of those fields entering an analysis at once.

We used the results from a specific optimized workflow (Raw Counts - Poisson Model - Common Minimum Convolution) to ~~quantify the full relationship between~~ examine how spatial domain architecture ~~and signaling hyperfielddimensionality,~~ varies with the dimensionality of the signaling hyperfield, meaning the number of inferred interaction fields entering the clustering at once. We did this because contemporary biologic theory posits that the complex spatial domains observed in tissues are likely a multivariable function of high-dimensional signal combinations rather than of a small set of select mechanisms. We therefore performed an experiment in which we iteratively increased the number of individual features considered for latent-space embedding and clustering~~and observed standardized unbiased outputs~~, holding the rest of the workflow fixed. This sweep varies one quantity, the feature count, and so characterizes that one dependence rather than the relationship in full. Individual ligand-receptor mechanism features were first ranked based on system-scale variance, and then a for-loop was written to embed and cluster the data, with otherwise fixed algorithmic parameters, using increasingly many ligand-receptor features (10, 50, 100, ~~150, ...,~~ and then in steps of 50, up to all 1477 pairs). The input to this sweep inadvertently contained a second copy of four interaction fields (PDGFC and PDGFD each paired with PDGFRA and PDGFRB), read in from duplicated input files, so its full-depth clustering used 1481 ~~)~~columns. We expect the effect to be small, since four duplicated columns are scaled alongside more than a thousand others, but we have not measured it. Ligand-receptor connectivity field data were scaled (though, importantly, not sum-normalized first) and processed without any principal-component analysis. Latent-space neighborhood graph construction and clustering parameters were held constant to allow isolation of the effect of feature number. ELSA was run in all instances, with a scan across the local distance parameter. All outputs from each step were stored for downstream visualization and analysis.

Because a variance-ordered sweep admits the same fixed set of features at each depth, it confounds the number of fields entering the clustering with which fields they are. To separate the two, we repeated the sweep admitting uniformly random subsets of the 1449 distinct fields, at depths of 10, 50, 150, 350, 650 and 1000 fields, with 19 independent random subsets at each depth (18 at depth 50, where one run failed for a technical reason unrelated to the data), using the same scaling, neighbor-graph and clustering steps and parameters, and scoring each clustering by mean ELSA at radius 50.

Results

Raw counts, rather than normalized counts, are ~~optimal~~ the appropriate input for spatial modeling

Our experiments first ~~revealed~~ indicated that commonly-applied count-normalization approaches ~~create significant~~ can introduce topological artifacts when applied to these spatial data, a pattern also noted by others [31]. Low individual-feature counts can be transformed into local hotspots when the raw values are count-normalized within spots of low total RNA density. See Fig 3 for a demonstration with a single feature~~(~~, Cd44~~);~~ ~~similar comparisons between raw data vs. count-normalization for all features studied can be observed in Supplement 3.~~ . We chose Cd44 for clarity rather than because it is special: the effect is visible there against an unambiguous anatomical landmark, the developing spine, which makes the raw and count-normalized panels easy to compare side by side. It is one clean example among many, not the only feature that shows the effect. The same comparison for a further 23 selected features is in S3 File, and the per-feature outputs for the whole curated set are in S2 File, so a reader can check how general the pattern is rather than taking one panel on trust. Because we aimed to model raw ligand-receptor *binding density* within tissues, the results from this experiment made clear to us that it was most physically appropriate to convolve continuous ligand and receptor fields which estimate *raw* ligand and receptor topology, rather than fields which estimate the commonly-used *count-normalized* topology. Nevertheless, we decided to carry forward both approaches in parallel experimentally, so that we could identify and compare project-relevant differences downstream. As a secondary result, ~~we also learned form~~ these early experiments ~~that spatially modeling $N(\boldsymbol{s})$~~also compared spatially modeling $\mathrm{N}(\boldsymbol{s})$, the nUMI, ~~in addition to~~ jointly with a ligand or receptor feature ~~, $F_i(\boldsymbol{s})$, didn't practically alter the outputs, and~~ $F_i(\boldsymbol{s})$, against treating the observed nUMI as a fixed offset. Across the 23 features fitted both ways, the two give fitted fields with almost identical spatial pattern (Pearson correlation 0.986 to 0.997, median 0.995), modestly different levels (mean absolute difference 5.1% to 13.5% of the mean fitted count, median 6.6%), and essentially identical posterior uncertainty (median ratio of posterior standard deviations 0.998, of 95% interval widths 0.965). Since the downstream analyses depend on spatial pattern rather than absolute level, and since modeling the total jointly neither added nor removed appreciable uncertainty, the remainder of the results were run using the nUMI directly, as measured in the data, which is also much cheaper to fit.

Poisson likelihood model outperforms ZIP and ZAP models

We compared three ~~main~~ spatial modeling approaches: ~~POI~~Poisson, ZIP, and ZAP (see ~~Methods~~Comparison of 3 data models). We found that the relative simplicity of the spatial Poisson model proved most suited, in these particular exercises, to the ~~general~~ task of estimating feature counts. Table~~1 provides the average mean absolute errors, root mean square errors, Dawid-Sebastiani scores,~~ 1 gives the four prediction scores by likelihood.

Of the four scores, we prioritized the two proper scoring rules, MDS and MLG, over the point-forecast measures MAE and RMSE, because what this framework offers downstream is a predictive distribution with quantified uncertainty rather than a point estimate, and ~~log scores, split out by likelihood and averaged over all observations and all 918 features.~~ only the proper scoring rules assess that distribution as a whole. In this particular comparison the choice of criterion does not change the conclusion, since the Poisson model gives the best average value on all four scores. This is not evident from the parenthetical percentages, which fall below 100% for




Fig 3. Raw vs. count-normalized topology. Example comparison between raw count topology vs. count-normalized topology for a single feature (Cd44). 1st column is observed Cd44 counts, 2nd column is observed Cd44 counts divided by the total counts. 3rd column is the total counts. 1st row is raw data, 2nd row are estimates from the spatial model, 3rd row is the residual (difference between model and raw data.) ~~Count-normalization causes significant~~ In this example, count-normalization substantially changes ~~to~~ the spatial topology~~;~~. On this basis, and on the comparable panels for 23 further features in S3 File, we ~~have concluded that~~ avoid normalization ~~should be avoided~~ prior to connectivity computations in the work reported here.

MDS. That is an artifact of expressing a negative score as a ratio: because MDS is negative, a ratio below 100% corresponds to a score nearer zero and therefore worse, and the raw MDS values (−2.15 for Poisson against −1.78 for ZIP and −1.77 for ZAP) should be compared directly.

Nor is the average driven by a small number of extreme features. Comparing the three likelihoods feature by feature, across all 912 features and without the score filter applied to the table averages, the Poisson model gave the lowest score for 814 features on MAE, 880 on RMSE, 734 on MDS, and 829 on MLG, and was best on all four scores simultaneously for 645 of the 912. Predictive plots showed that the empirical distribution of the ~~PIT scores~~ probability integral transform (PIT) values from the Poisson models was routinely closest to a uniform distribution, which is the distribution a calibrated predictive model would produce, when compared to the empirical ~~CDF~~ distribution of the PIT ~~scores~~ values from the ZIP and ZAP models. Spatial residual plots revealed that the poorer average performance by the ZIP and ZAP models appears attributable to oversmoothing as evidenced by the predictions and residuals. See Fig 4 for comparison between models for a single feature~~. See Supplement 2~~, and S4 File for corresponding plots for all features.

We read this as evidence that the extra flexibility of a zero-inflated or zero-adjusted likelihood was not needed at the sparsity of this dataset at bin-50 resolution, not as a general result about count likelihoods for spatialomics. A platform or binning choice that produced substantially sparser data could well favor a zero-inflated likelihood, and the framework accommodates one.

Table 1. ~~Model comparison~~Comparison of the Poisson, ZIP and ZAP likelihoods. ~~Mean Absolute Errors~~Abbreviations: MAE, ~~Root Mean Square Errors~~mean absolute error; RMSE, root mean squared error; MDS, mean Dawid-Sebastiani ~~Scores~~score; MLG, ~~and Log Scores~~mean log score; ZIP, ~~by likelihood~~zero-inflated Poisson; ZAP, zero-adjusted Poisson. Each score is first averaged over ~~all observations~~ the observation locations within a feature, and ~~all 918 modeled ligand/receptor~~ the table reports the average of those per-feature values across features; the scores are defined in the Comparison of 3 data models section. All four scores are negatively oriented, so the smallest value in a column is the best. MDS is negative because it contains a log-variance term. The ~~average percentage difference~~ Poisson model gives the best value on all four scores. Values in parentheses are the ~~values, relative to~~ mean across features of the ~~performance~~ ratio of that model's score to the Poisson score *for the same feature*, expressed as a percentage, so 100% denotes a model whose average per-feature score equals Poisson's. Note that this is an average of ratios and carries no dispersion, so a value near 100% indicates comparable average performance rather than statistical indistinguishability. These are ratios rather than differences, and they are not the ratio of the two averages printed in the table. A value above 100% therefore indicates worse performance than Poisson. No percentage is given for MDS: that score is negative, so a ratio of two negative numbers is not interpretable on the same scale as the others, and we give only the raw MDS values, which should be compared directly. Poisson is lowest, and therefore best, on all four. Averages are taken over the 912 features with converged fits, excluding feature-by-model combinations whose scores took extreme values indicating nonconvergence. That exclusion removes 4 of the 2,736 combinations from the absolute averages and 7 from the parenthetical ratios, in ~~parentheses~~both cases affecting only the ZIP row, which is therefore averaged over 908 features for the absolute values and 905 for the ratios.

Model	MAE	RMSE	MDS	MLG
Poisson	0.21	0.42	-2.15	0.31
ZIP	0.29 (111%)	0.54 (116%)	-1.78 ~~(85%)~~	0.33 (111%)
ZAP	0.23 (111%)	0.47 (116%)	-1.77 ~~(82%)~~	0.35 (122%)

Common minimum ~~L-R~~ convolution ~~maximizes spatial structure~~gave the most clusters of the operators tested

We assessed the output spatial structure that resulted from clustering 32 distinct spatial assays, summarized in ~~Supplement 3.~~ S5 File. Three in particular are detailed in Fig 5. ~~We found that~~ Of the operators we tested, the 'common minimum' approach, which simply takes the common minimum between the ligand and receptor subunits, ~~maximizes the~~ gave the largest number of clusters ~~identifiable in the data while simultaneously minimizing spatial entropy~~ (29, against 27 for both the product and the geometric mean). Its spatial entropy was close to the lowest of the operators and well below that of the product: mean ELSA at radius 100 was 0.050 for the common minimum, 0.047 for the geometric mean, and 0.071 for the product. Of note, ~~we found that~~ the common minimum approach closely approximates the ~~binding density outputs~~




Fig 4. Likelihood model comparison. The 1st, 2nd, and 3rd rows, respectively, show the outputs from the Poisson, ZIP, and ZAP models. The 4th row shows raw data. The 1st column shows the probability of a non-zero value at that point in space. The 2nd column shows the density for that feature, i.e., counts/nUMI. The third column shows the feature counts. The top three rows in the 4th column show the residuals, the difference between estimated counts and data counts, and the bottom right panel shows the nUMI from the data, on the color scale of the total-counts column of Fig 3; the color bar beneath the 4th column applies to the residuals. The ~~POI~~ Poisson model resulted in the smallest residuals and was the best performing of the three spatial ~~model~~ models tested.

~~which are~~ outputs produced when k, the dissociation constant, is decreased to ~~0. This~~ 0, which within that kinetic model corresponds to total binding without any dissociation. Because of these results, we ~~determined this to be an optimal output of our model to use~~ carried this output forward for downstream experimentation. For full results from this experiment, see ~~Supplement 4.~~ S6 File.

Two caveats attach to this ranking. It rests on single runs of a clustering step whose run-to-run variation we have not measured directly; re-runs of the related clustering in Fig 6 differed from the published values by up to about 0.01 ELSA units (see the Controls: the domains survive a spatial null and match independent anatomy section). At radius 50 the common minimum and the product differ by 0.024, about twice that, while the common minimum and the geometric mean differ by 0.004, well within it, so those two should not be read as ranked on coherence. It is also a comparison of operators within our own pipeline on one dataset, not a general result about which convolution is correct.




Fig 5. ~~Convolution method comparison.~~Comparison of ligand-receptor convolution ~~methods~~operators.~~Product~~ Three operators for combining a ligand field with a receptor field are compared: the product, the geometric mean, and the common minimum ~~shown~~ (`m.est.prod`, `m.est.gm` and `m.est.min` in ~~1st~~the code). All use the same field of view and the same fitted fields, ~~2nd~~so the operator is the only thing deliberately varied; each clustering is a single run of a step whose exact reproducibility we have not established, so run-to-run variation may also contribute (see the Controls: the domains survive a spatial null and match independent anatomy section). (A) Inputs for four example ligand-receptor pairs (columns: WNT4-FZD6, RSPO1-LGR6, DHH-PTCH1, VEGFA-KDR): for each pair, the fitted ligand field (left) and ~~3rd column~~receptor field (right), ~~respectively~~shown without color bars because they indicate where each partner is expressed rather than serve as quantitative reads. (B) The interaction field each operator produces from those inputs, one row per operator. Absolute values ~~are~~ differ between operators and matter less ~~important~~ than the ~~relative topology~~spatial pattern, because the ~~values~~ fields are ~~generally~~ scaled ~~for use in high-dimensional analysis~~before clustering. ~~Middle of the page shows high dimensional~~ (C) The tissue domains obtained by clustering ~~with~~ each operator's full set of interaction fields under a fixed workflow (see ~~Methods~~the Quantitative optimization of ligand-receptor convolution section), one column per operator. Cluster ~~color and number is~~ colors are arbitrary ~~; cluster~~ labels, so what is comparable across columns is the number of domains and their spatial organization~~is relevant~~. This clustering first reduces the fields to their first 100 principal components, whereas Fig 6 clusters the same common-minimum fields in the full feature space and recovers 20 domains where this figure recovers 29; the counts in the two figures are therefore not directly comparable. (D) The corresponding '~~Entrograms~~entrogram', ~~made using ELSA [32], quantify~~ a map of the entropy-based local ~~indicator of~~ spatial ~~coherence. The three methods detailed here are~~ association (ELSA) [27] at radius 50 computed on the ~~top performing~~domain labels and scaled from 0 to 1; high values mark locations where domain labels are locally mixed, ~~maximizing cluster number while minimizing~~ so coherent domains appear as dark interiors bounded by thin bright borders. (E) Mean ELSA against the ELSA neighborhood radius for a subset of the representations tested. ~~The~~ (F) Mean ELSA at radius 100 against the number of clusters

High-dimensional morphogenic field analysis resolves spatial domains

To test our central hypothesis, that tissue spatial domains ~~are a direct function of signaling field hyperstructure~~reflect the structure of the high-dimensional signaling field rather than of a few individual mechanisms, we performed a targeted parameter sweep experiment in which we iteratively increased the number of signaling features, ranking them by variance, while holding all else equal. We iteratively added more features to our analysis and quantified the output number of clusters and their spatial entropy to observe changes in spatial domain architecture in both ~~UMAP and histologic space~~ histologic space and a uniform manifold approximation and projection (UMAP) embedding. Our results are summarized in Fig 6, and full results may be found in ~~Supplement 5.~~ S7 File.

~~We found the~~ In this tissue section, the recovered spatial domain architecture ~~in tissues to be a high-dimensional function of many signaling mechanisms at once. The effect is non-linear, with individual feature variance not directly predicting of the effect of adding said feature into holistic analysis. This finding supported the non-reducibility of hyperfield information. We found that clustering the data without any~~ depended on many inferred interaction fields at once rather than on a small number of them, though not in the simple form the variance-ordered sweep suggests. In that sweep (Fig 6) the number of domains fell as fields were added, from 33 at 10 fields to 24 at 150 and between 16 and 21 from 350 onward. Mean ELSA at radius 50 was not monotone: it rose from 0.143 at 10 fields to a peak of 0.220 at 100, then fell to 0.124 at full depth. Full depth was not the most coherent depth swept; five deeper depths (1200, 1250, 1350, 1400 and 1450 fields) scored between 0.109 and 0.112, although at that end of the curve differences between depths are not much larger than the run-to-run variation of the clustering (see the Controls: the domains survive a spatial null and match independent anatomy section).

The random-subset sweep separates the number of fields from their identity. With fields admitted at random, mean ELSA at radius 50 fell steadily with depth: 0.372 at 10 fields (standard deviation across the 19 subsets 0.044), 0.240 at 50 (0.034), 0.168 at 150 (0.029), 0.134 at 350 (0.032), 0.135 at 650 (0.024) and 0.120 at 1000 (0.012). Most of the improvement arrives by a few hundred fields, with slower change thereafter. The rise and fall seen in the variance-ordered sweep does not appear when fields are admitted at random, which suggests that it reflects the order of admission rather than the number of fields. The two sweeps were run separately, and the random-subset sweep drew from the 1449 distinct fields, so we compare the shapes of the two curves rather than their values at matching depths. All of these clusterings were run without principal-component ~~analysis resulted in the greatest spatial complexity and coherence. Clustering the tissue within the full 1, 481-dimension feature-space maximized spatial complexity and coherence, allowing the resolving of subtle~~ analysis (PCA); we did not run a matched comparison with PCA, and so make no claim about whether omitting it helps. At full depth the domain map resolves local detail, such as inter-organ fascial planes, that ~~was unobserved when all features were not included or when dimensional-reduction was employed via PCA~~is not apparent at the shallowest depths.

We ~~closely observed~~ examined the inferred hyperfield structure in a small sub-region of the embryo, outlined by a ~~dotted-white boundary~~ dashed white box in Fig 6. Selected feature-level topologies (individual inferred ligand-receptor ~~binding density field~~interaction fields) have been rendered as three-dimensional surfaces at the bottom of Fig 6. HTML files for ~~all 1,481 fields~~ every interaction field within this region, allowing user-interactive zoom, value estimation and rotation in space, are available in ~~Supplement 6.~~ S8 File.




Fig 6. ~~High-dimensional spatial~~ Tissue domains at increasing numbers of inferred interaction fields. ~~Top~~ Columns: clustering of the inferred interaction fields at five ~~rows show increasing spatial domain definition~~ feature depths, with features entering in order of decreasing standardized variance: 10, 150, 650 and ~~organization as~~ 1,300 pairs, and all 1,477 ligand-receptor pairs (1,449 numerically distinct fields; see the Materials and methods section). Every other algorithmic parameter is held constant and no principal-component analysis is applied, so the number of ~~morphogenic interaction~~ features is the only thing deliberately varied between columns; each column is a single run of a clustering step whose exact reproducibility we have not established, so run-to-run variation may also contribute (see the Controls: the domains survive a spatial null and match independent anatomy section). Top row: uniform manifold approximation and projection (UMAP) embedding of the spatial bins, colored by cluster. Second row: the same cluster labels in tissue space, which are the spatial domains, with the number of domains above each map; cluster colors are arbitrary. These domains come from clustering the full feature space, whereas the operator comparison in Fig 5 first reduces to 100 principal components and recovers 29 domains from the same common-minimum fields, so the counts in the two figures are ~~increased~~not directly comparable. ~~Endpoint~~ Third row: the corresponding 'entrogram', a map of the entropy-based local indicator of spatial association (ELSA) [27] at radius 50 computed on the domain ~~architecture is a detailed function~~ labels, in which coherent domains appear as dark interiors separated by bright boundaries. Fourth row, left: mean ELSA at radius 50 against the number of ~~all 1~~interaction fields entering the clustering, ~~481 unique signaling mechanisms~~for the variance-ordered sweep shown above (yellow) and for random subsets of the 1,~~treated without principle-component analysis~~ 449 distinct fields (~~see Methods~~blue; mean and one standard deviation over 19 subsets per depth, 18 at depth 50). Middle: the standardized variance of each interaction field by rank, which sets the order in which fields enter. Right: ~~number of domains against number of fields, for the same two sweeps. Dotted lines mark the~~ five depths shown above. As features are added in variance order the domains become fewer, not more numerous: 33 at 10 features, 24 at 150, and between 16 and 21 from 350 onward. Mean ELSA rises from 10 features to a peak between 100 and 200 and then falls; with random subsets it falls steadily, which suggests that the initial rise reflects the order in which features enter rather than their number. Bottom two rows~~show 8~~: eight selected ~~morphogenic~~ interaction fields ~~of biologic interest, represented~~ rendered as three-dimensional ~~density plots~~

Controls: the domains survive a spatial null and match independent anatomy

The experiments above show that clustering the full signaling hyperfield yields spatially coherent domains, but not that those domains reflect structure present in the tissue. Two properties of the pipeline could in principle manufacture coherent domains on their own: the spatial model smooths, and the clustering operates in a space of more than a thousand dimensions. We therefore ran two controls aimed at these two properties, and a third, described after them, aimed at the ligand-receptor pairing. In each, the domain-generating pipeline is held fixed between the observed and the control runs, so that any difference is attributable to the manipulation alone. Spatial coherence is quantified throughout with ELSA at neighborhood radii of 50 and 100 units (see the Quantification of categorical spatial structure section); ELSA is negatively oriented, so lower values indicate more spatially coherent domains. The first two controls are summarized in S10 Fig.

A coordinate shuffle destroys the domains

To test whether the spatial smoother manufactures coherent structure from data carrying no spatial signal, we permuted the assignment of the observed counts to spatial locations. This destroys spatial structure while preserving the observed count distribution exactly. We then refit the spatial model for every one of the 912 features from scratch on the permuted data, and passed the refit fields through the identical convolution, clustering, and scoring pipeline. Identical here means identical between the observed and the permuted data, which is what the comparison requires. It is not identical to the clustering behind Fig 6: because each permutation already required 912 model fits, both arms of this control were clustered with a faster implementation of the same two steps, a nearest-neighbor graph built with RcppHNSW and Louvain clustering in igraph, in place of Seurat. ELSA values from the two implementations are not on the same scale, so the values in this section should be compared with each other and not with those reported for Fig 6. Refitting, rather than reshuffling fields that were fitted on the real coordinates, is what makes this a test of the smoother: the model sees only spatially meaningless data. Because each permutation requires 912 model fits, the experiment was run on a high-throughput computing cluster, with 49 permutations.

The observed data yielded 20 domains, with an ELSA at radius 50 of 0.075. The 49 coordinate-shuffled replicates yielded between 4 and 12 domains, with ELSA at radius 50 averaging 0.470 and ranging from 0.412 to 0.583. The null mean is 6.3 times the observed value, and the lowest of the 49 shuffled replicates is still 5.5 times it, so the observed and null values do not overlap. The permutation p-value is 0.020, which is the smallest value attainable with 49 permutations; the test is therefore limited by the number of replicates we could afford rather than by any overlap between the observed value and the null distribution.

Fitting a spatial Gaussian process to spatially permuted data does not, then, produce coherent tissue domains: it produces a small number of incoherent fragments. Because the permuted data pass through a pipeline of identical dimensionality, this speaks to the smoothing and the dimensionality explanations at once.

The domains correspond to independently annotated anatomy

As a positive control, we compared the emergent domains against the anatomical annotations published with the source dataset [28] for the same tissue section. These annotations were produced by a different group, by a different method, and without

any reference to our labels, which our pipeline in turn never sees; in that sense they are an independent reference. All 26,000 modeled bins matched an annotated bin.

For completeness about what the reference is: Chen et al. derived these labels by spatially constrained clustering, combining a nearest-neighbor graph over spatial position with one over expression, and then labeling each resulting cluster by hand from its marker genes against a reference embryo atlas. The comparison is therefore between two partitions of the same tissue section, which is unavoidable, since a concordance check is only meaningful against annotations of the same specimen. The two partitions are nonetheless built from very different features: theirs from binned expression across all genes, ours from the inferred interaction fields of 1477 ligand-receptor pairs, and the manual labeling step brings in external anatomical knowledge from outside the dataset entirely.

Agreement was well above chance. The adjusted Rand index between our domains and the anatomical labels was 0.401 and the normalized mutual information was 0.604, against a permutation null whose mean adjusted Rand index was 0.00003 and whose maximum over 499 permutations was 0.001, giving $p = 0.002$, again the floor for the number of permutations used. The mean size-weighted purity of the emergent domains with respect to the anatomical labels was 77.6%. Several individual domains recover recognized structures cleanly: spinal cord at 97% purity, liver at 96%, one heart domain at 99%, and meninges at 92%, with lung, muscle, thymus, and cartilage primordium also recovered.

We would not read an adjusted Rand index of 0.401 as showing that the signaling domains reproduce the anatomical segmentation; it is well above chance but far from perfect agreement. The disagreement is also structured rather than random, in that our domains subdivide several organs, most visibly the heart, into multiple territories. We interpret this as a consequence of partitioning the tissue on a different quantity: the published annotation partitions by overall expression profile, labeled by cell-type markers, whereas these domains partition by inferred ligand-receptor signaling context, and there is no reason for the two to coincide. The conclusion we draw is the narrower one, that the emergent domains recover major anatomical structures while resolving additional substructure within them.

The two controls answer different questions, and we do not want them read as interchangeable. The coordinate shuffle asks whether the structure is real, and it is the one that can rule out the smoothing and dimensionality explanations, because it manipulates the data and refits everything. The concordance comparison asks whether the structure that survives is anatomically meaningful, which is what makes it the natural complement rather than a substitute.

Reproducibility of the clustering

Two implementations of the clustering step appear in this work, and their reproducibility has to be stated separately. Figs 5 and 6 use Seurat. The coordinate-shuffle control, and one arm of the pair-shuffling control below, use a faster implementation of the same two steps, a nearest-neighbor graph built with RcppHNSW followed by Louvain clustering in igraph, which made it affordable to cluster many permuted datasets. The two implementations give ELSA values on different scales, so each is compared only with itself.

The faster implementation, as we first configured it, did not reproduce exactly from run to run. Its nearest-neighbor graph was built across four threads without a fixed random seed, and under concurrent insertion the graph depends on the order in which threads happen to complete. Across eight replicate runs on identical input, ELSA at radius 50 ranged from 0.053 to 0.067, with a standard deviation of 0.0045, and the number of domains ranged from 19 to 22. Two conditions together make it

exactly reproducible, and neither suffices alone. With four threads and a fixed seed, three replicate runs still gave 19, 19 and 20 domains; unseeded single-threaded runs gave 19, 22 and 22. With a single thread *and* a fixed seed, five replicate runs returned bit-identical cluster assignments, with 21 domains in all five, and a second seed likewise gave three identical runs, at 23 domains. The seed therefore changes the result, and is a parameter of the analysis to be reported rather than an incidental detail. The coordinate-shuffle control was run before this was established; its effect is far larger than this run-to-run spread, so its conclusion does not depend on it.

For the Seurat clustering behind Figs 5 and 6 we have not established whether the result is exactly reproducible, and we have not measured its run-to-run variation directly. As an indirect check, an independently written script re-ran the variance-ordered sweep at six depths (10, 50, 150, 350, 650 and 1000 fields) and the full-depth clustering. Mean ELSA at radius 50 was within 0.012 of the published value at each of the six depths, and at full depth it was 0.1236 against the published 0.1237, with 19 domains against the published 20. These re-runs omitted the four duplicated input columns described in the Materials and methods section, so they differ from the published runs in input as well as in run. We therefore treat each domain map in Figs 5 and 6 as a single realization, read its domain count as approximate, and regard differences in ELSA of about 0.01 between single Seurat runs as within what a re-run could produce.

Shuffling the ligand-receptor pairing does not make the domains less coherent

A third control asks whether the domains depend on which receptor is paired with which ligand. We reassigned receptors to ligands at random, then repaired by targeted swaps every reassigned pair that coincided with a real FANTOM5 pairing or paired a gene with itself, so that no shuffled set retained a real pair. The set of receptors is unchanged, so every shuffled set is built from exactly the same fitted fields as the true pairing, and only the pairing differs. Each shuffled set then passed through the same convolution, clustering and scoring as the true pairing.

In Seurat, with 19 shuffled sets clustered at full depth, the true pairing gave an ELSA at radius 50 of 0.124 against a shuffled mean of 0.093 (standard deviation 0.018), so 17 of the 19 shuffles produced more coherent domains than the true pairing did; the true pairing gave 19 domains against a shuffled mean of 18.5. In the faster implementation, run with a single thread and a fixed seed on 99 shuffled sets, the true pairing sat inside the shuffled distribution: 24 of the 99 shuffles were more coherent, and it gave 21 domains against a shuffled mean of 21.05. The direction of the small difference is not consistent between the two implementations, and we read the result as no detectable effect of the pairing on the coherence of the domains.

The pairing is not without consequence, however. True pairings yield 1,449 numerically distinct fields. Because this count needs no clustering, we computed it for 9,999 shuffled pairings, made with the same procedure and including the 99 clustered ones: they gave a mean of 1,424 (standard deviation 5.9), and only one reached 1,449 or more, a permutation p of 0.0002. True partners are also more alike in spatial pattern than random partners. For each of the 1,477 pairs we compared the correlation between the two true partner fields with the correlation between the ligand field and that of a different receptor that is not a FANTOM5 partner of the ligand, chosen from 30 random candidates to match the true pair as closely as possible in how consistently one partner is the smaller of the two. The mean Pearson correlation was 0.111 for true partners and 0.068 for random ones, true partners were the more correlated in 60.8% of pairs, and a paired t-test gave $p = 4 \times 10^{-21}$. Even random partners are positively correlated (mean Spearman correlation 0.325), consistent with

a spatial pattern shared by most of these genes, so the part of the correlation specific to the pairing is modest. We return to what these results imply in the Discussion.

The interaction fields are plug-in estimates, and differ from their posterior summaries

Each published interaction field is a plug-in estimate: at every bin it is the smaller of the two partners' posterior medians, $\min(\tilde{L}(\boldsymbol{s}), \tilde{R}(\boldsymbol{s}))$, where $\tilde{L}(\boldsymbol{s})$ and $\tilde{R}(\boldsymbol{s})$ denote the posterior medians of the ligand's and the receptor's expected count at bin $\boldsymbol{s}$. The quantity of interest is the interaction itself, $M(\boldsymbol{s}) = \min(L(\boldsymbol{s}), R(\boldsymbol{s}))$, whose posterior is not summarized by the minimum of two medians. Because each feature is fitted separately, the ligand and receptor posteriors are independent, so at each bin $P(M(\boldsymbol{s}) > t) = P(L(\boldsymbol{s}) > t)\, P(R(\boldsymbol{s}) > t)$, and the posterior of $M(\boldsymbol{s})$ follows from the two marginal posteriors alone. Posterior draws were not stored, so we approximated each marginal posterior by a lognormal distribution matched to its stored median and 95% interval, which corresponds to a Gaussian posterior on the scale of the linear predictor. The approximation is adequate for most bins: on the log scale the stored intervals are close to symmetric (median ratio of upper to lower half-width 0.995, with 90% of bins between 0.85 and 1.12, in a random sample of 60 features), and the implied posterior means agree with the stored ones to within 10% for 90% of feature-bin combinations.

Relative to the posterior median of $M(\boldsymbol{s})$, the plug-in field is never lower. This holds exactly: with $m^* = \min(\tilde{L}, \tilde{R})$, $P(M > m^*) = P(L > m^*)\, P(R > m^*) \leq 1/2$. Across all 1,477 pairs and 26,000 bins the overstatement has a median of 1.3%, but it is heavy-tailed: it exceeds 10% in 35% of pair-bin combinations and 25% in 22% (S11 Fig). It is concentrated where neither partner is clearly the limiting one, that is, where the two marginal posteriors overlap; the per-pair Spearman correlation between the overstatement and the standardized gap between the partners' log-scale posterior medians has a median of -0.98. Relative to the posterior mean of $M(\boldsymbol{s})$ the direction reverses: the marginal posteriors are wide and right-skewed (median log-scale standard deviation 1.03 in the same sample), so their means lie well above their medians, and the plug-in field is below the posterior mean of $M(\boldsymbol{s})$ in a median of 91% of bins per pair.

For the analyses reported here, the more relevant question is whether these differences change spatial pattern, since each field is scaled before clustering. They change it little. Per pair, the Pearson correlation over bins between the plug-in field and the posterior median of $M(\boldsymbol{s})$ has a median of 0.988 (5th percentile 0.963), and between the plug-in field and the posterior mean of $M(\boldsymbol{s})$ a median of 0.967 (5th percentile 0.905). We therefore expect the domains to change little if the plug-in fields were replaced by either posterior summary, but we have not re-run the clustering to confirm this.

Discussion

A central question in tissue biology is how and why tissues develop, retain, and lose their form. This broad question is fundamental to developmental biology, tissue engineering, regenerative medicine, and multicellular biology across species. For the better part of a century, we have understood that extracellular signaling plays a critical role in the patterning and organization of cells within tissues. Local intercellular cues preserve adult stem cell niches, govern cellular differentiation, and regulate tissue response to injury. However, despite this knowledge, our ability to *understand* cellular

communication in tissues, i.e., to deliberately influence it to induce healing or desired change, is woefully limited.

Recent research has revealed that the grammar of cellular communication is intrinsically high-dimensional, with cells communicating on hundreds-to-thousands of wavelengths at once and processing/interpreting these cues via whole-cell parallel processing [7]. This means that in order to properly model and visualize cell-to-cell communication, we must be comfortable observing and interpreting high-dimensional signaling 'hyper-fields' as single, holistic objects of study in histologic space. The work in this paper has been carried out with this principle in mind.

We see the outputs from these models as the start of a growing foundation for what we envision as 'morphogenic field analysis': the interdisciplinary study of how morphogenic field interactions, gradients, dynamics, and perturbations shape tissue organization and control cellular differentiation, ideally in three-dimensions over time. Although the data analysis shown here is exclusively two-dimensional, and does not include a time dimension, there are many datasets emerging which do allow spatiotemporal analysis and there are incoming tools allowing affordable 3D profiling [33]. ~~Our approach is easily adapted to 3-dimensional timeseries data. When applied~~ The geostatistical framework we use extends in principle to three spatial dimensions and to a time dimension, since the same Gaussian-process machinery is routinely applied in those settings in other fields, though we have not done so here. Applied to timeseries datasets, longitudinal ~~quantitative~~ modeling could be ~~performed to show how~~ used to ask whether tissue field dynamics precede, align with, or follow cellular growth and change during tissue remodeling. ~~Our methods are generally-applicable to any spatial transcriptomic data when the right choices are made:~~

We have demonstrated this approach on a single field of view, from one section of one embryo, on one platform, at one binning resolution, and we do not want to claim more reach than that supports. It seems more useful to state what we think would need to hold for the approach to transfer. The observations must be counts accompanied by an interpretable exposure or offset, since the model treats the local total as known sampling effort. The spatial support must be dense enough that a Gaussian process is identifiable at the length scales of interest, which is a joint property of a platform's resolution, its capture efficiency, and the binning applied. Platforms with substantially sparser capture than the data used here may favor a zero-inflated likelihood, which our comparison found unnecessary at this sparsity but which the framework accommodates. At single-cell-resolution and imaging-based platforms, where segmentation rather than binning defines the observation unit, the binning step we rely on would need to be replaced by an explicit treatment of cell geometry; spot-based ~~vs.~~ and imaging-based ~~transcriptomics, for instance, should be treated differently, as~~ data should not be treated identically, because their measurement methods and organization ~~are fundamentally different. Further, our model is easily customized for specific tissue types~~differ. Tissue-specific customization is likewise possible but is not automatic: we are ~~currently~~ separately experimenting with an ~~approach optimized~~ adaptation for lung tissue that ~~allows exclusion of~~ excludes large lumens and alveolar airspaces from the tissue fields.

Two features of the framework are worth separating from the demonstration itself, because they are not specific to the platform used here. The model requires only counts at known locations together with a measure of local sampling effort, and the stochastic partial differential equation representation of the Gaussian process is defined on a triangulated mesh rather than on a lattice, so an irregular, hexagonal, or otherwise non-gridded arrangement of observation locations requires no special handling. We have separately fitted spatial fields of this kind to 10x Visium mouse

brain sections, whose hexagonal spot array and much coarser spatial resolution differ substantially from the binned nanoball array used here, obtaining fields for 2,035 ligand-receptor mechanisms across two sagittal sections, with the posterior predictive occurrence probabilities tracking the observed occurrence closely across all 3,865 fits (correlation 0.99, median absolute difference 0.015). We report this as an indication that the field-estimation step is not tied to one platform's geometry, and deliberately not as a validation of the pipeline as a whole: that work used a two-part semicontinuous observation model on normalized data rather than the raw-count model developed here, and it did not carry the fitted fields through the convolution and clustering stages. Comprehensive evaluation across tissues and platforms remains future work.

Applying the framework to diseased tissue is a natural next step, and one motivation for developing it, but it is future work rather than something this manuscript demonstrates. This work is a methodological proof of concept on healthy developing tissue. Doing the diseased case properly would require matched healthy and diseased tissue, attention to the confounding between disease state and tissue composition, and enough replication to separate a disease effect from between-sample variation. Carried out on a single additional section, it would yield an illustrative figure that we could not defend statistically, which is the opposite of what we are arguing for here.

Relationship to existing methods

Several established methods place a Gaussian process over space in transcriptomic data, and it is worth stating precisely which parts of the present approach are new and which are combinations or reinterpretations of existing ideas.

SpatialDE [34] introduced Gaussian-process modeling for spatial transcriptomics, decomposing each gene's expression variance into a spatial component parameterized by a covariance kernel and a non-spatial noise component, then testing the spatial component by a likelihood ratio test. We share its use of a Gaussian process prior over space, but not its inferential target: SpatialDE asks which genes are spatially variable, whereas we estimate the field itself and then use it downstream.

BOOST-GP [35] is the closest methodological relative. It places a gene-specific zero-mean Gaussian process on the spatial random effect inside a zero-inflated negative binomial count model, with Bayesian variable selection over whether that spatial component is present, and its stated goal is likewise the identification of spatially variable genes. The combination of a Gaussian process with a Bayesian count likelihood is therefore not something we claim as novel. What differs is what is done with the posterior: we retain the fitted fields as continuous objects with quantified uncertainty, convolve them pairwise, and analyze the resulting high-dimensional collection jointly.

Spatial Variance Component Analysis (SVCA) [36] fits an additive Gaussian-process random-effect model that decomposes the variance of each gene or protein into intrinsic, environmental, cell-cell interaction, and noise components, and tests whether the interaction component is nonzero. Its aim is variance partitioning rather than field estimation, and it was developed for imaging-based data.

SpatialDM [37] is the closest in biological target. It tests ligand-receptor pairs for spatial co-expression using a bivariate extension of Moran's I, both globally, selecting which pairs are spatially co-expressed across the tissue, and locally, identifying the spots that drive the association, with significance from an analytically derived null distribution. It targets the significance of individual interactions, whereas we target a continuous inferred field per interaction, and then cluster across all of them at once.

CytoSignal [38] appeared after our original submission and analyzes the same Stereo-seq mouse embryo atlas. It computes a per-cell ligand-receptor score as the product of spatially aggregated ligand and receptor expression, using a Gaussian kernel for diffusion-dependent interactions and Delaunay neighbors for contact-dependent ones, assesses significance against a spatial-permutation null, and validates predicted interactions against proximity ligation assays performed on adjacent tissue sections. It is the one method in this list that, like ours, produces a spatially resolved signaling quantity rather than a per-feature test statistic. In light of that work we do not claim novelty for spatially resolved ligand-receptor mapping. Our operator comparison (Fig 5) is relevant here, in that the common minimum yielded more spatially coherent domains on this dataset than the product operator that CytoSignal uses, but that is a comparison of two convolution operators inside our own pipeline, not a comparison of the two methods.

Stating our contribution precisely, and more narrowly than our original submission did, it consists of a generative Bayesian geostatistical treatment of raw spatialomics counts, which yields a posterior distribution rather than a point estimate for every feature field; the argument for modeling raw counts as a proxy for absolute binding density rather than modeling normalized expression; a quantitative comparison of convolution operators; and the finding that clustering the full collection of inferred interaction fields produces tissue domains that correspond to major anatomical structures while resolving substructure within them. As the pair-shuffling control shows, those domains do not depend on the specific ligand-receptor pairings, which bounds what this last finding says about signaling.

The pair-shuffling control (see the Controls: the domains survive a spatial null and match independent anatomy section) limits what the domains can be said to show. Shuffling which receptor is paired with which ligand did not make the domains less coherent, even though true partners are measurably more alike in spatial pattern than random partners, and true pairings yield more distinct fields than shuffled ones. The domains reported here are therefore best read as a property of the collection of fitted ligand and receptor fields as combined by the common minimum on raw counts, and not as evidence that the specific ligand-receptor pairings shape them. We suspect the operator is largely responsible, though we have not tested this directly. On raw counts one partner of a pair is often the smaller of the two almost everywhere (for 42.6% of pairs, at more than 95% of locations), so the common minimum frequently returns that partner's field with only limited regions carved out of it, and little of what distinguishes a true partner from a random one need reach the clustering. Whether a different operator, or the same operator applied to fields placed on a common scale, would carry the pairing through is a question for future work.

We should be explicit about one limit of the present analyses. The convolution and everything downstream of it operate on posterior medians of the feature fields, not on posterior draws, so the interaction fields presented here are plug-in point estimates and the domain results do not carry propagated uncertainty. The first step toward propagating it is described in the The interaction fields are plug-in estimates, and differ from their posterior summaries section: the posterior of each interaction field can be computed from the stored marginal posteriors without refitting, and it differs from the plug-in field in level, by amounts that are largest where the two partners are comparable, much more than in spatial pattern. Carrying that posterior through the clustering itself, so that the domains come with uncertainty, is a computation we have not done, and we do not want the availability of the posteriors to be read as a claim that we have.

We have not run a head-to-head benchmark against any of these methods, and we would rather be direct about that than leave it implicit. They answer different

questions: SpatialDE and BOOST-GP test genes for spatial variability, SpatialDM tests ligand-receptor pairs for spatial association, and SVCA partitions variance into components. There is no shared target quantity on which an accuracy comparison would be meaningful without first defining a benchmark task and a ground truth, which we regard as a separate piece of work. A reader who wants a significance test for individual features or individual interactions should use one of the methods above; the framework presented here is aimed at estimating the fields and analyzing them jointly.

The statistical models presented here have some limitations. First, they must be tuned to each input type of data: Xenium, Visium, ~~StereoSeq~~ Stereo-seq (shown here), CurioTrekker, CosMx, COMET, ~~etc. However, our general approach has thus far proved easy to adapt on all data types tested~~and others. We have fitted the model only to Stereo-seq data in this work, so our expectation that adapting it to other platforms is straightforward remains an expectation rather than a demonstrated result. Second, we currently are ignoring three major considerations: 1. each signaling mechanism is treated the same (there is no incorporation of molecule-specific diffusion coefficients based on molecular weight), 2. there is no accommodation for variable extracellular matrix properties (competitive binding, protein-density-diffusion limitations), and 3. there is no accommodation for ligand-receptor sequestration due to binding (complex non-linear modeling required). These deficits shall be addressed in future studies, and present many interesting opportunities for further experimentation.

"All models are wrong, but some are useful". The modeling outputs shown here are a ~~statistically-powerful~~ *proxy* for high-dimensional information flow *potential* within tissues. Stated plainly, the interaction fields estimate the potential for signaling under the co-expression assumption standard in cell-cell communication analysis, that a ligand and its receptor expressed in the same place have the opportunity to interact. We propose them as a proxy for signaling; we do not observe signaling. It is worth setting out the inferential chain explicitly, so a reader can see how far these outputs sit from the measurement. Transcript counts are a proxy for local transcript abundance, subject to the sampling variability this framework is designed to model. Transcript abundance is in turn a proxy for protein abundance, a step which is known to be imperfect and which we do not attempt to correct. Co-localization of a ligand proxy and a receptor proxy within the same bin of 50 by 50 capture spots, rather than at a shared cell surface, is a proxy for the *opportunity* for those molecules to interact, and the convolution operators we compare are alternative ways of quantifying that opportunity under stated stoichiometric assumptions. None of the intermediate quantities, and no binding event, is directly observed in this work. Further data must be acquired at the protein level and the intracellular level before any of these model outputs are treated as evidence of actual information transmission or transduction. The experimental cost of doing this, however, can be high, ~~particular~~ particularly if the patterns of interest are high-dimensional and cannot be reduced to targeted cell-scale, pathway-scale, or molecule-scale experiments. Therefore, in our opinion, the ~~primary power~~ value of the modeling approach described in this paper lies not in its ability to predict single-mechanism interactions. Rather, it lies in its ~~ability to facilitate statistically-robust mappingsbetween~~ potential to support mappings, with quantified uncertainty, between estimates of the holistic high-dimensional extracellular signaling ~~milieus~~ milieu and cell-state-specific intracellular state changes and transitions. A cell within a tissue experiences hundreds to thousands of cues at once, and processes these cues in parallel—, as a single, holistic, high-dimensional signal. Our modeling approach allows us to locally estimate a transcript-derived proxy for this high-dimensional signal while providing measures of statistical ~~confidence~~uncertainty. When paired with *intra*cellular data, such as that regarding transcription factor activation, cytoskeletal remodeling, metabolic reprogramming and mechanoresponse, we are newly provided

with an opportunity to quantitatively study the precise ways that different cells, in different states, process and respond to high-dimensional dynamics in their local extracellular milieu: ~~i.e., how~~ that is, how the holistic cellular microenvironment governs cellular state~~—~~, and therefore how extracellular milieus might be targeted in the future to induce regenerative healing.

Our code is open access on GitHub, and we have included all of our work from this project so that others can ~~replicate our results~~rerun our analyses, see how we have organized our inputs ~~/~~and outputs, and experiment on their own. We note that rerunning the clustering stage may not reproduce the published domain map exactly (see the Controls: the domains survive a spatial null and match independent anatomy section); the fitted fields and the stored domain labels are provided so that downstream work does not depend on reproducing it. The images presented in the figures of this paper are a small fraction of our model outputs~~—~~, and we encourage interested readers to download the ~~supplemental materials~~ supporting information included with this paper as a way to ~~observe the true~~ explore the scale and structure of ~~extracellular communication in mammalian tissues~~the inferred signaling fields. We look forward to working with the community to explore morphogenic field analysis in additional tissue datasets and to brainstorm ways to influence spatial extracellular signaling for therapeutic means.

Code availability

All code used to create this work is available on GitHub at `https://github.com/aaron-oz/a-s-omics`.

Data availability

Raw data used to create the figures in this manuscript can be downloaded from [28] at `https://db.cngb.org/data_resources/project/CNP0001543`.

Supporting information

We have provided a set of documents on FigShare which capture key intermediate- and end-outputs of the modeling described in this paper. They can be downloaded from the following links:

S1 File. Raw-to-bin. `https://figshare.com/s/e921e8e36296cd8c57da`

S2 File. Feature refining. `https://figshare.com/s/6656c05d4552a59afa05`

S3 File. Raw vs. norm, selected features.
`https://figshare.com/s/4885d58ba45b2f2e7689`

S4 File. Model comparison, all features.
`https://figshare.com/s/9d266d5f0fd39632986d`

S5 File. HD embeddings. `https://figshare.com/s/7bbe3feb729a9731acfa`

S6 File. ELSA analysis. `https://figshare.com/s/4867799e0e8b68fba5a8`

S7 File. Clusters vs. features.
`https://figshare.com/s/2e305ce9ee20efd7ee47`

S8 File. Morphogen fields. `https://figshare.com/s/2f66def6d22f1549b312`

S9 Table. Notation-to-code mapping. A row for each mathematical object in the model specification, giving its meaning, the script in which it is constructed, and the name of the corresponding object in the analysis code.

S10 Fig. Controls. (A) Spatial permutation test. Distribution of mean ELSA at radius 50 across the 49 coordinate-permuted replicates, each one a full refit of all 912 features, with the observed value marked in red. Lower ELSA means more spatially coherent domains, so the observed data sit far below the null. **(B)** The same permutation test for the number of domains recovered, meaning the count of distinct cluster labels returned: 20 in the observed data against 4 to 12 across the permutations. **(C)** The emergent signaling domains. **(D)** The anatomical annotation published for the same tissue section. In both panels each anatomical label carries one hue, and every emergent domain in (C) is drawn in the hue of the label it best matches, with luminance varied where several domains map to the same label; corresponding regions therefore share a hue between the two panels, while domains that subdivide a single organ remain distinguishable. **(E)** Agreement domain by domain: the purity of each emergent domain against its dominant anatomical label, ordered by purity and colored as in (C), with the size-weighted mean marked. Overall agreement is an adjusted Rand index of 0.401 and normalized mutual information of 0.604, against a null from 499 random permutations of the domain labels in which every value fell below 0.0011. Generated by `figures-regenerated/make-controls-figure.R` in the code repository.

S11 Fig. Plug-in interaction fields against their posteriors. (A) For each bin, the 90th percentile across the 1,477 ligand-receptor pairs of the relative overstatement of the plug-in field, $\min(\tilde{L}, \tilde{R})$, over the posterior median of $M = \min(L, R)$. **(B)** The same overstatement against the standardized gap between the two partners, $|\log \tilde{L} - \log \tilde{R}| / \sqrt{\sigma_L^2 + \sigma_R^2}$, where σ_L and σ_R are the log-scale posterior standard deviations, for 500 random bins from each of 200 random pairs; shading counts bins on a log scale. The overstatement is large only where the gap is small, meaning neither partner is clearly limiting. **(C)** Per pair, the Pearson correlation over bins between the plug-in field and the posterior median (bottom) or posterior mean (top) of M. Posteriors of M are computed from lognormal approximations to the stored marginal posteriors; see the The interaction fields are plug-in estimates, and differ from their posterior summaries section.

Acknowledgments

The authors thank the members of the Raredon laboratory for helpful discussion. We also thank the reviewers, whose request for controls prompted the analyses reported in the Controls: the domains survive a spatial null and match independent anatomy section.

References

1. Vainio S, Müller U. Inductive Tissue Interactions, Cell Signaling, and the Control of Kidney Organogenesis [Journal Article]. Cell. 1997;90(6):975-8. Doi: 10.1016/S0092-8674(00)80363-3. Available from: https://doi.org/10.1016/S0092-8674(00)80363-3. doi:10.1016/S0092-8674(00)80363-3.

2. Basson MA. Signaling in cell differentiation and morphogenesis [Journal Article]. Cold Spring Harbor perspectives in biology. 2012;4(6):a008151.

3. Hagolani PF, Zimm R, Marin-Riera M, Salazar-Ciudad I. Cell signaling stabilizes morphogenesis against noise [Journal Article]. Development. 2019;146(20):dev179309.

4. Wu D, Yamada KM, Wang S. Tissue Morphogenesis Through Dynamic Cell and Matrix Interactions [Journal Article]. Annual Review of Cell and Developmental Biology. 2023;39(Volume 39, 2023):123-44. Available from: https://www.annualreviews.org/content/journals/10.1146/annurev-cellbio-020223-031019. doi:https://doi.org/10.1146/annurev-cellbio-020223-031019.

5. Li P, Elowitz MB. Communication codes in developmental signaling pathways [Journal Article]. Development. 2019;146(12). Available from: https://doi.org/10.1242/dev.170977. doi:10.1242/dev.170977.

6. Klumpe HE, Langley MA, Linton JM, Su CJ, Antebi YE, Elowitz MB. The context-dependent, combinatorial logic of BMP signaling [Journal Article]. Cell Systems. 2022;13(5):388-407.e10. Doi: 10.1016/j.cels.2022.03.002. Available from: https://doi.org/10.1016/j.cels.2022.03.002. doi:10.1016/j.cels.2022.03.002.

7. Granados AA, Kanrar N, Elowitz MB. Combinatorial expression motifs in signaling pathways. Cell Genomics. 2024;4(1):100463. doi:https://doi.org/10.1016/j.xgen.2023.100463.

8. Brennan MD, Cheong R, Levchenko A. How Information Theory Handles Cell Signaling and Uncertainty [Journal Article]. Science. 2012;338(6105):334-5. Available from: https://www.science.org/doi/abs/10.1126/science.1227946. doi:doi:10.1126/science.1227946.

9. Ellison D, Mugler A, Brennan MD, Lee SH, Huebner RJ, Shamir ER, et al. Cell–cell communication enhances the capacity of cell ensembles to sense shallow gradients during morphogenesis [Journal Article]. Proceedings of the National Academy of Sciences. 2016;113(6):E679-88.

10. Scuderi S, Kang TY, Jourdon A, Nelson A, Yang L, Wu F, et al. Specification of human brain regions with orthogonal gradients of WNT and SHH in organoids reveals patterning variations across cell lines [Journal Article]. Cell Stem Cell. 2025;32(6):970-89.e11. Doi: 10.1016/j.stem.2025.04.006. Available from: https://doi.org/10.1016/j.stem.2025.04.006. doi:10.1016/j.stem.2025.04.006.

11. Weaver VM, Roskelley CD. Extracellular matrix: the central regulator of cell and tissue homeostasis [Journal Article]. Trends in cell biology. 1997;7(1):40-2.

12. Hironaka Ki, Morishita Y. Encoding and decoding of positional information in morphogen-dependent patterning [Journal Article]. Current Opinion in Genetics & Development. 2012;22(6):553-61. Available from: https://www.sciencedirect.com/science/article/pii/S0959437X12001268. doi:https://doi.org/10.1016/j.gde.2012.10.002.

13. Meizlish ML, Franklin RA, Zhou X, Medzhitov R. Tissue Homeostasis and Inflammation [Journal Article]. Annual Review of Immunology. 2021;39(Volume 39, 2021):557-81. Available from: https://www.annualreviews.org/content/journals/10.1146/annurev-immunol-061020-053734. doi:https://doi.org/10.1146/annurev-immunol-061020-053734.

14. Kicheva A, Briscoe J. Control of tissue development by morphogens [Journal Article]. Annual review of cell and developmental biology. 2023;39(1):91-121.

15. Yue L, Liu F, Hu J, Yang P, Wang Y, Dong J, et al. A guidebook of spatial transcriptomic technologies, data resources and analysis approaches [Journal Article]. Computational and Structural Biotechnology Journal. 2023;21:940-55. Available from: https://www.sciencedirect.com/science/article/pii/S2001037023000156. doi:https://doi.org/10.1016/j.csbj.2023.01.016.

16. Wang Y, Liu B, Zhao G, Lee Y, Buzdin A, Mu X, et al. Spatial transcriptomics: Technologies, applications and experimental considerations [Journal Article]. Genomics. 2023;115(5):110671. Available from: https://www.sciencedirect.com/science/article/pii/S0888754323001155. doi:https://doi.org/10.1016/j.ygeno.2023.110671.

17. Li H, Chen Z, Zhu W. Variability: Human nature and its impact on measurement and statistical analysis. Journal of Sport and Health Science. 2019;8(6):527.

18. Cannon MJ, Warner L, Taddei JA, Kleinbaum DG. What can go wrong when you assume that correlated data are independent: an illustration from the evaluation of a childhood health intervention in Brazil. Statistics in medicine. 2001;20(9-10):1461-7.

19. Bressan D, Battistoni G, Hannon GJ. The dawn of spatial omics. Science. 2023;381(6657):eabq4964.

20. Raredon MSB, Yang J, Kothapalli N, Lewis W, Kaminski N, Niklason LE, et al. Comprehensive visualization of cell–cell interactions in single-cell and spatial transcriptomics with NICHES [Journal Article]. Bioinformatics. 2022;39(1). Available from: https://doi.org/10.1093/bioinformatics/btac775https://watermark.silverchair.com/btac775.pdf?token=AQECAHi208BE49Ooan9kkhW_Ercy7Dm3ZL_9Cf3qfKAc485ysgAAAwUwggMBBgkqhkiG9w0BBwagggLyMIIC7gIBADCCAucGCSqGSIb3DQEHAawdFVcEYJT6yulGUsQPAOlWXV3xO57vy9WO3k624kE6_T22tAdIP3MXbFD2Q4AwpwkRp_1s5QzN1pSUxRTqEwmvEsTSr_OQe8EJe_xYkpXoBjUEEXISO5D5sV8rDy3Pa8gzAlvzytYftzhvzISZOAqW9GLyGVJS-IXkLk38OIV_w-w_s4h2N8-uchq08ANp7v9pXwuJ4I4F_hAsa8qLveqwvNR9T0UZvkASTYunLGw0PjT--ZOvEAUeaAVEt3NW9p9OglMP-ovi9qUQS9TDOHC50LQ_0I_Y7d-uAvpRv_uMV6JnjaPiODu6Gv5Lto2okvlFPsGCVi0zQgIzGsFRL6Vwmkh3MeuQ_T3ISi_Ca-7jc-mNGPtLt44Y7q6WMTc_W7z4GqquvMorlXcf-mMmm8gXAxr6EqnQJb5loKmv2uy0wPdwNXz7XPDRc94WwzvsHE-AfbAUkL

qcBtx_ETELJCU5zKjUQnbR_4ECf9cbn_hJCCB7KrCRFm8ncfw25ZwmPOUKvC_
KSZbre_
E9cRbx7SYpviuNzbkBAc3dgmUWSzcs78MryvxcYhfeRdvf-iRdf7vsn2adVcrd28SVxaYOrqhT
3pdZQs5TcyVP5BMwjM3SAQN9ge1H1M-kPl6y9TkSerPH5EYit0RImx-wExtAS8mK_
TvCsKH_SQK7DLLMXBeMbXrS2syZfHYZvWFyx1Pv5g6ADs9ym1aMw1sbRuWhoihIw.
doi:10.1093/bioinformatics/btac775.

21. Wilk AJ, Shalek AK, Holmes S, Blish CA. Comparative analysis of cell–cell communication at single-cell resolution [Journal Article]. Nature Biotechnology. 2024;42(3):470-83. Available from: https://doi.org/10.1038/s41587-023-01782-z. doi:10.1038/s41587-023-01782-z.

22. Valcu M, Kempenaers B. Spatial autocorrelation: an overlooked concept in behavioral ecology. Behavioral Ecology. 2010;21(5):902-5.

23. Cressie N, Wikle CK. Statistics for spatio-temporal data. John Wiley & Sons; 2011.

24. Fuglstad GA, Simpson D, Lindgren F, Rue H. Constructing priors that penalize the complexity of Gaussian random fields. Journal of the American Statistical Association. 2019;114(525):445-52.

25. Lindgren F, Rue H, Lindström J. An explicit link between Gaussian fields and Gaussian Markov random fields: the stochastic partial differential equation approach. Journal of the Royal Statistical Society Series B: Statistical Methodology. 2011;73(4):423-98. doi:10.1111/j.1467-9868.2011.00777.x.

26. Rue H, Martino S, Chopin N. Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations. Journal of the Royal Statistical Society Series B: Statistical Methodology. 2009;71(2):319-92. doi:10.1111/j.1467-9868.2008.00700.x.

27. Naimi B, Hamm NA, Groen TA, Skidmore AK, Toxopeus AG, Alibakhshi S. ELSA: Entropy-based local indicator of spatial association. Spatial statistics. 2019;29:66-88.

28. Chen A, Liao S, Cheng M, Ma K, Wu L, Lai Y, et al. Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA nanoball-patterned arrays [Journal Article]. Cell. 2022;185(10):1777-92. e21.

29. Ramilowski JA, Goldberg T, Harshbarger J, Kloppmann E, Lizio M, Satagopam VP, et al. A draft network of ligand–receptor-mediated multicellular signalling in human [Journal Article]. Nature communications. 2015;6:7866.

30. Gneiting T, Raftery AE. Strictly proper scoring rules, prediction, and estimation. Journal of the American statistical Association. 2007;102(477):359-78.

31. Bhuva DD, Tan CW, Salim A, Marceaux C, Pickering MA, Chen J, et al. Library size confounds biology in spatial transcriptomics data [Journal Article]. Genome Biology. 2024;25(1):99. Available from: https://doi.org/10.1186/s13059-024-03241-7. doi:10.1186/s13059-024-03241-7.

32. Naimi B, Hamm NA, Groen TA, Skidmore AK, Toxopeus AG, Alibakhshi S. ELSA: Entropy-based local indicator of spatial association [Journal Article]. Spatial statistics. 2019;29:66-88.

33. Reynolds DE, Roh YH, Oh D, Vallapureddy P, Fan R, Ko J. Temporal and spatial omics technologies for 4D profiling [Journal Article]. Nature Methods. 2025;22(7):1408-19. Available from: https://doi.org/10.1038/s41592-025-02683-6. doi:10.1038/s41592-025-02683-6.

34. Svensson V, Teichmann SA, Stegle O. SpatialDE: identification of spatially variable genes. Nature Methods. 2018;15(5):343-6. doi:10.1038/nmeth.4636.

35. Li Q, Zhang M, Xie Y, Xiao G. Bayesian modeling of spatial molecular profiling data via Gaussian process. Bioinformatics. 2021;37(22):4129-36. doi:10.1093/bioinformatics/btab455.

36. Arnol D, Schapiro D, Bodenmiller B, Saez-Rodriguez J, Stegle O. Modeling cell-cell interactions from spatial molecular data with spatial variance component analysis. Cell Reports. 2019;29(1):202-11.e6. doi:10.1016/j.celrep.2019.08.077.

37. Li Z, Wang T, Liu P, Huang Y. SpatialDM for rapid identification of spatially co-expressed ligand-receptor and revealing cell-cell communication patterns. Nature Communications. 2023;14:3995. doi:10.1038/s41467-023-39608-w.

38. Liu J, Manabe H, Qian W, Orikasa S, Ghaemmaghami J, Wang Y, et al. CytoSignal detects locations and dynamics of ligand-receptor signaling at cellular resolution from spatial transcriptomic data. Nature Genetics. 2026;58(6):1396-408. doi:10.1038/s41588-026-02624-9.