
An Agent-based Model of Voluntary Incentive Programs for Sustainable Biofuel Development



An Agent-based Model of Voluntary Incentive Programs for Sustainable Biofuel Development

Robert Zupko¹, Himamshu Ghimre¹, Peter Nouhan¹, and Mark Rouleau¹
Michigan Technological University, Houghton, Michigan, USA

Keywords: agent-based modeling, biofuels, geographic information systems, non-industrial private forest owners, policy

1 Research Purpose

Future demand for energy is expected to increase well beyond growth in proven fossil fuel reserves. Reliance on diminishing reserves will also amplify climate change. Renewable energy makes it possible for society to overcome these two interrelated problems. Second-generation bioenergy from woody biomass offers a promising renewable alternative for regions with ready access to forests, especially those incapable of fully harnessing wind, hydro, or solar technologies. However, the success of bioenergy greatly depends on the proper management of biomass harvesting activity to ensure bioenergy remains a sustainable renewable alternative.

Our research uses ForestSim, an agent-based model (ABM) of non-industrial private forest (NIPF) owners, to evaluate voluntary incentive programs (VIPs) and their ability to produce sustainable bioenergy outcomes. Under a VIP landowners receive a financial incentive for compliance with program guidelines. We compare traditional financial incentivization mechanisms to a newly emerging VIP scheme known as an agglomeration bonus [10]. These programs provide additional incentives based upon participation rate as an incentive for neighbors to recruit others. Model results show that agglomeration bonuses allow for better centralized management of forest resources and produce more sustainable outcomes.

2 Background

Previous research into NIPF owners has started with work in economics in an attempt to determine willingness-to-harvest with regards to woody biomass supply and the price points involved [1, 4]. Recently there has been growing interest in the use of ABMs in the context of CHANS [2]. With regards to bioenergy, ABMs have been used to evaluate factors affecting crop adoption and decision practices [7, 11, 13]. These ABMs exist within a broader research space that focuses on agriculture [5, 12], generalized land-use change [3, 19, 15], and policy analysis [20]. The field of forestry is starting also starting to see the adoption of ABMs being used. Previous work has focused on forest management [6] along

with deforestation and reforestation cycles [18]. Finally, an ABM developed to model harvest decisions by small scale forest owners [16]. Our work is similar to the work of Leahy et al., but expands upon it by evaluating biomass production in the context of sustainability.

3 Model Design

ForestSim is a CHANS model of forest growth and harvest cycles in the Western Upper Peninsula (UP) of Michigan, USA. The Western UP is a heavily forested region with approximately 90% of the forest ownership divided between state and federal public forests, large-tract private forests for timber harvesting, and NIPF owners [17, 21]. The region also has high energy costs due to reliance upon generating stations in neighboring states which makes it an ideal candidate for local power generation using biofuel. Due to the region's forest ownership structure, NIPF owners are expected to have a significant impact upon the biomass available for generation.

The environment is based upon even-aged forest stands wherein each pixel represents approximately one hundred and twenty by one hundred and twenty meters. Stands are allocated a forest type based upon data from the National Land Cover Database 2011 (NLCD) [14]. After forest types are assigned, a grid of Perlin noise is generated and values are used allocate the average diameter at breast height (d.b.h.) for the trees in the stand. Forestry stocking tables are then used to determine the number of trees in the stand, with overall stocking values for the region corresponding to observed conditions [21]. Agents will use the stand values along with forest management plans to determine the best time to harvest.

NIPF agents are embedded within the landscape using an overlaid, digitized parcel boundary map to assign agents to all properties with a minimum of ten forested acres. Decision-making schemes were developed based upon semistructured interviews with local forest owners and a general survey of NIPF owners in the UP [17, 8]. Agents are assigned a schema that is categorized as either rational economic decision making, or optimizing for ecosystem services. The economic decision making scheme contains rules that have the agent optimize for profits from their land. In contrast, the ecosystem services rules have the agent optimize for intangible uses of their land. Logistic regression was used to generate probabilities of adopting alternative management activities based on correlations with key forest owner attributes socioeconomic status, residency, attitudes toward alternative forest management options, and so forth. Agents use their assigned schema to determine whether and how to participate in a simulated market for biomass under alternative VIP incentivization scenarios.

When the model is running, the forest model is subjected to a growth cycle that is based upon typical trees for representational trees in the region (e.g. Acer Rubrum, Pinus Strobus, etc.) [9]. Natural thinning is also taken into account by the model and overstocked tree stands will be subjected to a randomized rate of die off. To allow for better performance at run time, regrowth and stocking

model is done through parallel processing by segmenting data structures and running them on separate threads, scaled to the number of cores available.

During the simulation, agents use the current state of their forest stands and assigned decision-making schema to make a "harvest, no harvest" decision. When an agent makes a harvest decision, biomass is harvested from the forest in accordance with standard forestry practices [22]. Harvested biomass is then allocated for use by a biorefinery as appropriate and the stand is reseeded. After all agents have had an opportunity to run, the forest stands are updated to reflect any growth that may have occurred.

4 Conclusion

ForestSim makes it possible to explore how alternative VIP incentivization schemes impact potentially competing dimensions of sustainable bioenergy. To do this ForestSim compares the simulated results of alternative schemes as part of a multidimensional sustainability scorecard. Consistent biomass volume results in a high score from the standpoint of bioenergy plants. Public access to forest lands results in a high score for the social dimension. Using this scorecard, ForestSim demonstrates that VIP incentivization with an agglomeration bonus produces optimal sustainability outcomes when compared to no incentivization and alternative VIP schemes. Ultimately, ForestSim may allow us to develop a generalized framework to evaluate policy options in other heavily forested regions of the United States currently underutilizing available bioenergy potential.

References

- [1] Aguilar, F.X., Cai, Z., D'Amato, A.W.: Non-industrial private forest owner's willingness-to-harvest: How higher timber prices influence woody biomass supply. Biomass and Bioenergy 71, 202–215 (Dec 2014)
- [2] An, L.: Modeling human decisions in coupled human and natural systems: Review of agent-based models. Modeling Human Decisions 229, 25–36 (Mar 2012)
- [3] Bakker, M.M., Alam, S.J., van Dijk, J., Rounsevell, M.D.A.: Land-use change arising from rural land exchange: an agent-based simulation model. Landscape Ecology 30(2), 273–286 (2015)
- [4] Beach, R.H., Pattanayak, S.K., Yang, J.C., Murray, B.C., Abt, R.C.: Econometric studies of non-industrial private forest management: a review and synthesis. Forest Policy and Economics 7(3), 261–281 (Mar 2005)
- [5] Bert, F.E., Podest, G.P., Rovere, S.L., Menndez, n.N., North, M., Tatara, E., Laciana, C.E., Weber, E., Toranzo, F.R.: An agent based model to simulate structural and land use changes in agricultural systems of the argentine pampas. Ecological Modelling 222(19), 3486–3499 (Oct 2011)
- [6] Bone, C., Dragievi, S.: Evaluating Spatio-temporal Complexities of Forest Management: An Integrated Agent-based Modeling and GIS Approach. Environmental Modeling & Assessment 14(4), 481–496 (2009)
- [7] Brown, C., Bakam, I., Smith, P., Matthews, R.: An agent-based modelling approach to evaluate factors influencing bioenergy crop adoption in north-east Scotland. GCB Bioenergy 8(1), 226–244 (Jan 2016)

- [8] Brunner, A., Currie, W.S., Miller, S.: Cellulosic ethanol production: Landscape scale net carbon strongly affected by forest decision making. Biomass and Bioenergy 83, 32–41 (Dec 2015)
- [9] Burns, R.M., Barbara, H.: Silvics of North America: 1. conifers; 2. hardwoods. No. 654 in Agriculture Handbook, United States Department of Agriculture, Forest Service, Washington, D.C. (1990)
- [10] Fooks, J.R., Higgins, N., Messer, K.D., Duke, J.M., Hellerstein, D., Lynch, L.: Conserving Spatially Explicit Benefits in Ecosystem Service Markets: Experimental Tests of Network Bonuses and Spatial Targeting. American Journal of Agricultural Economics 98(2), 468–488 (Mar 2016)
- [11] Gan, J., Langeveld, J.W.A., Smith, C.T.: An Agent-Based Modeling Approach for Determining Corn Stover Removal Rate and Transboundary Effects. Environmental Management 53(2), 333–342 (2014)
- [12] Guzy, M.R., Smith, C.L., Bolte, J.P., Hulse, D.W., Gregory, S.V.: Policy Research Using Agent-Based Modeling to Assess Future Impacts of Urban Expansion into Farmlands and Forests. Ecology and Society 13(1) (2008)
- [13] Heairet, A., Choudhary, S., Miller, S., Xu, M.: Beyond life cycle analysis: Using an agent-based approach to model the emerging bio-energy industry. In: Proceedings of 2012 IEEE International Symposium on Sustainable Systems and Technology (ISSST). pp. 1–5. Boston, MA (May 2012)
- [14] Homer, C.G., Dewitz, J.A., Yang, L., Jin, S., Danielson, P., Xian, G., Coulston, J., Herold, N.D., Wickham, J.D., Megown, K.: Completion of the 2011 National Land Cover Database for the conterminous United States-Representing a decade of land cover change information. Photogrammetric Engineering and Remote Sensing 81(5), 345–354 (2015)
- [15] Kelley, H., Evans, T.: The relative influences of land-owner and landscape heterogeneity in an agent-based model of land-use. Ecological Economics 70(6), 1075–1087 (Apr 2011)
- [16] Leahy, J.E., Reeves, E.G., Bell, K.P., Straub, C.L., Wilson, J.S.: Agent-Based Modeling of Harvest Decisions by Small Scale Forest Landowners in Maine, USA. International Journal of Forestry Research 2013, 12 (2013)
- [17] Lind-Riehl, J., Jeltema, S., Morrison, M., Shirkey, G., Mayer, A.L., Rouleau, M., Winkler, R.: Family legacies and community networks shape private forest management in the western Upper Peninsula of Michigan (USA). Land Use Policy 45, 95–102 (May 2015)
- [18] Manson, S.M., Evans, T.: Agent-based modeling of deforestation in southern Yucatn, Mexico, and reforestation in the Midwest United States. Proceedings of the National Academy of Sciences 104(52), 20678–20683 (Dec 2007)
- [19] Matthews, R.B., Gilbert, N.G., Roach, A., Polhill, J.G., Gotts, N.M.: Agent-based land-use models: a review of applications. Landscape Ecology 22(10), 1447–1459 (2007)
- [20] Morgan, F.J., Daigneault, A.J.: Estimating Impacts of Climate Change Policy on Land Use: An Agent-Based Modelling Approach. PLoS ONE 10(5), e0127317 (May 2015)
- [21] Pugh, S.A., Pedersen, L.D., Heym, D.C., Piva, R.J., Woodall, C.W., Barnett, C.J., Kurtz, C.M., Moser, W.K.: Michigan's Forests 2009. Tech. Rep. NRS-66, U.S. Department of Agriculture, Forest Service, Northern Research Station, Newtown Square, PA (2012)
- [22] Wenger, K.F. (ed.): Forestry Handbook. John Wiley & Sons, Inc., second edition edn. (1984)