

FOUNDATIONS OF STATISTICAL DECISION MAKING

Measuring Uncertainty

May 29, 2018

Aaron R. Baggett, Ph.D.

Department of Physical Therapy
University of Mary Hardin-Baylor
PHTH 7147: Critical Inquiry I



PREVIEW

Preview

1. May 29, 2018

- *Fundamentals of Statistical Inference: Measuring Uncertainty*

2. June 5, 2018

- *Fundamentals of Statistical Decision Making: Comparing Multiple Groups*

3. June 5, 2018

- *Fundamentals of Statistical Decision Making: Relationships and Prediction*

OUTLINE

Outline

- Descriptive vs. inferential statistics
- The normal distribution
- Comparing groups
- Statistical/practical significance

Resources

- Slides, data, and handouts available at:

bit.ly/umhb_dpt

STATISTICS

- Experimentation and observation:
 1. Measurement of uncertainty
 2. Examination of the consequences of that uncertainty

- Two fundamental branches
 1. Descriptive statistics
 - Summarize data
 - Condense larger themes
 2. Inferential statistics
 - Infer meaning
 - Test predictions

EXAMPLE

Low Birth Weight Study

- Baystate Medical Center, Springfield, MA.
- Sample of 189 births in 1986
- Risk factors in low birth weight babies

Low Birth Weight Study

Age	Weight	Race	Smoking Status	Birth Weight
19	182	Black	Non-Smoker	5.56
33	155	Other	Non-Smoker	5.62
20	105	White	Smoker	5.64
21	108	White	Smoker	5.72
18	107	White	Smoker	5.73
21	124	Other	Non-Smoker	5.78

DESCRIPTIVE STATISTICS

Descriptive Statistics

- How many babies were born at low birth weight (< 5.5 lbs.)?
- How many mothers smoked during pregnancy?
- How much did the average baby weigh?
 - Given mothers' smoking status
 - Given mothers' race

Descriptive Statistics

Question:

Do babies born to mothers who smoked during pregnancy weigh less than those born to mothers who did not?

Descriptive Statistics

Question:

Do babies born to mothers who smoked during pregnancy weigh less than those born to mothers who did not?

- How should we answer this question?

Descriptive Statistics

Question:

[ON AVERAGE], do babies born to mothers who smoked during pregnancy weigh less than those born to mothers who did not?

Smoking Status	n	Min.	Max.	M	SD
Non-Smoker	115	2.25	11.00	6.74	1.66
Smoker	74	1.56	9.34	6.11	1.46

Descriptive Statistics

Question:

1. Based on our sample, what are we left to assume about the weights of babies *in the population* born to smoking and non-smoking mothers?

Descriptive Statistics

Question:


1. Based on our sample, what are we left to assume about the weights of babies *in the population* born to smoking and non-smoking mothers?
 - That the sample estimates represent the population parameters

Smoking Status	n	Min.	Max.	M	SD
Non-Smoker	115	2.25	11.00	6.74	1.66
Smoker	74	1.56	9.34	6.11	1.46


Descriptive Statistics

- In fact, we assume that the population distribution of baby weights is “normal”


Normal Distribution




Normal Distribution



Normal Distribution



Normal Distribution



INFERENCEAL STATISTICS

Inferential Statistics

- More useful than descriptives
- Allow for making predictions or generalizations
- Key to hypothesis testing
- Two varieties:
 1. 95% confidence intervals (CIs)
 2. Null-hypothesis significance testing (NHST)

Inferential Statistics


Question:

Do babies born to mothers who smoked during pregnancy weigh less than those born to mothers who did not?

Inferential Statistics::Confidence Intervals

- Since we are interested in the mean difference in birth weights *in the population*, a first inferential step is to calculate a 95% confidence interval
- Confidence intervals are a plausible range of values for a population parameter
- Point estimates often may not represent the population parameter
- CIs are more likely to capture the population parameter than a point estimate alone

Inferential Statistics::Confidence Intervals



Inferential Statistics::Confidence Intervals

- 95% CI:

$$(M_1 - M_2) \pm 2 \times \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

Inferential Statistics::Confidence Intervals

- 95% CI:

$$(6.11_{NS} - 6.74_S) \pm 2 \times \sqrt{\frac{2.12_{NS}}{74_{NS}} + \frac{2.75_S}{115_S}}$$

Inferential Statistics::Confidence Intervals

- 95% CI:


$$-0.63 \pm (2 \times 0.229) = \\ (-1.09, -0.17)$$

- Thus, we can be 95% confident that, in the population, the true difference in birth weight of babies born to smoking mothers compared to those born to non-smoking mothers is between -1.09 and -0.17 lbs. less, on average.

Inferential Statistics::Confidence Intervals

- In other words, if we replicated this study 25 times, 24 of the 25 replications would include the true population parameter

Inferential Statistics::Confidence Intervals



Inferential Statistics

Question:

Do babies born to mothers who smoked during pregnancy weigh [STATISTICALLY SIGNIFICANTLY] less than those born to mothers who did not?

- How should we answer this question?

Inferential Statistics::Hypothesis Testing

- What do we mean by statistical significance?

Inferential Statistics::Hypothesis Testing

- What do we mean by statistical significance?
- Observed differences which exceed “normality.”

Inferential Statistics::Hypothesis Testing

- We usually consider differences beyond ± 2 SDs from M to be “statistically significant”
- **NOTE:** Statistical significance \neq practical significance

Low Birth Weight Study

Question:

- Do babies born to mothers who smoked during pregnancy weigh less than those born to mothers who did not?

Low Birth Weight Study

Hypotheses:

- H_0 : There is no mean difference in the birth weight of babies born to mothers who did and did not smoke during pregnancy
 - $(M_{non-smoker} - M_{smoker} = 0)$
- H_1 : There is some difference in the birth weight of babies born to mothers who did and did not smoke during pregnancy
 - $(M_{non-smoker} - M_{smoker} \neq 0)$

Low Birth Weight Study

- Let's test our hypothesis using an independent-samples t -test
 - IV: Mothers' smoking status (smoker, non-smoker)
 - DV: Baby birth weight

$$t = \frac{\overline{X}_{non-smokers} - \overline{X}_{smokers}}{\sqrt{\frac{s_{non-smokers}^2}{N_{non-smokers}} + \frac{s_{smokers}^2}{N_{smokers}}}}$$

Results

Table 1: Results of Independent-Samples *t*-Test


	Non-Smokers			Smokers			<i>t</i> (187)	<i>p</i>	ω^2
	<i>n</i>	<i>M</i>	<i>SD</i>	<i>n</i>	<i>M</i>	<i>SD</i>			
Baby birth weight	115	6.74	1.66	47	6.11	1.46	2.63	0.009	0.008

Note: *M* = Mean; *SD* = Standard deviation


Results

- Assuming the null hypothesis, in reality, is true, the probability of obtaining a mean difference in birth weight ≥ 0.62 lbs. is 0.009 (0.90%)
- Birth weights appear to differ statistically significantly

Results



Results



Results

- But, is the difference of $M = 0.62$ lbs. meaningful?

Results

- But, is the difference of $M = 0.62$ lbs. meaningful?
- A meaningful difference implies practicality or usefulness in the real world

Results

- But, is the difference of $M = 0.62$ lbs. meaningful?
- A meaningful difference implies practicality or usefulness in the real world
- Effect size (ω^2): Proportion of variance explained in the model

Results

- But, is the difference of $M = 0.62$ lbs. meaningful?
- A meaningful difference implies practicality or usefulness in the real world
- Effect size (ω^2): Proportion of variance explained in the model
- Smoking status explains 0.009 (0.90%) of the variance in baby birth weight

Results

- But, is the difference of $M = 0.62$ lbs. meaningful?
- A meaningful difference implies practicality or usefulness in the real world
- Effect size (ω^2): Proportion of variance explained in the model
- Smoking status explains 0.009 (0.90%) of the variance in baby birth weight
- Thus, $100\% - 0.991\% = 99.10\%$ of the variance in baby birth weight is left unexplained

RECAP

Recap

- Descriptive statistics allow us to summarize data from a sample
- Inferential statistics allow us to predict and generalize about a population
- Hypothesis testing allows us to construct a sense of meaning about the world

Next Time

- Making decisions using hypothesis testing and prediction
 - Statistical variables
 - Multiple group comparisons (ANOVA)
 - Predicting outcomes (Regression)