PK (ADME)
PD (named activity)
Risk/severity (unspecified mechanism)
Therapeutic efficacy (unspecified mechanism)
low test support (n<20)

DrugBank S1: per-type fusion-minus-graph-only AUPRC (62 evaluable types)

[16] #Drug1 may decrease the neuromuscular blocking acti ...
[83] #Drug1 may increase the vasopressor activities of # ...
[22] #Drug1 may increase the stimulatory activities of # ...
[56] #Drug1 may increase the nephrotoxic activities of # ...
[5] #Drug1 may increase the anticoagulant activities of ...
[20] #Drug1 may increase the neuromuscular blocking acti ...
[54] #Drug1 may increase the hyponatremic activities of ...
[33] #Drug1 may increase the immunosuppressive activitie ...
[32] The risk or severity of QTc prolongation can be inc ...
[47] #Drug1 may increase the myopathic rhabdomyolysis ac ...
[19] #Drug1 may increase the QTc-prolonging activities o ...
[52] #Drug1 may increase the antiplatelet activities of ...
[28] #Drug1 may decrease the diuretic activities of #Dru ...
[37] #Drug1 may increase the vasodilatory activities of ...
[59] #Drug1 may increase the hypotensive activities of # ...
[65] The risk or severity of bleeding can be increased w ...
[68] #Drug1 may increase the analgesic activities of #Dr ...
[9] #Drug1 may increase the antihypertensive activities ...
[24] #Drug1 may increase the atrioventricular blocking (...
[63] #Drug1 may increase the serotonergic activities of ...
[10] The serum concentration of the active metabolites o ...
[2] The bioavailability of #Drug2 can be decreased when ...
[36] #Drug1 may decrease the antihypertensive activities ...
[80] #Drug1 may increase the thrombogenic activities of ...
[26] #Drug1 may increase the neuroexcitatory activities ...
[11] #Drug1 may decrease the anticoagulant activities of ...
[31] #Drug1 may increase the sedative activities of #Dru ...
[57] #Drug1 may decrease the cardiotoxic activities of # ...
[53] #Drug1 may increase the bradycardic activities of # ...
[46] The metabolism of #Drug2 can be decreased when comb ...
[39] #Drug1 may increase the respiratory depressant acti ...
[69] The therapeutic efficacy of #Drug2 can be decreased ...
[48] The risk or severity of adverse effects can be incr ...
[81] #Drug1 may increase the arrhythmogenic activities o ...
[17] #Drug1 can cause an increase in the absorption of # ...
[75] #Drug1 may decrease the sedative activities of #Dru ...
[76] The serum concentration of the active metabolites o ...
[72] The serum concentration of #Drug2 can be increased ...
[38] #Drug1 may increase the constipating activities of ...
[15] #Drug1 may increase the central nervous system depr ...
[71] #Drug1 may decrease the excretion rate of #Drug2 wh ...
[29] #Drug1 may increase the orthostatic hypotensive act ...
[66] #Drug1 can cause a decrease in the absorption of #D ...
[21] #Drug1 may increase the adverse neuromuscular activ ...
[23] #Drug1 may increase the hypocalcemic activities of ...
[8] #Drug1 may increase the hypoglycemic activities of ...
[50] #Drug1 may increase the hypercalcemic activities of ...
[3] The metabolism of #Drug2 can be increased when comb ...
[84] #Drug1 may increase the tachycardic activities of # ...
[74] The serum concentration of #Drug2 can be decreased ...
[60] #Drug1 may decrease the stimulatory activities of # ...
[79] #Drug1 may increase the hepatotoxic activities of # ...
[73] #Drug1 may increase the fluid retaining activities ...
[70] #Drug1 may increase the hypertensive activities of ...
[7] The therapeutic efficacy of #Drug2 can be increased ...
[1] #Drug1 may increase the anticholinergic activities ...
[4] #Drug1 may decrease the vasoconstricting activities ...
[82] #Drug1 may increase the hypokalemic activities of # ...
[67] #Drug1 may increase the hyperkalemic activities of ...
[13] #Drug1 may decrease the bronchodilatory activities ...
[35] #Drug1 may increase the antipsychotic activities of ...
[18] #Drug1 may increase the vasoconstricting activities ...

−0.4 −0.2 0.0 0.2 0.4 0.6
Δ AUPRC (fusion=concat − graph-only), mean ± std over 3 seeds

DrugBank S2: per-type fusion-minus-graph-only AUPRC (28 evaluable types)

[5] #Drug1 may increase the anticoagulant activities of ...
[56] #Drug1 may increase the nephrotoxic activities of # ...
[39] #Drug1 may increase the respiratory depressant acti ...
[20] #Drug1 may increase the neuromuscular blocking acti ...
[26] #Drug1 may increase the neuroexcitatory activities ...
[59] #Drug1 may increase the hypotensive activities of # ...
[19] #Drug1 may increase the QTc-prolonging activities o ...
[11] #Drug1 may decrease the anticoagulant activities of ...
[48] The risk or severity of adverse effects can be incr ...
[66] #Drug1 can cause a decrease in the absorption of #D ...
[52] #Drug1 may increase the antiplatelet activities of ...
[71] #Drug1 may decrease the excretion rate of #Drug2 wh ...
[36] #Drug1 may decrease the antihypertensive activities ...
[69] The therapeutic efficacy of #Drug2 can be decreased ...
[53] #Drug1 may increase the bradycardic activities of # ...
[9] #Drug1 may increase the antihypertensive activities ...
[46] The metabolism of #Drug2 can be decreased when comb ...
[29] #Drug1 may increase the orthostatic hypotensive act ...
[76] The serum concentration of the active metabolites o ...
[8] #Drug1 may increase the hypoglycemic activities of ...
[72] The serum concentration of #Drug2 can be increased ...
[24] #Drug1 may increase the atrioventricular blocking (...
[74] The serum concentration of #Drug2 can be decreased ...
[3] The metabolism of #Drug2 can be increased when comb ...
[15] #Drug1 may increase the central nervous system depr ...
[82] #Drug1 may increase the hypokalemic activities of # ...
[70] #Drug1 may increase the hypertensive activities of ...
[84] #Drug1 may increase the tachycardic activities of # ...

−0.4 −0.2 0.0 0.2 0.4 0.6
Δ AUPRC (fusion=concat − graph-only), mean ± std over 3 seeds