
PK (ADME)
PD (named activity)
Risk/severity (unspecified mechanism)
Therapeutic efficacy (unspecified mechanism)
low test support (n<20)
[16] #Drug1 may decrease the neuromuscular blocking activities of #D...
[83] #Drug1 may increase the vasopressor activities of #Drug2.
[22] #Drug1 may increase the stimulatory activities of #Drug2.
[56] #Drug1 may increase the nephrotoxic activities of #Drug2.
[5] #Drug1 may increase the anticoagulant activities of #Drug2.
[20] #Drug1 may increase the neuromuscular blocking activities of #D...
[54] #Drug1 may increase the hyponatremic activities of #Drug2.
[33] #Drug1 may increase the immunosuppressive activities of #Drug2.
[32] The risk or severity of QTc prolongation can be increased when ...
[47] #Drug1 may increase the myopathic rhabdomyolysis activities of ...
[19] #Drug1 may increase the QTc-prolonging activities of #Drug2.
[52] #Drug1 may increase the antiplatelet activities of #Drug2.
[28] #Drug1 may decrease the diuretic activities of #Drug2.
[37] #Drug1 may increase the vasodilatory activities of #Drug2.
[59] #Drug1 may increase the hypotensive activities of #Drug2.
[65] The risk or severity of bleeding can be increased when #Drug1 i...
[68] #Drug1 may increase the analgesic activities of #Drug2.
[9] #Drug1 may increase the antihypertensive activities of #Drug2.
[24] #Drug1 may increase the atrioventricular blocking (AV block) ac...
[63] #Drug1 may increase the serotonergic activities of #Drug2.
[10] The serum concentration of the active metabolites of #Drug2 can...
[2] The bioavailability of #Drug2 can be decreased when combined wi...
[36] #Drug1 may decrease the antihypertensive activities of #Drug2.
[80] #Drug1 may increase the thrombogenic activities of #Drug2.
[26] #Drug1 may increase the neuroexcitatory activities of #Drug2.
[11] #Drug1 may decrease the anticoagulant activities of #Drug2.
[31] #Drug1 may increase the sedative activities of #Drug2.
[57] #Drug1 may decrease the cardiotoxic activities of #Drug2.
[53] #Drug1 may increase the bradycardic activities of #Drug2.
[46] The metabolism of #Drug2 can be decreased when combined with #D...
[39] #Drug1 may increase the respiratory depressant activities of #D...
[69] The therapeutic efficacy of #Drug2 can be decreased when used i...
[48] The risk or severity of adverse effects can be increased when #...
[81] #Drug1 may increase the arrhythmogenic activities of #Drug2.
[17] #Drug1 can cause an increase in the absorption of #Drug2 result...
[75] #Drug1 may decrease the sedative activities of #Drug2.
[76] The serum concentration of the active metabolites of #Drug2 can...
[72] The serum concentration of #Drug2 can be increased when it is c...
[38] #Drug1 may increase the constipating activities of #Drug2.
[15] #Drug1 may increase the central nervous system depressant (CNS ...
[71] #Drug1 may decrease the excretion rate of #Drug2 which could re...
[29] #Drug1 may increase the orthostatic hypotensive activities of #...
[66] #Drug1 can cause a decrease in the absorption of #Drug2 resulti...
[21] #Drug1 may increase the adverse neuromuscular activities of #Dr...
[23] #Drug1 may increase the hypocalcemic activities of #Drug2.
[8] #Drug1 may increase the hypoglycemic activities of #Drug2.
[50] #Drug1 may increase the hypercalcemic activities of #Drug2.
[3] The metabolism of #Drug2 can be increased when combined with #D...
[84] #Drug1 may increase the tachycardic activities of #Drug2.
[74] The serum concentration of #Drug2 can be decreased when it is c...
[60] #Drug1 may decrease the stimulatory activities of #Drug2.
[79] #Drug1 may increase the hepatotoxic activities of #Drug2.
[73] #Drug1 may increase the fluid retaining activities of #Drug2.
[70] #Drug1 may increase the hypertensive activities of #Drug2.
[7] The therapeutic efficacy of #Drug2 can be increased when used i...
[1] #Drug1 may increase the anticholinergic activities of #Drug2.
[4] #Drug1 may decrease the vasoconstricting activities of #Drug2.
[82] #Drug1 may increase the hypokalemic activities of #Drug2.
[67] #Drug1 may increase the hyperkalemic activities of #Drug2.
[13] #Drug1 may decrease the bronchodilatory activities of #Drug2.
[35] #Drug1 may increase the antipsychotic activities of #Drug2.
[18] #Drug1 may increase the vasoconstricting activities of #Drug2.
−0.4
−0.2
0.0
0.2
0.4
0.6
Δ AUPRC, Fusion (GIN+text) − Graph-only (GIN), mean ± st