

MA 105 D1 Lecture 8

Ravi Raghunathan

Department of Mathematics

August 17, 2017

The Darboux integral

Riemann integration

Partitions

Definition: Given a closed interval $[a, b]$, a **partition** P of $[a, b]$ is simply a collections of points

$$P = \{a = x_0 < x_1 < \dots < x_{n-1} < x_n = b\}.$$

Partitions

Definition: Given a closed interval $[a, b]$, a **partition** P of $[a, b]$ is simply a collections of points

$$P = \{a = x_0 < x_1 < \dots < x_{n-1} < x_n = b\}.$$

We can think of the points of the partition as dividing the original interval $[a, b]$ into sub-intervals $I_j = [x_{j-1}, x_j]$, $1 \leq j \leq n$. Indeed $I = \cup_j I_j$ and if two sub-interval intersect, they have at most one point in common. Hence, the notation “partition”.

Partitions

Definition: Given a closed interval $[a, b]$, a **partition** P of $[a, b]$ is simply a collections of points

$$P = \{a = x_0 < x_1 < \dots < x_{n-1} < x_n = b\}.$$

We can think of the points of the partition as dividing the original interval $[a, b]$ into sub-intervals $I_j = [x_{j-1}, x_j]$, $1 \leq j \leq n$. Indeed $I = \cup_j I_j$ and if two sub-interval intersect, they have at most one point in common. Hence, the notation “partition”.

Definition: A partition $P' = \{a = x'_0 < x'_1 < \dots < x'_m = b\}$ is said to be a **refinement** of the partition P if for each $x_i \in P$, there exists an $x'_j \in P'$ such that $x_i = x'_j$.

Partitions

Definition: Given a closed interval $[a, b]$, a **partition** P of $[a, b]$ is simply a collections of points

$$P = \{a = x_0 < x_1 < \dots < x_{n-1} < x_n = b\}.$$

We can think of the points of the partition as dividing the original interval $[a, b]$ into sub-intervals $I_j = [x_{j-1}, x_j]$, $1 \leq j \leq n$. Indeed $I = \cup_j I_j$ and if two sub-interval intersect, they have at most one point in common. Hence, the notation “partition”.

Definition: A partition $P' = \{a = x'_0 < x'_1 < \dots < x'_m = b\}$ is said to be a **refinement** of the partition P if for each $x_i \in P$, there exists an $x'_j \in P'$ such that $x_i = x'_j$.

Intuitively, a refinement P' of a partition P will break some of the sub-intervals in P into smaller sub-intervals. **Any two partitions have a common refinement.**

Lower and Upper sums

Given a partition $P = \{a = x_0 < x_1 < \dots < x_{b-1} < x_n = b\}$ and a function $f : [a, b] \rightarrow \mathbb{R}$, we define two associated quantities. First we set:

$$M_i = \sup_{x \in [x_{i-1}, x_i]} f(x) \quad \text{and} \quad m_i = \inf_{x \in [x_{i-1}, x_i]} f(x), \quad 1 \leq i \leq n$$

Definition: We define the **Lower sum** as


$$L(f, P) = \sum_{j=1}^n m_j (x_j - x_{j-1}).$$

Similarly, we can define the **Upper sum** as


$$U(f, P) = \sum_{j=1}^n M_j (x_j - x_{j-1}).$$

In case the words “infimum” and “supremum” bother you, you can think “minimum” and “maximum” most of the time since we will usually be dealing with continuous functions on $[a, b]$.

Lower and Upper Darboux sums



A picture for a non-monotonic function



One basic example

In order to illustrate what we are saying we will take the following basic example. Let $[a, b] = [0, 1]$ and let $f(x) = x$.

One basic example

In order to illustrate what we are saying we will take the following basic example. Let $[a, b] = [0, 1]$ and let $f(x) = x$.

One of the most natural partitions on an interval is a partition that divides the interval into sub-intervals of equal length. For $[0, 1]$, this is

$$P_n = \{0 < 1/n < 2/n < \dots < (n-1)/n < 1\}.$$

One basic example

In order to illustrate what we are saying we will take the following basic example. Let $[a, b] = [0, 1]$ and let $f(x) = x$.

One of the most natural partitions on an interval is a partition that divides the interval into sub-intervals of equal length. For $[0, 1]$, this is

$$P_n = \{0 < 1/n < 2/n < \dots < (n-1)/n < 1\}.$$

On the interval $I_j = [\frac{j-1}{n}, \frac{j}{n}]$, where does the function $f(x) = x$ take its minimum? its maximum?

One basic example

In order to illustrate what we are saying we will take the following basic example. Let $[a, b] = [0, 1]$ and let $f(x) = x$.

One of the most natural partitions on an interval is a partition that divides the interval into sub-intervals of equal length. For $[0, 1]$, this is

$$P_n = \{0 < 1/n < 2/n < \dots < (n-1)/n < 1\}.$$

On the interval $I_j = [\frac{j-1}{n}, \frac{j}{n}]$, where does the function $f(x) = x$ take its minimum? its maximum?

Clearly, the minimum $m_j = \frac{j-1}{n}$ is attained at $\frac{j-1}{n}$ and the maximum $M_j = \frac{j}{n}$ at $\frac{j}{n}$. And finally, $\frac{j}{n} - \frac{j-1}{n} = \frac{1}{n}$, for all $1 \leq j \leq n$.

One basic example

In order to illustrate what we are saying we will take the following basic example. Let $[a, b] = [0, 1]$ and let $f(x) = x$.

One of the most natural partitions on an interval is a partition that divides the interval into sub-intervals of equal length. For $[0, 1]$, this is

$$P_n = \{0 < 1/n < 2/n < \dots < (n-1)/n < 1\}.$$

On the interval $I_j = [\frac{j-1}{n}, \frac{j}{n}]$, where does the function $f(x) = x$ take its minimum? its maximum?

Clearly, the minimum $m_j = \frac{j-1}{n}$ is attained at $\frac{j-1}{n}$ and the maximum $M_j = \frac{j}{n}$ at $\frac{j}{n}$. And finally, $\frac{j}{n} - \frac{j-1}{n} = \frac{1}{n}$, for all $1 \leq j \leq n$.

An example of a refinement of P_n is P_{2n} , or, more generally, P_{kn} for any natural number k .

The Darboux integrals

We now define the lower Darboux integral of f by

$$L(f) = \sup\{L(f, P) : P \text{ is a partition of } [a, b]\},$$

where the supremum is taken over all partitions of $[a, b]$.

The Darboux integrals

We now define the lower Darboux integral of f by

$$L(f) = \sup\{L(f, P) : P \text{ is a partition of } [a, b]\},$$

where the supremum is taken over all partitions of $[a, b]$.

and similarly the upper Darboux integral of f by

$$U(f) = \inf\{U(f, P) : P \text{ is a partition of } [a, b]\},$$

and again the infimum is over all partitions of $[a, b]$. (This time there is no escaping inf and sup!)

The Darboux integrals

We now define the lower Darboux integral of f by

$$L(f) = \sup\{L(f, P) : P \text{ is a partition of } [a, b]\},$$

where the supremum is taken over all partitions of $[a, b]$.

and similarly the upper Darboux integral of f by

$$U(f) = \inf\{U(f, P) : P \text{ is a partition of } [a, b]\},$$

and again the infimum is over all partitions of $[a, b]$. (This time there is no escaping inf and sup!)

If $L(f) = U(f)$, then we say that f is Darboux-integrable and define

$$\int_a^b f(t) dt := U(f) = L(f).$$

This common value of the two integrals is called the Darboux integral.

Back to the example

Let us calculate $L(f, P_n)$ and $U(f, P_n)$ in the example we gave.

Back to the example

Let us calculate $L(f, P_n)$ and $U(f, P_n)$ in the example we gave.

$$L(f, P_n) = \sum_{j=1}^n \frac{(j-1)}{n} \cdot \frac{1}{n} = \sum_{j=0}^{n-1} \frac{j}{n^2}.$$

Back to the example

Let us calculate $L(f, P_n)$ and $U(f, P_n)$ in the example we gave.

$$L(f, P_n) = \sum_{j=1}^n \frac{(j-1)}{n} \cdot \frac{1}{n} = \sum_{j=0}^{n-1} \frac{j}{n^2}.$$

This can be evaluated explicitly:

$$L(f, P_n) = \frac{n(n-1)}{2} \cdot \frac{1}{n^2} = \frac{1}{2} - \frac{1}{2n}.$$

Back to the example

Let us calculate $L(f, P_n)$ and $U(f, P_n)$ in the example we gave.

$$L(f, P_n) = \sum_{j=1}^n \frac{(j-1)}{n} \cdot \frac{1}{n} = \sum_{j=0}^{n-1} \frac{j}{n^2}.$$

This can be evaluated explicitly:

$$L(f, P_n) = \frac{n(n-1)}{2} \cdot \frac{1}{n^2} = \frac{1}{2} - \frac{1}{2n}.$$

Similary, we can check that

$$U(f, P_n) = \frac{n(n+1)}{2} \cdot \frac{1}{n^2} = \frac{1}{2} + \frac{1}{2n}.$$

Back to the example

Let us calculate $L(f, P_n)$ and $U(f, P_n)$ in the example we gave.

$$L(f, P_n) = \sum_{j=1}^n \frac{(j-1)}{n} \cdot \frac{1}{n} = \sum_{j=0}^{n-1} \frac{j}{n^2}.$$

This can be evaluated explicitly:

$$L(f, P_n) = \frac{n(n-1)}{2} \cdot \frac{1}{n^2} = \frac{1}{2} - \frac{1}{2n}.$$

Similary, we can check that

$$U(f, P_n) = \frac{n(n+1)}{2} \cdot \frac{1}{n^2} = \frac{1}{2} + \frac{1}{2n}.$$

Can we conclude that the Darboux integral is $1/2$ by letting $n \rightarrow \infty$?

Back to the example

Let us calculate $L(f, P_n)$ and $U(f, P_n)$ in the example we gave.

$$L(f, P_n) = \sum_{j=1}^n \frac{(j-1)}{n} \cdot \frac{1}{n} = \sum_{j=0}^{n-1} \frac{j}{n^2}.$$

This can be evaluated explicitly:

$$L(f, P_n) = \frac{n(n-1)}{2} \cdot \frac{1}{n^2} = \frac{1}{2} - \frac{1}{2n}.$$

Similary, we can check that

$$U(f, P_n) = \frac{n(n+1)}{2} \cdot \frac{1}{n^2} = \frac{1}{2} + \frac{1}{2n}.$$

Can we conclude that the Darboux integral is $1/2$ by letting $n \rightarrow \infty$? Unfortunately, no.

Useful properties of the Darboux sums

Since, for any partition P , $L(f, P) \leq U(f, P)$, we have

$$L(f) \leq U(f).$$

Useful properties of the Darboux sums

Since, for any partition P , $L(f, P) \leq U(f, P)$, we have

$$L(f) \leq U(f).$$

In fact, for any two partitions P_1 and P_2 , we have

$$L(f, P_1) \leq U(f, P_2).$$

Useful properties of the Darboux sums

Since, for any partition P , $L(f, P) \leq U(f, P)$, we have

$$L(f) \leq U(f).$$

In fact, for any two partitions P_1 and P_2 , we have

$$L(f, P_1) \leq U(f, P_2).$$

This is easy to see - the lower sum computes the sum of the areas of rectangles that lie entirely below the curve while the upper sum computes the sum of the areas of rectangles whose “tops” lie above the curve.

Useful properties of the Darboux sums

Since, for any partition P , $L(f, P) \leq U(f, P)$, we have

$$L(f) \leq U(f).$$

In fact, for any two partitions P_1 and P_2 , we have

$$L(f, P_1) \leq U(f, P_2).$$

This is easy to see - the lower sum computes the sum of the areas of rectangles that lie entirely below the curve while the upper sum computes the sum of the areas of rectangles whose “tops” lie above the curve.

One of the most useful properties of the Darboux sums is the following. If P' is a refinement of P then obviously

$$L(f, P) \leq L(f, P') \leq U(f, P') \leq U(f, P).$$

Riemann Sums

There is another way of getting at the integral due to Riemann which may be a little more intuitive and is better for calculation. This is done via Riemann sums.

Riemann Sums

There is another way of getting at the integral due to Riemann which may be a little more intuitive and is better for calculation. This is done via Riemann sums.

To define the notion of a Riemann sum we need one more piece of data.

Riemann Sums

There is another way of getting at the integral due to Riemann which may be a little more intuitive and is better for calculation. This is done via Riemann sums.

To define the notion of a Riemann sum we need one more piece of data. Suppose that for each of the intervals I_j we are given a point $t_j \in I_j$. We will denote the collection of points t_j by t . The pair (P, t) is sometimes called a **tagged partition**.

Riemann Sums

There is another way of getting at the integral due to Riemann which may be a little more intuitive and is better for calculation. This is done via Riemann sums.

To define the notion of a Riemann sum we need one more piece of data. Suppose that for each of the intervals I_j we are given a point $t_j \in I_j$. We will denote the collection of points t_j by t . The pair (P, t) is sometimes called a **tagged partition**.

Definition: We define the **Riemann sum** associated to the function f , and the tagged partition (P, t) by

$$R(f, P, t) = \sum_{j=1}^n f(t_j)(x_j - x_{j-1}).$$

The norm of a partition

As must be clear, the Lower sum, Upper sum and Riemann sum all give approximations to the area between the lines $x = a$ and $x = b$ and between the curve $y = f(x)$ and the x -axis and

$$L(f, P) \leq R(f, P, t) \leq U(f, P).$$

The point is to make this statement quantitatively precise.

The norm of a partition

As must be clear, the Lower sum, Upper sum and Riemann sum all give approximations to the area between the lines $x = a$ and $x = b$ and between the curve $y = f(x)$ and the x -axis and

$$L(f, P) \leq R(f, P, t) \leq U(f, P).$$

The point is to make this statement quantitatively precise.

We define the **norm** of a partition P (denoted $\|P\|$) by

$$\|P\| = \max_j \{x_j - x_{j-1}\}, \quad 1 \leq j \leq n.$$

The norm of a partition

As must be clear, the Lower sum, Upper sum and Riemann sum all give approximations to the area between the lines $x = a$ and $x = b$ and between the curve $y = f(x)$ and the x -axis and

$$L(f, P) \leq R(f, P, t) \leq U(f, P).$$

The point is to make this statement quantitatively precise.

We define the **norm** of a partition P (denoted $\|P\|$) by

$$\|P\| = \max_j \{|x_j - x_{j-1}|\}, \quad 1 \leq j \leq n.$$

The norm gives some measure of the “size” of a partition, in particular, it allows us to say whether a partition is big or small.

The norm of a partition

As must be clear, the Lower sum, Upper sum and Riemann sum all give approximations to the area between the lines $x = a$ and $x = b$ and between the curve $y = f(x)$ and the x -axis and

$$L(f, P) \leq R(f, P, t) \leq U(f, P).$$

The point is to make this statement quantitatively precise.

We define the **norm** of a partition P (denoted $\|P\|$) by

$$\|P\| = \max_j \{|x_j - x_{j-1}|\}, \quad 1 \leq j \leq n.$$

The norm gives some measure of the “size” of a partition, in particular, it allows us to say whether a partition is big or small.

When the size of the partition is small, it means that **every interval in the partition is small**.

The Riemann integral

Definition 1: A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be **Riemann integrable** if for some $R \in \mathbb{R}$ and every $\epsilon > 0$ there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon,$$

whenever $\|P\| < \delta$. In this case R is called the **Riemann integral** of the function f on the interval $[a, b]$.

The Riemann integral

Definition 1: A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be **Riemann integrable** if for some $R \in \mathbb{R}$ and every $\epsilon > 0$ there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon,$$

whenever $\|P\| < \delta$. In this case R is called the **Riemann integral** of the function f on the interval $[a, b]$.

In other words, for all sufficiently “small” or “fine” partitions, the Riemann sums must be within ϵ of R .

The Riemann integral

Definition 1: A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be **Riemann integrable** if for some $R \in \mathbb{R}$ and every $\epsilon > 0$ there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon,$$

whenever $\|P\| < \delta$. In this case R is called the **Riemann integral** of the function f on the interval $[a, b]$.

In other words, for all sufficiently “small” or “fine” partitions, the Riemann sums must be within ϵ of R .

Notice, that as long as $\|P\|$ is small, **it doesn't matter exactly where the x_j 's or the t_j 's are in the interval $[a, b]$.**

The Riemann integral

Definition 1: A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be **Riemann integrable** if for some $R \in \mathbb{R}$ and every $\epsilon > 0$ there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon,$$

whenever $\|P\| < \delta$. In this case R is called the **Riemann integral** of the function f on the interval $[a, b]$.

In other words, for all sufficiently “small” or “fine” partitions, the Riemann sums must be within ϵ of R .

Notice, that as long as $\|P\|$ is small, **it doesn't matter exactly where the x_j 's or the t_j 's are in the interval $[a, b]$.**

Also notice that if P' is a refinement of P , then $\|P'\| \leq \|P\|$.

The Riemann integral continued

Intuitively, we can see that the smaller or finer the partition, the better the area under the curve is represented by the Riemann sum.

The Riemann integral continued

Intuitively, we can see that the smaller or finer the partition, the better the area under the curve is represented by the Riemann sum.

The reason that the Riemann integral is useful is because the definition we have given is actually equivalent to the following apparently weaker definition.

The Riemann integral continued

Intuitively, we can see that the smaller or finer the partition, the better the area under the curve is represented by the Riemann sum.

The reason that the Riemann integral is useful is because the definition we have given is actually equivalent to the following apparently weaker definition.

Definition 2: A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be **Riemann integrable** if for some $R \in \mathbb{R}$ and every $\epsilon > 0$ there exists a partition P such that for every tagged refinement of (P', t') of P with $\|P'\| \leq \delta$,

$$|R(f, P', t') - R| < \epsilon.$$

The Riemann integral continued

Intuitively, we can see that the smaller or finer the partition, the better the area under the curve is represented by the Riemann sum.

The reason that the Riemann integral is useful is because the definition we have given is actually equivalent to the following apparently weaker definition.

Definition 2: A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be **Riemann integrable** if for some $R \in \mathbb{R}$ and every $\epsilon > 0$ there exists a partition P such that for every tagged refinement of (P', t') of P with $\|P'\| \leq \delta$,

$$|R(f, P', t') - R| < \epsilon.$$

The nice thing about the above definition is that one only has to check that $|R(f, P', t') - R|$ is small for **refinements of a fixed partition, and not for all partitions.**

Note: the slide in class had P in one place, and (P, t) in another. I have corrected this.

Back to our example

Using Definition 2 of the Riemann integral it is easy to see that the function $f(x) = x$ is Riemann integrable.

Back to our example

Using Definition 2 of the Riemann integral it is easy to see that the function $f(x) = x$ is Riemann integrable.

Let $\epsilon > 0$ be arbitrary. For our fixed partition we take $P = P_n$ where $n > \frac{1}{2\epsilon}$ is some fixed number. The statements between the parentheses are true but not logically essential for our proof. [The Riemann sum is trapped between the upper and lower sums:

$$L(f, P_n) = \frac{1}{2} - \frac{1}{2n} \leq R(f, P_n, t) \leq U(f, P_n) = \frac{1}{2} + \frac{1}{2n},$$

for any choice t of points in the intervals I_j , that is,

Back to our example

Using Definition 2 of the Riemann integral it is easy to see that the function $f(x) = x$ is Riemann integrable.

Let $\epsilon > 0$ be arbitrary. For our fixed partition we take $P = P_n$ where $n > \frac{1}{2\epsilon}$ is some fixed number. The statements between the parentheses are true but not logically essential for our proof. [The Riemann sum is trapped between the upper and lower sums:

$$L(f, P_n) = \frac{1}{2} - \frac{1}{2n} \leq R(f, P_n, t) \leq U(f, P_n) = \frac{1}{2} + \frac{1}{2n},$$

for any choice t of points in the intervals I_j , that is,

$$|R(f, P_n, t) - \frac{1}{2}| < \frac{1}{2n} < \epsilon,$$

because of our choice of n .]

Back to our example

Using Definition 2 of the Riemann integral it is easy to see that the function $f(x) = x$ is Riemann integrable.

Let $\epsilon > 0$ be arbitrary. For our fixed partition we take $P = P_n$ where $n > \frac{1}{2\epsilon}$ is some fixed number. The statements between the parentheses are true but not logically essential for our proof. [The Riemann sum is trapped between the upper and lower sums:

$$L(f, P_n) = \frac{1}{2} - \frac{1}{2n} \leq R(f, P_n, t) \leq U(f, P_n) = \frac{1}{2} + \frac{1}{2n},$$

for any choice t of points in the intervals I_j , that is,

$$|R(f, P_n, t) - \frac{1}{2}| < \frac{1}{2n} < \epsilon,$$

because of our choice of n .] Moreover, if (P', t') is any refinement of P_n we have

$$L(f, P_n) \leq L(f, P') \leq R(f, P', t') \leq U(f, P') \leq U(f, P_n),$$

Back to our example

Using Definition 2 of the Riemann integral it is easy to see that the function $f(x) = x$ is Riemann integrable.

Let $\epsilon > 0$ be arbitrary. For our fixed partition we take $P = P_n$ where $n > \frac{1}{2\epsilon}$ is some fixed number. The statements between the parentheses are true but not logically essential for our proof. [The Riemann sum is trapped between the upper and lower sums:

$$L(f, P_n) = \frac{1}{2} - \frac{1}{2n} \leq R(f, P_n, t) \leq U(f, P_n) = \frac{1}{2} + \frac{1}{2n},$$

for any choice t of points in the intervals I_j , that is,

$$|R(f, P_n, t) - \frac{1}{2}| < \frac{1}{2n} < \epsilon,$$

because of our choice of n .] Moreover, if (P', t') is any refinement of P_n we have

$$L(f, P_n) \leq L(f, P') \leq R(f, P', t') \leq U(f, P') \leq U(f, P_n),$$

whence it follows that

$$|R(f, P', t') - \frac{1}{2}| < \epsilon.$$

The example continued

As the preceding example shows, Definition 2 of the Riemann integral is really easy to work with. Why do we then care about Definition 1 or the Darboux integral?

The example continued

As the preceding example shows, Definition 2 of the Riemann integral is really easy to work with. Why do we then care about Definition 1 or the Darboux integral?

The reason is that while Definition 2 is good for showing that a given function is Riemann integrable, the other definitions are often better for proving the *abstract properties* of integrals.

The example continued

As the preceding example shows, Definition 2 of the Riemann integral is really easy to work with. Why do we then care about Definition 1 or the Darboux integral?

The reason is that while Definition 2 is good for showing that a given function is Riemann integrable, the other definitions are often better for proving the *abstract properties* of integrals.

In fact, this will be clear in the tutorial exercises. You will see that sometimes the Darboux integral is better than the Riemann integral.

The example continued

As the preceding example shows, Definition 2 of the Riemann integral is really easy to work with. Why do we then care about Definition 1 or the Darboux integral?

The reason is that while Definition 2 is good for showing that a given function is Riemann integrable, the other definitions are often better for proving the *abstract properties* of integrals.

In fact, this will be clear in the tutorial exercises. You will see that sometimes the Darboux integral is better than the Riemann integral.

Before going any further we will formally state what we have already been referring to for several slides.

Comparison with the Darboux integral

Theorem 20: The Riemann integral (using either definition) exists if and only if the Darboux integral exists and in this case the two integrals are equal.

Comparison with the Darboux integral

Theorem 20: The Riemann integral (using either definition) exists if and only if the Darboux integral exists and in this case the two integrals are equal.

With this theorem in hand, we see that the function $f(x) = x$ is also Darboux integrable.

Comparison with the Darboux integral

Theorem 20: The Riemann integral (using either definition) exists if and only if the Darboux integral exists and in this case the two integrals are equal.

With this theorem in hand, we see that the function $f(x) = x$ is also Darboux integrable.

How does one prove Theorem 20? It is not too hard but it takes some work and is roundabout.

Comparison with the Darboux integral

Theorem 20: The Riemann integral (using either definition) exists if and only if the Darboux integral exists and in this case the two integrals are equal.

With this theorem in hand, we see that the function $f(x) = x$ is also Darboux integrable.

How does one prove Theorem 20? It is not too hard but it takes some work and is roundabout.

The easiest way is to proceed as follows. It is clear that if f is Riemann integrable in the sense of Definition 1, it is Riemann integrable in the sense of Definition 2.

Comparison with the Darboux integral

Theorem 20: The Riemann integral (using either definition) exists if and only if the Darboux integral exists and in this case the two integrals are equal.

With this theorem in hand, we see that the function $f(x) = x$ is also Darboux integrable.

How does one prove Theorem 20? It is not too hard but it takes some work and is roundabout.

The easiest way is to proceed as follows. It is clear that if f is Riemann integrable in the sense of Definition 1, it is Riemann integrable in the sense of Definition 2. Next, one shows that if f is Riemann integrable in the sense of Definition 2, then it is Darboux integrable.

Comparison with the Darboux integral

Theorem 20: The Riemann integral (using either definition) exists if and only if the Darboux integral exists and in this case the two integrals are equal.

With this theorem in hand, we see that the function $f(x) = x$ is also Darboux integrable.

How does one prove Theorem 20? It is not too hard but it takes some work and is roundabout.

The easiest way is to proceed as follows. It is clear that if f is Riemann integrable in the sense of Definition 1, it is Riemann integrable in the sense of Definition 2. Next, one shows that if f is Riemann integrable in the sense of Definition 2, then it is Darboux integrable. And finally, one can show that if the Darboux integral exists, then the Riemann integral exists in the sense of Definition 1. An interested student can try this as an exercise.

The main theorem for Riemann integration

From now on we will use any of the three definitions - the Darboux definition, Definition 1 and Definition 2 for the integral interchangeably and we will use only the words Riemann integral.

The main theorem for Riemann integration

From now on we will use any of the three definitions - the Darboux definition, Definition 1 and Definition 2 for the integral interchangeably and we will use only the words Riemann integral.

The main theorem of Riemann integration is the following:

The main theorem for Riemann integration

From now on we will use any of the three definitions - the Darboux definition, Definition 1 and Definition 2 for the integral interchangeably and we will use only the words Riemann integral.

The main theorem of Riemann integration is the following:

Theorem 21: Let $f : [a, b] \rightarrow \mathbb{R}$ be a function that is bounded, and continuous at all but finitely many points of $[a, b]$. Then f is Riemann integrable on $[a, b]$.

The main theorem for Riemann integration

From now on we will use any of the three definitions - the Darboux definition, Definition 1 and Definition 2 for the integral interchangeably and we will use only the words Riemann integral.

The main theorem of Riemann integration is the following:

Theorem 21: Let $f : [a, b] \rightarrow \mathbb{R}$ be a function that is bounded, and continuous at all but finitely many points of $[a, b]$. Then f is Riemann integrable on $[a, b]$.

In fact, one can allow even countably many discontinuities and the Theorem will remain true.

Exercise 1: Those of you who have an extra interest in the course should think about trying to prove both Theorem 21 and the extension to countably many discontinuities (**Warning:** there is one crucial fact about continuous functions that we have not covered that you will have to discover for yourself).