

## PH-107 (2017) Tutorial Sheet 12

\* Problems to be done in tutorial.

### **Drude, Sommerfeld Model, Band Energy**

**P101.** Find the velocity of copper wire whose cross-sectional area is 1 mm when the wire carries a current of 10 A. Assume that each copper atom contributes one electron to the electron gas.

**P102\*.** Using Drude model of an electron gas, calculate the relaxation time for Cu at room temperature given that its free electron density is  $8.5 \times 10^{28} \text{ m}^{-3}$  and its resistivity is  $1.5 \times 10^{-8} \Omega \text{m}$ .

**P103\*.** A copper wire of cross-sectional area  $3.3 \times 10^{-6} \text{ m}^2$  is carrying a current of 25 A. One meter length of this wire has a resistance of  $5.8 \times 10^{-3} \text{ ohms}$  at room temperature. Calculate the conductivity, average drift velocity of the electrons, electron mobility and mean free path of the electrons in Drude and Sommerfeld models at room temperature. [Take atomic weight of Cu as 63.5 and density as 8.94 g/cc]

**P104:** For Sodium the conductivity at 300 K is  $2.17 \times 10^7 \text{ ohm}^{-1} \text{m}^{-1}$  and the effective mass of the electron is 1.2 times the mass of the free electron. Calculate the relaxation time and the mean free path. Calculate the drift velocity of the electrons in an electric field of 100V/m. [Density of Na = 970 kg/m<sup>3</sup>, Atomic weight 23]

**P105\*:** Use the equation  $m(dv/dt + v/\tau) = -eE$  for the electron drift velocity  $v$ , show that the conductivity at frequency  $\omega$  is

$$\sigma(\omega) = \sigma(0) \left[ \frac{1 + i\omega\tau}{1 + (\omega\tau)^2} \right]$$

where  $\sigma(0) = ne^2\tau / m$

[Problem from book by Charles Kittel 6.6]

**P106:** Using the data given and any other constants, evaluate the Fermi energy of the alkali metals.

	Li	Na	K	Rb	Cs
<b>Density (g/cc)</b>	0.534	0.971	0.860	1.530	1.870
<b>Atomic weight</b>	6.939	22.99	39.102	85.47	132.905

**P107\*:** The Fermi energy of Cu is 7.04 eV. Calculate the velocity and de Broglie wavelength of electrons at the Fermi energy of Cu. Can these electrons be diffracted by a crystal?

**P108\*:** The energy of an electron in a band of 1-dimensional solid, as a function of wavenumber  $k$ , is given by

$$E(k) = 2 - \cos(ka) \quad \text{for } k \in \left[-\frac{\pi}{a}, \frac{\pi}{a}\right]$$

**(a)** Plot  $E$  versus  $k$ .

**(b)** Find the speed ( $v$ ) and effective mass ( $m^*$ ) of the electron as a function of  $k$ .

**(c)** Plot  $m^*$  versus  $k$ .