

Sequential continuity

Ravi Raghunathan

Department of Mathematics

Autumn 2017, IIT Bombay, Mumbai

Sequential continuity

The definition

Let us recall the definition of sequential continuity:

The definition

Let us recall the definition of sequential continuity:

Definition: A function $f : (a, b) \rightarrow \mathbb{R}$ is said to be sequentially continuous at a point $c \in (a, b)$ if for **every** sequence x_n in (a, b) such that $\lim_{n \rightarrow \infty} x_n = c$, $\lim_{n \rightarrow \infty} f(x_n) = f(c)$.

The definition

Let us recall the definition of sequential continuity:

Definition: A function $f : (a, b) \rightarrow \mathbb{R}$ is said to be sequentially continuous at a point $c \in (a, b)$ if for **every** sequence x_n in (a, b) such that $\lim_{n \rightarrow \infty} x_n = c$, $\lim_{n \rightarrow \infty} f(x_n) = f(c)$.

Theorem: A function $f : (a, b) \rightarrow \mathbb{R}$ is continuous at c if and only if it is sequentially continuous at c .

The definition

Let us recall the definition of sequential continuity:

Definition: A function $f : (a, b) \rightarrow \mathbb{R}$ is said to be sequentially continuous at a point $c \in (a, b)$ if for **every** sequence x_n in (a, b) such that $\lim_{n \rightarrow \infty} x_n = c$, $\lim_{n \rightarrow \infty} f(x_n) = f(c)$.

Theorem: A function $f : (a, b) \rightarrow \mathbb{R}$ is continuous at c if and only if it is sequentially continuous at c .

The theorem is often useful in its contrapositive form, that is, to show that a function is discontinuous at a point c it is enough to show that it is not sequentially continuous at the point c , i.e., that **there is at least one sequence** x_n such that $\lim_{n \rightarrow \infty} x_n = c$, but $\lim_{n \rightarrow \infty} f(x_n) \neq f(c)$.

Continuity implies sequential continuity

Suppose f is continuous at c . Let $\epsilon > 0$ be given. Because f is continuous at c , there exists $\delta > 0$ such that

$$|f(x) - f(c)| < \epsilon \quad \text{whenever} \quad 0 < |x - c| < \delta. \quad (1)$$

Continuity implies sequential continuity

Suppose f is continuous at c . Let $\epsilon > 0$ be given. Because f is continuous at c , there exists $\delta > 0$ such that

$$|f(x) - f(c)| < \epsilon \quad \text{whenever} \quad 0 < |x - c| < \delta. \quad (1)$$

Since $\lim_{n \rightarrow \infty} x_n = c$, for the $\delta > 0$ chosen above, there exists $N \in \mathbb{N}$ such that $|x_n - c| < \delta$, whenever $n > N$.

Continuity implies sequential continuity

Suppose f is continuous at c . Let $\epsilon > 0$ be given. Because f is continuous at c , there exists $\delta > 0$ such that

$$|f(x) - f(c)| < \epsilon \quad \text{whenever} \quad 0 < |x - c| < \delta. \quad (1)$$

Since $\lim_{n \rightarrow \infty} x_n = c$, for the $\delta > 0$ chosen above, there exists $N \in \mathbb{N}$ such that $|x_n - c| < \delta$, whenever $n > N$.

By equation (1) above, it follows that $|f(x_n) - f(c)| < \epsilon$ for all $n > N$. This shows that $\lim_{n \rightarrow \infty} f(x_n) = f(c)$. □

Sequential Continuity implies Continuity

This is the slightly harder direction. We will prove it by contradiction.

Sequential Continuity implies Continuity

This is the slightly harder direction. We will prove it by contradiction.

Suppose f is sequentially continuous at c but not continuous at c .

Sequential Continuity implies Continuity

This is the slightly harder direction. We will prove it by contradiction.

Suppose f is sequentially continuous at c but not continuous at c .

Since f is not continuous, there exists **some** $\epsilon > 0$, such that for **any** $\delta > 0$ there is a point x such that

$$0 < |x - c| < \delta \quad \text{and} \quad |f(x) - f(c)| \geq \epsilon.$$

Sequential Continuity implies Continuity

This is the slightly harder direction. We will prove it by contradiction.

Suppose f is sequentially continuous at c but not continuous at c .

Since f is not continuous, there exists **some** $\epsilon > 0$, such that for **any** $\delta > 0$ there is a point x such that

$$0 < |x - c| < \delta \quad \text{and} \quad |f(x) - f(c)| \geq \epsilon.$$

Fix ϵ as above. For each $n \in \mathbb{N}$, let $\delta = 1/n$. Then there exists x_n such that $0 < |x_n - c| < 1/n$ and $|f(x_n) - f(c)| \geq \epsilon$. Clearly $\lim_{n \rightarrow \infty} x_n = c$, but $\lim_{n \rightarrow \infty} f(x_n) \neq f(c)$.

Sequential Continuity implies Continuity

This is the slightly harder direction. We will prove it by contradiction.

Suppose f is sequentially continuous at c but not continuous at c .

Since f is not continuous, there exists **some** $\epsilon > 0$, such that for **any** $\delta > 0$ there is a point x such that

$$0 < |x - c| < \delta \quad \text{and} \quad |f(x) - f(c)| \geq \epsilon.$$

Fix ϵ as above. For each $n \in \mathbb{N}$, let $\delta = 1/n$. Then there exists x_n such that $0 < |x_n - c| < 1/n$ and $|f(x_n) - f(c)| \geq \epsilon$. Clearly $\lim_{n \rightarrow \infty} x_n = c$, but $\lim_{n \rightarrow \infty} f(x_n) \neq f(c)$.

This contradicts the sequential continuity of f at c . Hence our assumption that f is not continuous at c must have been false. \square

An everywhere discontinuous function

Let us return to the Exercise 3 of the Optional Exercises. Consider the function

$$f(x) = \begin{cases} 0 & \text{if } x \text{ is rational;} \\ 1 & \text{if } x \text{ is irrational.} \end{cases}$$

We will show that there is no point at which f is continuous. We will use the following two facts (which are intuitively obvious).

An everywhere discontinuous function

Let us return to the Exercise 3 of the Optional Exercises. Consider the function

$$f(x) = \begin{cases} 0 & \text{if } x \text{ is rational;} \\ 1 & \text{if } x \text{ is irrational.} \end{cases}$$

We will show that there is no point at which f is continuous. We will use the following two facts (which are intuitively obvious).

Fact 1: We can find a rational number between any two real numbers.

An everywhere discontinuous function

Let us return to the Exercise 3 of the Optional Exercises. Consider the function

$$f(x) = \begin{cases} 0 & \text{if } x \text{ is rational;} \\ 1 & \text{if } x \text{ is irrational.} \end{cases}$$

We will show that there is no point at which f is continuous. We will use the following two facts (which are intuitively obvious).

Fact 1: We can find a rational number between any two real numbers.

Fact 2: We can find an irrational number between any two real numbers.

We will prove both these facts a little later. For now we will assume that they are true.

Suppose that c is rational. Then, by Fact 1, in every interval $(c, c + 1/n)$, $n \in \mathbb{N}$, we can find an irrational number, say x_n . Clearly $\lim_{n \rightarrow \infty} x_n = c$. Since $f(x_n) = 1$ for all x_n , with $n \in \mathbb{N}$ and $f(c) = 0$, $|f(x_n) - f(c)| = 1$.

Suppose that c is rational. Then, by Fact 1, in every interval $(c, c + 1/n)$, $n \in \mathbb{N}$, we can find an irrational number, say x_n . Clearly $\lim_{n \rightarrow \infty} x_n = c$. Since $f(x_n) = 1$ for all x_n , with $n \in \mathbb{N}$ and $f(c) = 0$, $|f(x_n) - f(c)| = 1$.

Thus, if $\epsilon = 1$ we see that there is no $\delta > 0$ such that $0 < |x - c| < \delta$ implies $|f(x) - f(c)| < \epsilon$. This shows that f is not continuous at c .

Suppose that c is rational. Then, by Fact 1, in every interval $(c, c + 1/n)$, $n \in \mathbb{N}$, we can find an irrational number, say x_n . Clearly $\lim_{n \rightarrow \infty} x_n = c$. Since $f(x_n) = 1$ for all x_n , with $n \in \mathbb{N}$ and $f(c) = 0$, $|f(x_n) - f(c)| = 1$.

Thus, if $\epsilon = 1$ we see that there is no $\delta > 0$ such that $0 < |x - c| < \delta$ implies $|f(x) - f(c)| < \epsilon$. This shows that f is not continuous at c .

If c is irrational, we use Fact 2 to show (in the same way as above) that f is not continuous at c . □

The proofs of Facts 1 and 2

We first prove Fact 1.

The proofs of Facts 1 and 2

We first prove Fact 1.

Suppose x and y are two real numbers with $x < y$. Choose n such that $1/n < y - x$.

The proofs of Facts 1 and 2

We first prove Fact 1.

Suppose x and y are two real numbers with $x < y$. Choose n such that $1/n < y - x$.

Consider the numbers of the form k/n with $k \in \mathbb{Z}$. There will be some $m \in \mathbb{Z}$ such that $m/n \leq x$ but $(m+1)/n > x$.

Clearly

$$\frac{m+1}{n} = \frac{m}{n} + \frac{1}{n} \leq x + \frac{1}{n} < y.$$

So $(m+1)/n$ is the desired rational number.

The proofs of Facts 1 and 2

We first prove Fact 1.

Suppose x and y are two real numbers with $x < y$. Choose n such that $1/n < y - x$.

Consider the numbers of the form k/n with $k \in \mathbb{Z}$. There will be some $m \in \mathbb{Z}$ such that $m/n \leq x$ but $(m+1)/n > x$.

Clearly

$$\frac{m+1}{n} = \frac{m}{n} + \frac{1}{n} \leq x + \frac{1}{n} < y.$$

So $(m+1)/n$ is the desired rational number.

To prove Fact 2, replace $1/n$ in the argument above by $\sqrt{2}/n$!