

MA-106 Linear Algebra

H. Ananthnarayan



Department of Mathematics
Indian Institute of Technology Bombay
Powai, Mumbai - 76

8th February 2018
D1 - Lecture 16

Random Attendance

1	170050001	Piyush Dilip Tibarewal	
2	170050013	Shrutesh Tapesh Patil	
3	170050030	Anurag Arun Kedia	
4	170050034	Rajdeep Yadav	
5	170050006	Kalsaria Sagar Pareshbhai	
6	170050042	Ajay	
7	170050066	Manas Shukla	
8	170050096	Ramolla Nikhil Reddy	
9	170070002	Thorve Rajesh Bhaskar	
10	170070033	Anurag Fogawat	Absent
11	170070049	Modhugu Rineeth	Absent
12	170070053	Ponnada Prem Sekhar	
13	17D070010	Shubham Anand Jain	
14	17D070013	Paras Vijay Bodake	Absent
15	17D070015	Nikhil Arvind Bhaladhare	Absent
16	17D070062	Jeetika	

Recall: Formula for Determinant

- **Using Permutations:** For $n \times n$ matrix $A = (a_{ij})$,

$$\det(A) = \sum_{\text{all permutations } P} (a_{1\alpha_1} \dots a_{n\alpha_n}) \det(P).$$

Here a permutation (i_1, i_2, \dots, i_n) of $(1, 2, \dots, n)$ corresponds to the

permutation matrix $P = \begin{bmatrix} e_{i_1}^T \\ \vdots \\ e_{i_n}^T \end{bmatrix}.$

- **Using Cofactors:** Let C_{1j} be the coefficient of a_{1j} in the expansion

$$\det(A) = \sum_{\text{all permutations } P} (a_{1\alpha_1} \dots a_{n\alpha_n}) \det(P)$$

- Then $\boxed{\det(A) = a_{11}C_{11} + a_{12}C_{12} + \dots + a_{1n}C_{1n}.}$

Note: $C_{1j} = (-1)^{1+j} \det(M_{1j})$, where M_{1j} is obtained from A by deleting the 1st row and j^{th} column.

Recall: Formula for Determinant

- **Expansion along i th row:** If C_{ij} is the coefficient of a_{ij} in the formula of $\det(A)$, then $\det(A) = a_{i1} C_{i1} + \dots + a_{in} C_{in}$.

Note: $C_{ij} = (-1)^{i+j} \det(M_{ij})$, where M_{ij} is obtained from A by deleting i -th row and j -th column.

Idea: Using $i - 1$ row exchanges, make A_{i*} the first row.

- **Expansion along j th column:** $C_{ij}(A^T) = C_{ji}(A)$ and $\det(A) = \det(A^T) \Rightarrow \det(A) = a_{1j} C_{1j}(A) + \dots + a_{nj} C_{nj}(A)$.

Computing determinants: Exercise

Example: Let $F_n = \begin{vmatrix} 1 & -1 & & & \\ 1 & 1 & -1 & & \\ & \ddots & \ddots & \ddots & \\ & & 1 & 1 & -1 \\ & & & 1 & 1 \end{vmatrix}$ be a $(1, 1, -1)$ tri-diagonal $n \times n$ matrix. Expanding along the first row, we get

$$F_n = F_{n-1} + (-1)^{1+2}(-1) \begin{vmatrix} & & & & \\ & 1 & -1 & & \\ & 1 & 1 & -1 & \\ & & \ddots & \ddots & \ddots \\ & & & 1 & 1 & -1 \\ & & & & 1 & 1 \end{vmatrix} = F_{n-1} + F_{n-2},$$

by expanding along first column.

Since $F_1 = \dots$, $F_2 = \dots$, the sequence F_n is $\dots, \dots, \dots, \dots$

Computing determinants: Examples

Find the determinants for the following examples

• $A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 \end{bmatrix}$. Expand along 1st column.

• $B = \begin{bmatrix} 3 & 2 & -1 \\ 0 & 2 & 0 \\ 1 & 1 & 1 \end{bmatrix}$. Expand along 2nd row.

• $E = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}$. Find the cofactor matrix C . Compute EC^T .

$$C = \begin{bmatrix} 3 & 2 & 1 \\ 2 & 4 & 2 \\ 1 & 2 & 3 \end{bmatrix} \quad EC^T = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix} \begin{bmatrix} 3 & 2 & 1 \\ 2 & 4 & 2 \\ 1 & 2 & 3 \end{bmatrix} = \begin{bmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix}$$

Note: $\det(E) = 4$ and $EC^T = 4I \Rightarrow E^{-1} = \frac{1}{\det(E)} C^T$.

Applications: 1. Computing A^{-1}

If $C = (C_{ij})$: cofactor matrix of A , then $A C^T = \det(A) I$ i.e.,

$$\begin{bmatrix} a_{11} & \dots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{n1} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} C_{11} & \dots & C_{n1} \\ \vdots & \ddots & \vdots \\ C_{1n} & \dots & C_{nn} \end{bmatrix} = \begin{bmatrix} \det(A) & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \det(A) \end{bmatrix}$$

Proof. We have seen that $a_{i1}C_{i1} + \dots + a_{in}C_{in} = \det(A)$. Now

$$a_{11}C_{21} + a_{12}C_{22} + \dots + a_{1n}C_{2n} = \det \begin{bmatrix} a_{11} & \dots & a_{1n} \\ a_{11} & \dots & a_{1n} \\ a_{31} & \dots & a_{3n} \\ \vdots & & \vdots \\ a_{n1} & \dots & a_{nn} \end{bmatrix} = 0.$$

Similarly, if $i \neq j$, then $a_{i1}C_{j1} + a_{i2}C_{j2} + \dots + a_{in}C_{jn} = 0$. □

Remark. If A is invertible, then $A^{-1} = \frac{1}{\det(A)} C^T$.

For $n \geq 4$, this is *not* a good formula to find A^{-1} .

Use elimination to find A^{-1} for $n \geq 4$.

Reading Slide - Exercise

Find the inverse of $A = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 3 & 0 \\ 0 & 4 & 1 \end{bmatrix}$.

$$\det(A) = \begin{vmatrix} 3 & 0 \\ 4 & 1 \end{vmatrix} + (-1)^{1+2}(2) \begin{vmatrix} 0 & 0 \\ 0 & 1 \end{vmatrix} = 3$$

$$C_{11} = 3, C_{12} = 0, C_{13} = 0$$

$$C_{21} = -2, C_{22} = 1, C_{23} = -4$$

$$C_{31} = 0, C_{32} = 0, C_{33} = 3.$$

$$\text{Hence } A^{-1} = \frac{1}{\det(A)} C^T = \frac{1}{3} \begin{bmatrix} 3 & -2 & 0 \\ 0 & 1 & 0 \\ 0 & -4 & 3 \end{bmatrix}$$

Check. $AA^{-1} = I$.

Applications: 2. Solving $Ax = b$

Cramer's rule: If A is invertible, the $Ax = b$ has a unique solution.

$$x = A^{-1}b = \frac{1}{\det(A)} C^T b = \frac{1}{\det(A)} \begin{bmatrix} C_{11} & \cdots & C_{n1} \\ \vdots & & \vdots \\ C_{1n} & \cdots & C_{nn} \end{bmatrix} \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$$

$$\text{Hence } x_j = \frac{1}{\det(A)} (b_1 C_{1j} + b_2 C_{2j} + \cdots + b_n C_{nj}) = \frac{1}{\det(A)} \det(B_j),$$

where B_j is obtained by replacing j^{th} column of A by b , and $\det(B_j)$ is computed along the j^{th} column.

Extra Reading - Applications: 3. A Formula for Pivots

Observation: If row exchanges are not required, then the first k pivots are determined by the top-left $k \times k$ submatrices \tilde{A}_k of A .

Example. If $A = [a_{ij}]_{3 \times 3}$, then $\tilde{A}_1 = (a_{11})$, $\tilde{A}_2 = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ and $\tilde{A}_3 = A$.

Assume the pivots are d_1, \dots, d_n , obtained without row exchange. Then

- $\det(\tilde{A}_1) = a_{11} = d_1$
- $\det(\tilde{A}_2) = d_1 d_2 = \det(A_1) d_2$
- $\det(\tilde{A}_3) = d_1 d_2 d_3 = \det(A_2) d_3$ etc.,
- If $\det(\tilde{A}_k) = 0$, then we need a row exchange in elimination.
- Otherwise the k -th pivot is $d_k = \det(\tilde{A}_k) / \det(\tilde{A}_{k-1})$

Extra Reading - Example

Example. Find if row exchange is required in Gauss Elimination of

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 4 & 4 \\ 1 & 4 & 9 \end{pmatrix} \text{ and find the pivots.}$$

Solution. $\tilde{A}_1 = (1)$, $\tilde{A}_2 = \begin{pmatrix} 1 & 1 \\ 1 & 4 \end{pmatrix}$, $\tilde{A}_3 = A$

$$\det(\tilde{A}_1) = 1, \quad \det(\tilde{A}_2) = 3$$

$$\det(\tilde{A}_3) = \begin{vmatrix} 1 & 1 & 1 \\ 0 & 3 & 3 \\ 0 & 3 & 8 \end{vmatrix} = \begin{vmatrix} 3 & 3 \\ 3 & 8 \end{vmatrix} = 15$$

Therefore row exchange is not required in elimination and the pivots are $d_1 = 1$, $d_2 = 3/1 = 3$, $d_3 = 15/3 = 5$.

Verify this directly by elimination!

Summary: Determinants

Let A and B $n \times n$, and t a scalar.

- $\det(A + B) \neq \det(A) + \det(B)$, and $\det(tA) = t^n \det(A)$.
- $\det(AB) = \det(A)\det(B)$.
- $\det(A) = \det(A^T)$.
- If A is orthogonal, i.e., $AA^T = I$, then $\det(A) =$
- If $A = [a_{ij}]$ is triangular, then $\det(A) =$
- A is invertible $\Leftrightarrow \det(A) \neq 0$. If this happens, then $\det(A^{-1}) =$
- If A and B are similar, i.e., $B = S^{-1}AS$ for an invertible matrix S , then $\det(B) =$
- If A is invertible, and d_1, \dots, d_n are the pivots of A , then $\det(A) =$

Eigenvalues and Eigenvectors: Motivation

- Solve for the differential equation for u : $du/dt = 3u$.

The solution is $u(t) = c e^{3t}$, $c \in \mathbb{R}$

With initial condition $u(0) = 2$, the solution is $u(t) = 2e^{3t}$.

- Consider the system of linear 1st order differential equations (ODE) with constant coefficients:

$$du_1/dt = 4u_1 - 5u_2,$$

$$du_2/dt = 2u_1 - 3u_2,$$

How does one find the solution?

- Write the system in matrix form $du/dt = Au$,

where $u(t) = \begin{pmatrix} u_1(t) \\ u_2(t) \end{pmatrix}$, $A = \begin{pmatrix} 4 & -5 \\ 2 & -3 \end{pmatrix}$.

- Assuming the solution is $u(t) = \begin{pmatrix} u_1(t) \\ u_2(t) \end{pmatrix} = e^{\lambda t} v = \begin{pmatrix} e^{\lambda t} x \\ e^{\lambda t} y \end{pmatrix}$, where

$v = \begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2$, we need to find λ and v .

Eigenvalues and Eigenvectors: Definition

We have $u_1' = 4u_1 - 5u_2$, $u_2' = 2u_1 - 3u_2$, where
 $u_1(t) = e^{\lambda t} x$, $u_2(t) = e^{\lambda t} y$

$$\lambda e^{\lambda t} x = 4e^{\lambda t} x - 5e^{\lambda t} y,$$

$$\lambda e^{\lambda t} y = 2e^{\lambda t} x - 3e^{\lambda t} y.$$

Cancelling $e^{\lambda t}$, we get

Eigenvalue problem: Find λ and $v = (x, y)^T$ satisfying

$$4x - 5y = \lambda x,$$

$$2x - 3y = \lambda y.$$

In the matrix form, it is $A v = \lambda v$.

This equation has two unknowns, λ and v .

If there exists a λ such that $A v = \lambda v$ has a non-zero solution v , then λ is called an **eigenvalue** of A and all *nonzero* v satisfying $A v = \lambda v$ are called **eigenvectors** of A associated to λ .

Q: Given A $n \times n$, how does one find its eigenvalues and eigenvectors?

Eigenvalues and Eigenvectors: Solving $Ax = \lambda x$

- Write $Av = \lambda v$ as $(A - \lambda I)v = 0$.
- λ is an eigenvalue of A
 - \Leftrightarrow there is a nonzero v in the nullspace of $A - \lambda I$
 - $\Leftrightarrow N(A - \lambda I) \neq 0$, i.e., $\dim(N(A - \lambda I)) \geq 1$,
 - $\Leftrightarrow A - \lambda I$ is singular
 - $\Leftrightarrow \det(A - \lambda I) = 0$.
- $\det(A - \lambda I)$ is a polynomial in the variable λ of degree n . Hence it has **at most** n roots $\Rightarrow A$ has at most n eigenvalues.
- $\det(A - \lambda I)$ is called the **characteristic polynomial** of A .
- If λ is an eigenvalue of A , then the nullspace of $A - \lambda I$ is called the **eigenspace** of A associated to eigenvalue λ .
- $\lambda = 0$ is an eigenvalue of $A \Leftrightarrow \det(A) = 0 \Leftrightarrow A$ is singular.

Eigenvalues and Eigenvectors: Example

To summarise: An eigenvalue of A is a root of its characteristic polynomial, and any non-zero vector in the corresponding eigenspace is an associated eigenvector.

Recall: The ODE system we want to solve is

$$u_1' = 4u_1 - 5u_2, \quad u_1(0) = 8, \quad u_2' = 2u_1 - 3u_2, \quad u_2(0) = 5.$$

The solutions are $u_1(t) = e^{\lambda t} x$, $u_2(t) = e^{\lambda t} y$, where $(x, y)^T$ is a solution of:

$$\begin{pmatrix} 4 & -5 \\ 2 & -3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \lambda \begin{pmatrix} x \\ y \end{pmatrix} \quad (Av = \lambda v)$$

The characteristic polynomial of A is

$$\begin{aligned} \det(A - \lambda I) &= \det \begin{pmatrix} 4 - \lambda & -5 \\ 2 & -3 - \lambda \end{pmatrix} \\ &= (4 - \lambda)(-3 - \lambda) + 10 \\ &= \lambda^2 - \lambda - 2 = (\lambda + 1)(\lambda - 2) \end{aligned}$$

The eigenvalues of A are $\lambda_1 = -1, \lambda_2 = 2$.

Eigenvalues and Eigenvectors: Example

Eigenvectors v_1 and v_2 associated to $\lambda_1 = -1$ and $\lambda_2 = 2$ respectively, are in:

$$N(A - \lambda_1 I) = N(A + I), \text{ and } N(A - \lambda_2 I) = N(A - 2I).$$

Solving $(A + I)v = 0$, i.e., $\begin{pmatrix} 5 & -5 \\ 2 & -2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = 0$, we get

$N(A + I) = \left\{ \begin{pmatrix} y \\ y \end{pmatrix} \mid y \in \mathbb{R} \right\}$ and $v_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ is an eigenvector associated to $\lambda_1 = -1$.

Similarly, solving $(A - 2I)v = 0$ gives $N(A - 2I) = \left\{ \begin{pmatrix} 5y \\ y \end{pmatrix} \mid y \in \mathbb{R} \right\}$.

In particular, $v_2 = \begin{pmatrix} 5 \\ 2 \end{pmatrix}$ is an eigenvector associated to $\lambda_2 = 2$.

Thus, the system $du/dt = Au$ has two special solutions

$$e^{-t}v_1 \text{ and } e^{2t}v_2.$$

Reading Slide - Complete Solution to ODE

Note: When two functions satisfy $du/dt = Au$, then so do their linear combinations.

Complete solution: $u(t) = c_1 e^{-t} v_1 + c_2 e^{2t} v_2$,

$$\text{i.e. } \begin{pmatrix} u_1(t) \\ u_2(t) \end{pmatrix} = c_1 e^{-t} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + c_2 e^{2t} \begin{pmatrix} 5 \\ 2 \end{pmatrix}.$$

$$\text{i.e. } u_1(t) = c_1 e^{-t} + 5c_2 e^{2t}, \quad u_2(t) = c_1 e^{-t} + 2c_2 e^{2t}.$$

If we put initial conditions (IC) $u_1(0) = 8$ and $u_2(0) = 5$, then

$$c_1 + 5c_2 = 8, \quad c_1 + 2c_2 = 5 \Rightarrow c_1 = 3, \quad c_2 = 1.$$

Hence the solution of the original ODE system with the given IC is

$$u_1(t) = 3e^{-t} + 5e^{2t}, \quad u_2(t) = 3e^{-t} + 2e^{2t}.$$

Examples

In some cases it is easy to find the eigenvalues.

Example: $A = \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix}$ is diagonal. Characteristic polynomial $(3 - \lambda)(2 - \lambda)$.

Eigenvalues: $\lambda_1 = 3, \lambda_2 = 2$.

Eigenvectors: $(A - 3I)v_1 = 0 \Rightarrow Av_1 = 3v_1 \Rightarrow v_1 = e_1$

Similarly, an eigenvector associated to λ_2 is $v_2 = e_2$

Further, \mathbb{R}^2 has a basis consisting of eigenvectors of A : $\{e_1, e_2\}$.

General case: If A is a diagonal matrix with diagonal entries $\lambda_1, \dots, \lambda_n$, then

Eigenvalues: $\lambda_1, \dots, \lambda_n$ Eigenvectors: e_1, \dots, e_n ,

which form a basis for \mathbb{R}^n .

Examples

Example: Projection onto the line $x = y$: $P = \begin{pmatrix} 1/2 & 1/2 \\ 1/2 & 1/2 \end{pmatrix}$.

$v_1 = \begin{pmatrix} 1 & 1 \end{pmatrix}^T$ projects onto itself $\Rightarrow \lambda_1 = 1$ with eigenvector v_1 .

$v_2 = \begin{pmatrix} 1 & -1 \end{pmatrix}^T \mapsto 0 \Rightarrow \lambda_2 = 0$ with eigenvector v_2 .

Further, $\{v_1, v_2\}$ is a basis of \mathbb{R}^2 .

Q: Do a collection of eigenvectors always form a basis of \mathbb{R}^n ?