

MA-106 Linear Algebra

H. Ananthnarayan



Department of Mathematics
Indian Institute of Technology Bombay
Powai, Mumbai - 76

13th February 2018
D1 - Lecture 18

Random Attendance: 13th Feb

1	170050010	Arjit Jain	Absent
2	170050023	Akshay Yadav	
3	170050026	Tushar Gautam	
4	170050054	Chaw Akametta Enling	
5	170050068	Killari Ramprasad	
6	170050069	Gulla Niranjan	
7	170050071	Yash Raj	
8	170050093	Somavarapu Tarun	
9	170050097	Mannem Sai Varshitha	
10	170050107	Sarvesh Mehtani	
11	170070014	Shriram Girish Lokhande	Absent
12	170070019	Titus Chakraborty	
13	170070029	Chandra Shekhar	
14	17D070001	Rathod Harekrissna Upendra	
15	17D070014	Nakrani Prajval Sushil	
16	17D070063	Prachi Goel	

Diagonalization: Example

Example: $A = \begin{pmatrix} 1 & 5 & 6 \\ 0 & 2 & -4 \\ 0 & 0 & 3 \end{pmatrix}$ is triangular.

Eigenvalues: $\lambda_1 = 1, \lambda_2 = 2, \lambda_3 = 3$.

Eigenvectors: $v_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, v_2 = \begin{pmatrix} 5 \\ 1 \\ 0 \end{pmatrix}, v_3 = \begin{pmatrix} -7 \\ -4 \\ 1 \end{pmatrix}$.

Further, $\mathcal{B} = \{v_1, v_2, v_3\}$ is a basis of \mathbb{R}^3 . Hence $P = (v_1 \ v_2 \ v_3)$ is invertible, and $AP = (Av_1 \ Av_2 \ Av_3) = (v_1 \ 2v_2 \ 3v_3) = P\Lambda$,

where $\Lambda = \begin{pmatrix} 1 & & \\ & 2 & \\ & & 3 \end{pmatrix}$. Thus $P^{-1}AP = \Lambda$, i.e., A is diagonalizable.

Diagonalization of a Matrix

Question: What is the advantage of a basis of \mathbb{R}^n consisting of eigenvectors?

Let A be an $n \times n$ matrix with n eigenvectors v_1, \dots, v_n , associated to eigenvalues $\lambda_1, \dots, \lambda_n$. If $\mathcal{B} = \{v_1, \dots, v_n\}$ is a basis of \mathbb{R}^n , then the matrix $P = (v_1 \ \cdots \ v_n)$ is invertible.

Moreover, $AP = A(v_1 \ \cdots \ v_n) = (Av_1 \ \cdots \ Av_n)$

$$= (\lambda_1 v_1 \ \cdots \ \lambda_n v_n) = P\Lambda, \text{ where } \Lambda = \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix}.$$

Therefore $P^{-1}AP = \Lambda$, i.e., A is similar to a diagonal matrix.

Thus: Eigenvectors diagonalize a matrix

Caution: $\Lambda P \neq P\Lambda$ in general.

When is A Diagonalizable?

Thus, we have proved: If an $n \times n$ matrix A has n linearly independent eigenvectors v_1, \dots, v_n , then A is diagonalizable.

Moreover, if $\lambda_1, \dots, \lambda_n$ are the corresponding eigenvalues, then $P^{-1}AP = \Lambda$, where the diagonalizing matrix is $P = (v_1 \ \cdots \ v_n)$, and

the diagonal matrix is $\Lambda = \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix}$, i.e., $P^{-1}AP = \Lambda$, where

The diagonal entries of Λ are eigenvalues of A and

The columns of P are corresponding eigenvectors of A .

Exercise: With the above notation, if $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is defined by $T(v) = Av$, and $\mathcal{B} = \{v_1, \dots, v_n\}$, then $[T]_{\mathcal{B}}^{\mathcal{B}} = \dots$.

Note: P need not be unique, e.g., replace v_1 by $2v_1$, etc.

When is A Diagonalizable?

Q: When does A have n linearly independent eigenvectors?

- If v_1, \dots, v_r are eigenvectors of A associated to **distinct** eigenvalues $\lambda_1, \dots, \lambda_r$, then v_1, \dots, v_r are linearly independent.

Proof. Suppose v_1, \dots, v_r are linearly dependent. Choose a linear relation involving minimum number of v_i 's, say

$$(1) \quad a_1 v_1 + \dots + a_t v_t = 0. \quad (1 < t \leq r, t \text{ is minimal, } a_i \neq 0)$$

Apply A to get $a_1 \lambda_1 v_1 + \dots + a_t \lambda_t v_t = 0 \quad (2)$

$$\lambda_1(1) - (2) \text{ gives } a_2(\lambda_1 - \lambda_2)v_2 + \dots + a_t(\lambda_1 - \lambda_t)v_t = 0,$$

which contradicts the minimality of t . □

- If A has n distinct eigenvalues, then A is diagonalizable.

Proof. If v_1, \dots, v_n are eigenvectors associated to distinct eigenvalues $\lambda_1, \dots, \lambda_n$, then $\{v_1, \dots, v_n\}$ is linearly independent.

Then $P = (v_1 \ \dots \ v_n)$ is invertible, and $P^{-1}AP = \Lambda$ as earlier. Hence A is diagonalizable. □

When is A Diagonalizable?

Theorem A is diagonalizable $\Leftrightarrow A$ has n linearly independent eigenvectors. In particular, \mathbb{R}^n has a basis consisting of eigenvectors of A .

Proof. (\Leftarrow): Done! To prove (\Rightarrow), assume $P = (v_1 \ \cdots \ v_n)$ is an invertible matrix such that $P^{-1}AP = \Lambda = \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix}$. Then

$AP = P\Lambda$, i.e. $(Av_1 \ \cdots \ Av_n) = (\lambda_1 v_1 \ \cdots \ \lambda_n v_n)$. Therefore

v_1, \dots, v_n are eigenvectors of A . They are linearly independent since P is invertible. \square

Question: Is every matrix is diagonalizable? **A:** No.

Examples: $Q = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ no eigenvalues (over \mathbb{R})!

$P = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ not enough eigenvectors!

Summary: Diagonalizability

Let A be $n \times n$.

- A is diagonalizable $\Leftrightarrow \mathbb{R}^n$ has a basis, say $\mathcal{B} = \{v_1, \dots, v_n\}$, of eigenvectors of A , associated to eigenvalues $\lambda_1, \dots, \lambda_n$. In this case, $P^{-1}AP = \Lambda$, where $P = (v_1 \ \cdots \ v_n)$ and Λ is a diagonal matrix with entries $\lambda_1, \dots, \lambda_n$.
- If A is diagonalizable, and T is the linear operator defined by $Tx = Ax$, then $[T]_{\mathcal{B}}^{\mathcal{B}} = \Lambda$. Thus diagonalization of A is the same as finding a basis w.r.t. which the matrix of T (defined by $Tx = Ax$) is diagonal.
- Eigenvectors associated to distinct eigenvalues are linearly independent. In particular, if A has n distinct eigenvalues, A is diagonalizable.

Question: If λ and μ are eigenvalues of A and B respectively, then are $\lambda\mu$ and $\lambda + \mu$ eigenvalues of AB and $A + B$ respectively?

Reading Slide - Eigenvalues of AB and $A + B$

- If λ is an eigenvalue of A , μ is an eigenvalue of B , is $\lambda\mu$ an eigenvalue of AB ?

False Proof. $ABx = A(\mu x) = \mu(Ax) = \lambda\mu x$.

This is false since A and B may not have same eigenvector x .

- **Ex:** $A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, $B = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, $AB = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$.

The eigenvalues of A and B are 0, 0 and that of AB are 1, 0.

- Eigenvalues of $A + B$ are NOT $\lambda + \mu$.

In above example, $A + B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ has eigenvalues 1, -1.

- If A and B have **same eigenvectors** associated to λ and μ , then $\lambda\mu$ and $\lambda + \mu$ are eigenvalues of AB and $A + B$ respectively.

Q: When do A and B have the same eigenvectors?

Extra Reading: Simultaneous Diagonalizability

- Assume A and B are diagonalizable. Then A and B have same eigenvector matrix S if and only if $AB = BA$.
- Proof.** (\Rightarrow) Assume $S^{-1}AS = \Lambda_1$ and $S^{-1}BS = \Lambda_2$, where Λ_1 and Λ_2 are diagonal matrices.

$$\text{Then } AB = (S\Lambda_1 S^{-1})(S\Lambda_2 S^{-1}) = S(\Lambda_1\Lambda_2)S^{-1} \\ \text{and } BA = S(\Lambda_2\Lambda_1)S^{-1}.$$

Since $\Lambda_1\Lambda_2 = \Lambda_2\Lambda_1$, we get $AB = BA$.

- (Part of \Leftarrow) Assume $AB = BA$.

If $Ax = \lambda x$, then $ABx = B(Ax) = B(\lambda x) = \lambda Bx$.

If $Bx = 0$, then x is an eigenvector of B , associated to $\mu = 0$.

If $Bx \neq 0$, then x and Bx both are eigenvectors of A , associated to λ .

Special case: Assume all the eigenspaces of A are one dimensional. Then $Bx = \mu x$ for some scalar $\mu \Rightarrow x$ is an eigenvector of B . We will not prove the general case. □

Eigenvalues of A^k

- If $Av = \lambda v$, then $A^2v = A(Av) = A(\lambda v) = \lambda(Av) = \lambda^2v$.

Similarly $A^k v = \lambda^k v$ for any $k \geq 0$.

Thus if v is an eigenvector of A with associated eigenvalue λ , then v is also an eigenvector of A^k with associated eigenvalue λ^k for $k \geq 0$. If A is invertible, then $\lambda \neq 0$. Hence, the same also holds for $k < 0$ since $A^{-1}v = \lambda^{-1}v$.

- If A is diagonalizable, then $P^{-1}AP = \Lambda$ is diagonal where columns of P are eigenvectors of A .

Since $(P^{-1}A^kP) = \Lambda^k$, which is diagonal, we see that A^k is diagonalizable, and the eigenvectors of A^k are same as eigenvectors of A . Similarly, the same also holds for $k < 0$ if A is invertible.

Reading Slide - Application: Fibonacci Numbers

Let $F_0 = 0$, $F_1 = 1$ and $F_k = F_{k-1} + F_{k-2}$ for $k \geq 2$ define the Fibonacci sequence. What is the k th term?

If $u_k = \begin{pmatrix} F_{k+1} \\ F_k \end{pmatrix}$, then $\begin{pmatrix} F_{k+1} \\ F_k \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} F_k \\ F_{k-1} \end{pmatrix}$, i.e.,

$u_k = Au_{k-1}$ for $n \geq 1$, where $A = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \Rightarrow u_k = A^k u_0$ for $k \geq 1$.

Characteristic polynomial of A : $\lambda^2 - \lambda - 1$

Eigenvalues: $\lambda_1 = \frac{1 + \sqrt{5}}{2}$, $\lambda_2 = \frac{1 - \sqrt{5}}{2}$.

There are 2 distinct eigenvalues \Rightarrow the associated eigenvectors x_1 and x_2 are linearly independent $\Rightarrow \{x_1, x_2\}$ is a basis for \mathbb{R}^2 .

Write $u_0 = c_1 x_1 + c_2 x_2$. Then $u_k = A^k u_0 = A^k (c_1 x_1 + c_2 x_2)$
 $= c_1 A^k x_1 + c_2 A^k x_2 = c_1 \left(\frac{1 + \sqrt{5}}{2} \right)^k x_1 + c_2 \left(\frac{1 - \sqrt{5}}{2} \right)^k x_2$.

Q: Find x_1 , x_2 , c_1 and c_2 and get the exact formula for F_k .

An Application: Predator-Prey Model

Let the owl and rat populations at time k be O_k and R_k respectively. Owls prey on the rats, so if there are no rats, the population of owls will go down by 50%. If there are no owls to prey on the rats, then the rat population will increase by 10%.

In particular, the rat and owl populations dependence is as follows:

$$\begin{aligned}O_{k+1} &= 0.5O_k + 0.4R_k \\R_{k+1} &= -pO_k + 1.1R_k\end{aligned}$$

The term $-p$ calculates the rats preyed by the owls.

Thus, if $P_k = \begin{pmatrix} O_k \\ R_k \end{pmatrix}$ and $A = \begin{pmatrix} 0.5 & 0.4 \\ -p & 1.1 \end{pmatrix}$, then $P_{k+1} = AP_k$ for all k . In particular, $P_k = A^k P_0$.

Exercise: If we start with a certain initial population of owls and rats, how many will be there in, say, 50 years, i.e., given P_0 , what is P_{50} ? What is the steady state, i.e., what is $\lim_{k \rightarrow \infty} P_k$?

An Application: Steady State

Suppose we have a system where the current state u_k depends on the previous one u_{k-1} linearly, i.e., $u_k = Au_{k-1}$. Then observe that $u_k = A^k u_0$. The steady state of the system is $u_\infty = \lim_{k \rightarrow \infty} (u_k)$. How do we find this?

- If u_0 is an eigenvector of A associated to λ , then $u_k = \lambda^k u_0$.
- Let v_1, \dots, v_r be eigenvectors of A associated respectively to $\lambda_1, \dots, \lambda_r$. If $u_0 \in \text{Span}\{v_1, \dots, v_r\}$, i.e., $u_0 = c_1 v_1 + \dots + c_r v_r$ for scalars c_1, \dots, c_r , then
$$u_k = A^k u_0 = c_1 A^k v_1 + \dots + c_r A^k v_r = c_1 \lambda_1^k v_1 + \dots + c_r \lambda_r^k v_r.$$
In particular, if A is diagonalizable, then there is a basis of \mathbb{R}^n of eigenvectors of A . Hence, this is applicable to every $u_0 \in \mathbb{R}^n$.

Let A be diagonalizable, and u_k represent population.

- Under what conditions will there be a population explosion?
- What conditions will force the population to become extinct?
- When does it stabilise (to a non-zero value)?

Hint: Depends on $|\lambda_i|$.

Complex Eigenvalues

Ex: Rotation by 90° in \mathbb{R}^2 is given by $Q = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. It has no eigenvectors since rotation by 90° changes the direction. It has no **real** eigenvectors.

Q has eigenvalues, but they are **not real**. $\det(Q - \lambda I) = \lambda^2 + 1 \Rightarrow \lambda_1 = i$ and $\lambda_2 = -i$, where $i^2 = -1$.

Let us compute the eigenvectors.

$$(Q - iI)x_1 = \begin{pmatrix} -i & -1 \\ 1 & -i \end{pmatrix} x_1 = 0 \Rightarrow x_1 = \begin{pmatrix} 1 \\ -i \end{pmatrix},$$

$$(Q + iI)x_2 = \begin{pmatrix} i & -1 \\ 1 & i \end{pmatrix} x_2 = 0 \Rightarrow x_2 = \begin{pmatrix} 1 \\ i \end{pmatrix}.$$

The eigenvalues, though imaginary, are distinct, hence eigenvectors are linearly independent.

$$\text{If } P = (x_1 \ x_2) = \begin{pmatrix} 1 & 1 \\ -i & i \end{pmatrix}, \text{ then } P^{-1}QP = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}.$$