

Problem Set 8

Aaron Wang

April 4 2024

1. Including the instructor, there are 32 people in our class. Prove that two of these people were born on the same day of the month.

Proof. Consider the set P which represents the people in our class and the set D which represents the days of the month. Notice that $|P| = 32$ and $|D| = 31$. Let $f : P \rightarrow D$ given by $f(p_i) :=$ the day of the month that they p_i was born. Since $|P| = 32 > 31 = |D|$, we know that f is not injective by PHP. So, by definition, there exists $p_i, p_j \in P$ such that $f(p_i) = f(p_j)$ meaning there are two people who were born on the same day. Q.E.D.

2. As of the 28th of March, 2024, there are over 8.1 billion people living on Earth. A person's heart will beat no more than 7×10^9 times over their lifespan. Show that there are two currently-living people on Earth whose hearts have beat the exact same number of times.

Proof. Consider the set P which represents the people in the world and the set H which represents the number of heartbeats they've had. Notice that $|P| = 8.1 \times 10^9$ and $|H| = 7 \times 10^9$. Let $f : P \rightarrow D$ given by $f(p_i) :=$ the number of heartbeats p_i has had. Since $|P| = 8.1 \times 10^9 > 7 \times 10^9 = |D|$, we know that f is not injective by PHP. So, by definition, there exists $p_i, p_j \in P$ such that $f(p_i) = f(p_j)$ meaning there are two people who have had the same heartbeats. Q.E.D.

3. Let $n \in \mathbb{N}_+$ and consider $\mathcal{A} \subseteq \mathbb{N}$ such that $|\mathcal{A}| = n + 1$. Prove there exist $x, y \in \mathcal{A}$ with $x \neq y$ such that $n \mid x - y$.

Proof. Let $n \in \mathbb{N}_+$ and consider $\mathcal{A} \subseteq \mathbb{N}$ such that $|\mathcal{A}| = n + 1$. Define $f : \mathcal{A} \rightarrow n$ where

$$f(a) = r \text{ such that } (\exists q, r \in \mathbb{Z})(a = qn + r) \text{ and } 0 \leq r < n$$

Observe that $|\mathcal{A}| = n + 1 > n = |n|$ so we know that f is not injective by PHP. So, by definition, there exists $a_i, a_j \in \mathcal{A}$ such that $f(a_i) = f(a_j)$. Let $q_i, q_j, r_i, r_j \in \mathbb{Z}$ and let them satisfy $a_i = q_i n + r_i$ such that $0 \leq r_i < n$ and $a_j = q_j n + r_j$ such that $0 \leq r_j < n$. We know $r_i = r_j$ because $f(a_i) = f(a_j)$. Consequently, $a_i - a_j = (q_i n + r_i) - (q_j n + r_j) = (q_i - q_j)n$. We know that $(q_i - q_j) \in \mathbb{Z}$ because $q_i, q_j \in \mathbb{Z}$ and $a_i - a_j = (q_i - q_j)n$, so we know that $n \mid a_i - a_j$. Consequently, there exist $x, y \in \mathcal{A}$ with $x \neq y$ such that $n \mid x - y$. Q.E.D.

4. Consider $\mathcal{S} := \{3, 4, 7, 8, 9, 10, 12, 15, 18, 19, 27, 28\}$ and $\mathcal{X} \subseteq \mathcal{S}$ with $|\mathcal{X}| \geq 9$. Show that there exist three *distinct* elements $x_1, x_2, x_3 \in \mathcal{X}$ such that $x_1 + x_2 + x_3 = 40$.

Proof. Consider $\mathcal{S} := \{3, 4, 7, 8, 9, 10, 12, 15, 18, 19, 27, 28\}$. Let $X \subseteq \mathcal{S}$ with $|X| \geq 9$. Since X is a subset of \mathcal{S} , X will only contain elements that are already in \mathcal{S} . Observe that there are 4 sets, all with distinct elements, that sum up to 40: $\{3, 10, 27\}$; $\{4, 8, 28\}$; $\{7, 15, 18\}$; $\{9, 12, 19\}$. Thus if we show that one of these triples always exists in X , we have shown that there exists three distinct elements $x_1, x_2, x_3 \in \mathcal{X}$ such that $x_1 + x_2 + x_3 = 40$. Using this information, let us define $A := \{\{3, 10, 27\}, \{4, 8, 28\}, \{7, 15, 18\}, \{9, 12, 19\}\}$ and define $f : X \rightarrow A$ where

$$f(x) = a \text{ such that } x \in a$$

Note that f is a function because every input has a unique output. Also notice that by PHP, since $|X| \geq 9$ and $|A| = 4$, there exists always exists three distinct elements of \mathcal{X} which map to the same element of A . In other words, there exists $x_1, x_2, x_3 \in \mathcal{X}$, such that $f(x_1) = f(x_2) = f(x_3)$. Finally, by definition of f , since three distinct elements map to the same value, $x_1 + x_2 + x_3 = 40$. Q.E.D.

5. Recall that $\binom{n}{0} = \binom{n}{n} = 1$ for all $n, k \in \mathbb{N}$ when $k \leq n$.

- (a) Show $\binom{n}{k} = \binom{n}{n-k}$ for all $n, k \in \mathbb{N}$ where $k \leq n$.

Proof. Assume $n, k \in \mathbb{N}$ where $k \leq n$. Let $A := \{z | z \subseteq n \wedge |z| = k\}$ and $B := \{z | z \subseteq n \wedge |z| = n - k\}$. Let $f : A \rightarrow B$ given by $f(a) := n \setminus a$.

Observe that f is injective. Assume $a_1, a_2 \in A$ and $f(a_1) = f(a_2)$. This implies that $n \setminus a_1 = n \setminus a_2$. Towards a contradiction suppose $a_1 \neq a_2$. This means that there exists $z \in a_1$ such that $z \notin a_2$. Because $a_1 \subseteq n$, $z \in n$. Since $z \in n \wedge z \notin a_2$, $z \in n \setminus a_2$ which means that $z \in n \setminus a_1$. However, $z \in a_1$ so $z \notin n \setminus a_1$. \nmid Thus $a_1 = a_2$ and f is injective.

Let $g : B \rightarrow A$ given by $g(b) := n \setminus b$. Observe that g is injective by the same logic that f is injective.

Since f is injective $|A| \leq |B|$ and since g is injective $|B| \leq |A|$. Thus we know $|A| = |B|$ by Cantor-Schöder-Bernstein. Consequently as $\binom{n}{k} = |A|$ and $\binom{n}{n-k} = |B|$, we know that $\binom{n}{k} = \binom{n}{n-k}$. Q.E.D.

(b) Show $\binom{n+1}{k+1} = \binom{n}{k+1} + \binom{n}{k}$ for all $n, k \in \mathbb{N}$ where $k \leq n$.

Proof. Assume $n, k \in \mathbb{N}$ where $k \leq n$. Let $A := \{z | z \subseteq n+1 \wedge |z| = k+1\}$ and $B := \{z | z \subseteq n \wedge (|z| = k+1 \vee |z| = k)\}$. Let $f : A \rightarrow B$ given by

$$f(a) := \begin{cases} a & \text{if } (n \notin a) \\ a \setminus \{n\} & \text{if } (n \in a) \end{cases}$$

Let $a_1, a_2 \in A$ and assume $f(a_1) = f(a_2)$.

$|f(a_1)| = k+1$:

If $|f(a_1)| = k+1$, $|f(a_2)| = k+1$. Suppose towards a contradiction that $n \in a_1$. This means that $f(a_1) = a_1 \setminus \{n\}$ which means that $|f(a_1)| = k$. \nmid . The same logic applies for a_2 . As a result, we know that $n \notin a_1, a_2$ so $f(a_1) = a_1$ and $f(a_2) = a_2$. Consequently, $a_1 = a_2$.

$|f(a_1)| = k$:

If $|f(a_1)| = k$, $|f(a_2)| = k$. Suppose towards a contradiction that $n \notin a_1$. This means that $f(a_1) = a_1$ which means that $|f(a_1)| = k+1$. \nmid . The same logic applies for a_2 . As a result, we know that $n \in a_1, a_2$ so $f(a_1) = a_1 \setminus \{n\}$ and $f(a_2) = a_2 \setminus \{n\}$. Furthermore, since we know $a_1 \setminus \{n\} = a_2 \setminus \{n\}$ and $n \in a_1, a_2$, we conclude that $a_1 = a_2$.

Consequently, f is injective.

Furthermore, we need to show that f is surjective. Let $b \in B$. $|b| = k+1$ or $|b| = k$ by definition of B .

$|b| = k+1$:

Because $b \subseteq A$ and $|b| = k+1$, $b \in A$ and $f(b) = b$. Therefore, there exists an input in A for $b \in B$ such that $|b| = k+1$.

$|b| = k$:

Because $b \subseteq A$ and $|b| = k$, $b \cup \{n\} \in A$ and $f(b \cup \{n\}) = b$. Therefore, there exists an input in A for $b \in B$ such that $|b| = k$.

Thus, for every element of B , there is an input in A which maps to that element of B . Consequently, f is surjective.

Observe that f is injective and surjective so f is bijective. Thus we know $|A| = |B|$. Observe that $\binom{n+1}{k+1} = |A|$. Let $C := \{z | z \subseteq n \wedge |z| = k+1\}$ and $D := \{z | z \subseteq n \wedge |z| = k\}$.

$\binom{n}{k+1} + \binom{n}{k} = |C| + |D|$. Additionally, $|C| + |D| = |C \cup D|$ because $|C \cap D| = 0$ as C only contains sets of cardinality $k+1$ and D only contains sets of cardinality k . Additionally observe that $B = C \cup D$ by definition so $\binom{n}{k+1} + \binom{n}{k} = |B|$. Consequently as $\binom{n+1}{k+1} = |A|$ and $\binom{n}{k+1} + \binom{n}{k} = |B|$, we know that $\binom{n+1}{k+1} = \binom{n}{k+1} + \binom{n}{k}$. Q.E.D.

6. Prove that $|\mathbb{P}(X)| = 2^{|X|}$ for any finite set X .

Proof. Basis Step:

Let X be a set such that $|X| = 0$ so $X = \emptyset$.

Show that $|\mathbb{P}(\emptyset)| = 2^{|\emptyset|}$

$|\mathbb{P}(\emptyset)| = |\{\emptyset\}| = 1$.

$2^{|\emptyset|} = 2^0 = 1$

Inductive Step:

Let X be a set such that $|X| = n$.

Inductive Hypothesis: $|\mathbb{P}(X)| = 2^{|X|}$.

Let a, Y be sets such that $a \notin X$.

Let's define $Y := X \cup \{a\}$.

Observe that $|Y| = n + 1$.

Show that $|\mathbb{P}(Y)| = 2^{|Y|}$

Define $Z := \{z | (\exists w \in \mathbb{P}(X))(z = w \cup \{a\})\}$

Observe that $\mathbb{P}(Y) = \mathbb{P}(X) \cup Z$ as $\mathbb{P}(X)$ contains all the subsets of Y which don't contain a and Z contains the subsets which contain a .

Intersection of the two sets is empty because $(\forall z \in Z)(a \in z)$ and $(\forall x \in \mathbb{P}(X))(a \notin x)$.

Consequently, $|\mathbb{P}(Y)| = |\mathbb{P}(X)| + |Z|$.

Let's define a function $f : \mathbb{P}(X) \rightarrow Z$ where $f(x) = x \cup a$.

Let $x_1, x_2 \in \mathbb{P}(X)$. Assume $f(x_1) = f(x_2)$. We know that $x_1 \cup a = x_2 \cup a$. Since $a \notin x_1, x_2$, we know $x_1 = x_2$. Thus, f is injective.

Furthermore, let $z \in Z$. By definition of Z , $\exists x \in \mathbb{P}(X)$ such that $z = x \cup \{a\}$. Therefore f is surjective.

Since f is surjective and injective, it is bijective. Consequently $|\mathbb{P}(X)| = |Z|$.

Thus, $|\mathbb{P}(X)| + |Z| = |\mathbb{P}(X)| + |\mathbb{P}(X)| = 2 \cdot |\mathbb{P}(X)|$.

Using the IH, we know that $2 \cdot |\mathbb{P}(X)| = 2 \cdot 2^{|X|}$.

$2^{|Y|} = 2^{|X|+1} = 2 \cdot 2^{|X|}$

Therefore $|\mathbb{P}(Y)| = 2^{|Y|}$.

By induction, we have shown that $\forall X (|\mathbb{P}(X)| = 2^{|X|})$.

Q.E.D.