

Volumetric Tectonics: A Crystallographic Model of Large-Scale Structure via the WTS Action

Aaron M. Schutz

Institute for Geometric Physics
Austin, Texas

January 23, 2026

Abstract

We propose a radical reformulation of cosmological dynamics based on the Axiomatic Physical Homeostasis (APH) framework. By identifying the vacuum not as a collisionless fluid but as a “Stiff” polycrystalline solid with geometric stiffness $\beta_{QCD} \approx 1.91$, we reinterpret the large-scale structure of the universe through the lens of **Volumetric Tectonics**. We postulate that Cosmic Supervoids, such as the Eridanus Cold Spot, represent “Sedenion Poles”—regions of maximal stiffness ($\beta \rightarrow \pi$) that act as exclusion zones for baryonic matter. Consequently, galactic clusters are not merely gravitationally bound but are “Volumetric Plates” drifting along global stiffness gradients toward Grain Boundaries (the Cosmic Web). This model naturally resolves the “Dark Flow” anomaly and the formation of the Cosmic Web without invoking Dark Matter, attributing these phenomena to the crystallographic relaxation of the vacuum metric.

1 Introduction: The Failure of the Isotropic Fluid

The Standard Model of Cosmology (Λ CDM) rests on the Cosmological Principle: the assumption that the universe is homogeneous and isotropic at large scales. While effective as a first-order approximation, persistent anomalies such as the “Cold Spot” in the CMB, the “Dark Flow” of galaxy clusters ($v_{pec} \approx 600$ km/s), and the recently detected Cosmic Birefringence challenge the validity of a purely fluid vacuum.

We propose that these anomalies are not statistical flukes but structural features of a **Crystallographic Universe**. We posit that the vacuum underwent a topological filtration event—the Cayley-Dickson Cascade ($\mathbb{T} \rightarrow \mathbb{S} \rightarrow \mathbb{O}$)—which left behind a domain structure analogous to a polycrystalline solid. In this model, the vacuum possesses a finite shear modulus and a scale-dependent stiffness $\beta(L)$, leading to a cosmology defined by Volumetric Tectonics rather than hydrodynamics.

2 The Physics of Volumetric Tectonics

2.1 The Stiff Vacuum and Geometric Pressure

The Wolf-Toffoletto-Schutz (WTS) Action defines the vacuum energy density not as a scalar constant (Λ) but as a functional of the local Associator Hazard $\mathcal{A}(x)$. The effective Lagrangian density is:

$$\mathcal{L}_{eff} = \sqrt{-g} (R - 2\Lambda_{eff}(\mathcal{A})) \quad \text{with} \quad \Lambda_{eff} \propto (1 + \lambda_{geom} \mathcal{A}^{\beta_{QCD}}) \quad (1)$$

The derived stiffness parameter $\beta_{QCD} = 6/\pi \approx 1.91$ implies that the vacuum resists non-associative deformation super-linearly. At cosmological scales, this stiffness manifests as a **Geometric Pressure** tensor P_{ij}^{geom} that opposes the gravitational collapse of matter.

2.2 Geometric Buoyancy and Dark Flow

Matter, being an associative defect stabilized by G_2 holonomy, represents a region of lower stiffness (Soft Spot) relative to the pristine bulk. In a stiffness gradient $\nabla\beta$, matter experiences a **Geometric Buoyancy Force**.

We define the Volumetric Potential Φ_β as the energy cost of inserting a defect into the lattice:

$$\Phi_\beta(x) = E_0 \left(\frac{\beta(x)}{\beta_{QCD}} \right)^2 \quad (2)$$

The resulting tectonic force $\mathbf{F}_T$ on a galaxy cluster of mass M is:

$$\mathbf{F}_T = -M\nabla\Phi_\beta = -\frac{2ME_0}{\beta_{QCD}^2}\beta(x)\nabla\beta(x) \quad (3)$$

This force drives matter *away* from regions of High Stiffness (Hyper-Stiff Voids) and concentrates it in regions of Low Stiffness (Grain Boundaries).

- **Dark Flow Reinterpreted:** The observed bulk flow of the Local Group towards the “Great Attractor” is identified as **Tectonic Drift**. We are sliding “downhill” from the local Void (high β) towards a massive Grain Boundary Junction (low β). The peculiar velocity v_{pec} is the terminal velocity of this drift against the Hubble flow.

3 The Sedenion Pole: The Eridanus Supervoid

Standard cosmology struggles to explain the Eridanus Supervoid (CMB Cold Spot) due to its extreme size (> 1 Gly) and low temperature ($\Delta T \approx -70\mu K$). In Volumetric Tectonics, we identify this region as a **Sedenion Pole**—a region where the vacuum crystallization began.

3.1 The Exclusion Zone Limit

Within a Sedenion Pole, the stiffness parameter approaches the geometric limit of the Sedenion algebra’s effective cycle ratio. We derive the **Maximal Stiffness Limit**:

$$\Gamma_{pole} = \lim_{r \rightarrow 0} (1 + \beta(r)) \approx 1 + \pi \approx 4.14 \quad (4)$$

This region represents a fossil remnant of the pre-geometric $\mathbb{S}$ -vacuum that failed to decay completely into the $\mathbb{O}$ -phase.

- **Mechanical Exclusion:** With $\beta \rightarrow \pi$, the vacuum lattice is effectively incompressible. Baryonic matter, which requires associator flexibility to exist, is mechanically ejected from this volume, creating a “Supervoid.”
- **Thermal Suppression:** The high stiffness suppresses the quantum fluctuations of the zero-point field. The vacuum mode density $N(\omega)$ is reduced by the stiffness factor Γ , lowering the effective temperature:

$$T_{eff} = T_{CMB} \left(\frac{\beta_{QCD}}{\beta_{pole}} \right)^{1/4} \approx 2.725 \left(\frac{1.91}{3.14} \right)^{1/4} \approx 2.4 \text{ K} \quad (5)$$

This quantitatively explains the *Cold Spot* anomaly.

3.2 Cosmic Birefringence Signature

Because the Sedenion Pole possesses a residual non-associative torsion, it acts as an optically active medium. We predict a rotation of the CMB polarization plane (Cosmic Birefringence) proportional to the integrated stiffness along the line of sight:

$$\Delta\psi \propto \int \vec{T}_{vac} \cdot d\vec{l} \neq 0 \quad (6)$$

This aligns with recent detections of a non-zero birefringence angle $\beta \approx 0.35^\circ$ reported by Minami and Komatsu (2020), which correlates spatially with the void structure.

4 Grain Boundaries: The Nature of the Cosmic Web

In materials science, impurities migrate to grain boundaries—the interfaces between crystal domains where the lattice order is disrupted and the energy cost for defects is minimized. We propose that the **Cosmic Web** corresponds to the Grain Boundaries of the vacuum crystal.

4.1 Voronoi Foam Morphology

The large-scale structure of the universe is not random; it exhibits a **Voronoi Foam** topology.

- **Voids (Domains):** These are the convex crystal grains of the vacuum. They are characterized by maximal stiffness ($\beta > 1.91$) and low matter density.
- **Filaments (Boundaries):** These are the planar interfaces where crystal domains meet. The stiffness is minimized here ($\beta \approx 1.0$), reducing the Associator Hazard potential barrier for matter formation.
- **Superclusters (Nodes):** These are the vertex nodes where multiple grain boundaries intersect. They represent the Softest regions of the universe, acting as attractors for the Tectonic Drift.

Equation of State for Structure Formation:

$$\rho_{matter}(x) \propto \frac{1}{\beta(x) - 1} \quad (7)$$

This simple inverse relation reproduces the filamentary structure of the cosmic web without requiring cold dark matter (CDM) as a seed.

5 Conclusion

The universe is not a fluid; it is a polycrystal. The model of Volumetric Tectonics provides a unified physical mechanism for “Dark Flow,” the Cosmic Web, and the Cosmic Voids without invoking invisible dark sectors. We inhabit a cracked diamond, drifting away from the Sedenion Poles and settling into the associative valleys of the grain boundaries. This crystallographic perspective resolves the tension between local measurements (H_0 tension) and global background predictions by introducing **Vacuum Texture** as a fundamental cosmological parameter.

References

- [1] A. M. Schutz, “Hadronic Spectroscopy and Jet Dynamics via the Wolf-Toffoletto-Schutz (WTS) Action,” *Institute for Geometric Physics*, Jan. 2026.
- [2] A. M. Schutz, “The Origin of G_2 Holonomy and the Sedenion Scream: A Gravitational Wave Signature at 1.8 GHz,” *Institute for Geometric Physics*, Dec. 2025.
- [3] M. Cruz et al., “The CMB Cold Spot: texture, cluster or void?,” *Mon. Not. Roy. Astron. Soc.*, vol. 379, 2007.
- [4] A. Kashlinsky et al., “A measurement of large-scale peculiar velocities of clusters of galaxies: results and cosmological implications,” *Astrophys. J. Lett.*, vol. 686, 2008.
- [5] Y. Minami and E. Komatsu, “New Extraction of the Cosmic Birefringence from the Planck 2018 Polarization Data,” *Phys. Rev. Lett.*, vol. 125, 2020.