

CO2 separation

18.04.2017

Prep list:

- Lauderdale paper
- Lauderdale slides
- Ncview control timmean
- Ncview hist timmean

CO₂ separation

Aaron Spring

18.04.2017

Lauderdale et al. 2016

Citation:

Lauderdale, J. M., S. Dutkiewicz, R. G. Williams, and M. J. Follows (2016), Quantifying the drivers of ocean-atmosphere CO₂ fluxes, *Global Biogeochem. Cycles*, 30, doi:10.1002/2016GB005400.

Quantifying the drivers of ocean-atmosphere CO₂ fluxes

Jonathan M. Lauderdale¹, Stephanie Dutkiewicz^{1,2}, Richard G. Williams³, and Michael J. Follows¹

¹Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA, ²Center for Global Change Science, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA, ³Department of Earth, Ocean and Ecological Sciences, School of Environmental Science, University of Liverpool, Liverpool, Merseyside, UK

Assumption: steady state

$$F_{\text{CO}_2} = \gamma_\theta \frac{F_{\text{heat}}}{\rho C_p} + \frac{F_W}{\rho_{\text{fw}}} \left(\gamma_S \bar{S} + \gamma_{A_T} \bar{A_T} - \bar{C_T} \right) - R_{C_T:P} \left(-\nabla \cdot (\vec{u}P) + \nabla \cdot (\kappa \nabla P) \right) h - \frac{1}{2} R_{\text{CaCO}_3} R_{C_T:P} \left(-\nabla \cdot (\vec{u}P) + \nabla \cdot (\kappa \nabla P) \right) h + \left(-\nabla \cdot (\vec{u}C_{\text{res}}) + \nabla \cdot (\kappa \nabla C_{\text{res}}) \right) h$$

CO₂ flux driver:
Heat flux
Freshwater flux
Organic matter sources and sinks
Calcium carbonate sources and sinks
Transport of disequilibrium component

negative CO₂ flux = outgassing!


Key Points:

- We have developed a quantitative framework for diagnosing regional drivers of air-sea CO₂ fluxes
- Components can be evaluated in a model or can be derived from operational data, climatologies, and ocean state estimates
- Model CO₂ fluxes result from a balance between air-sea heat fluxes, biological activity, and disequilibrium

Table 2. Values of the Linear Solubility Coefficients Used in the Attribution of Saturated Carbon Changes^a



Coefficient	Gradient	Units
γ_θ	-8.72	mmol C m ⁻³ °C ⁻¹
γ_S	-5.93	mmol C m ⁻³ psu ⁻¹
γ_{A_T}	0.81	mmol C (mmol eq) ⁻¹

^aCoefficients were empirically diagnosed by calculating C_{sat} over a range of values for temperature, salinity, or alkalinity while holding the others (including atmospheric CO₂) at surface mean values [Lewis and Wallace, 1998; Goodwin and Lenton, 2009; Ito and Follows, 2013] and finding the gradient by linear regression.



What can we learn from this?

- Fresh water: alkalinity flux and dilution flux cancel out
- Bio and residual are largest contributors, but depend on each other by definition



Adaptation to my large ensemble runs

- Goal: identify drivers of CO₂ variability

$$F_{\text{CO}_2} = \gamma_\theta \frac{F_{\text{heat}}}{\rho C_p} + \frac{F_W}{\rho_{\text{fw}}} (\gamma_S \bar{S} + \gamma_{A_T} \bar{A_T} - \bar{C_T})$$

γ_θ : aoflnhwo: net heat flux over water
 γ_S : pem: water flux to ocean [m/s]
 γ_{A_T} : CO₂ flux driver: Heat flux
 $\bar{C_T}$: Freshwater flux

- (caex90+coex90)

biology

+ change in ensmean co2flux since 1870

climate change


+ residual

circulation

Table 2. Values of the Linear Solubility Coefficients Used in the Attribution of Saturated Carbon Changes^a

Coefficient	Gradient	Units
γ_θ	+8.72	mmol C m ⁻³ °C ⁻¹
γ_S	+5.93	mmol C m ⁻³ psu ⁻¹
γ_{A_T}	-0.81	mmol C (mmol eq) ⁻¹

^aCoefficients were empirically diagnosed by calculating C_{sat} over a range of values for temperature, salinity, or alkalinity while holding the others (including atmospheric CO₂) at surface mean values [Lewis and Wallace, 1998; Goodwin and Lenton, 2009; Ito and Follows, 2013] and finding the gradient by linear regression.



$$\text{CO}_2\text{flux} = \text{heatflux} + \text{freshwater} + \text{biology} + \text{climate change} + \text{residual} + \text{circulation}$$


positive CO₂flux = outgassing!

Comparing Lauderdale to pi-control

NCL plots: normal view or ncview live



Comparing Lauderdale to pi-control




CO₂flux=heatflux+freshwater+biology+climate change+residual circulation

$$\text{CO}_2\text{flux} = \text{heatflux} + \text{freshwater} + \text{biology} + \text{climate change} + \text{residual circulation}$$


My results 1980-2004

MPI-ESM LE 1980-2004 40-50°S



40-50°S	positive carbon sink trend		negative carbon sink trend		MPI-ESM LE
	PgC/8yrs	% contribution	PgC/8yrs	% contribution	PgC/yr
CO ₂ flux	-0.15	100	-0.03	100	-0.46±0.06
heat flux	-0.04	25	0.12	-366	0.13±0.06
fresh water	0.01	-4	-0.01	27	0.02±0.00
biology	0.03	-19	-0.07	197	1.09±0.03
climate change	-0.03	17	-0.01	16	-0.23±0.01
circulation	-0.12	83	-0.08	225	0.72±0.06

MPI-ESM LE 1980-2004 50-60°S



50-60°S	positive carbon sink trend		negative carbon sink trend		MPI-ESM LE
	PgC/8yrs	% contribution	PgC/8yrs	% contribution	PgC/yr
CO ₂ flux	-0.43	100	0.58	100	-0.05±0.19
heat flux	-0.04	9	0.06	11	0.14±0.05
fresh water	-0.00	1	0.00	1	0.04±0.00
biology	-0.17	39	0.17	30	-0.76±0.08
climate change	-0.06	13	-0.01	-2	-0.22±0.02
circulation	-0.17	38	0.35	60	0.75±0.12

Table 1: The trends of CO₂ flux and its contributions for the zonal band of 50-60°S indicate circulation as the most variable contribution





Limitations

- × Thermal pump via heat flux and not SST
- × Cooling SST trends (m178) can have negative trend in CO2 flux contribution due to heat flux, but should be positive (thermal)
- × I tried $\text{bio} = \text{fddtdic} - \text{co2flux}$, but too high values

My results 1980-2005 ensmean

50-60°S	positive carbon sink trend		negative carbon sink trend		MPI-ESM LE PgC/yr
	PgC/8yrs	% contribution	PgC/8yrs	% contribution	
CO ₂ flux	-0.43	100	0.58	100	-0.05±0.19
heat flux	-0.04	9	0.06	11	0.14±0.05
fresh water	-0.00	1	0.00	1	0.04±0.00
biology	-0.17	39	0.17	30	-0.76±0.08
climate change	-0.06	13	-0.01	-2	-0.22±0.02
circulation	-0.17	38	0.35	60	0.75±0.12

Table 1: The trends of CO₂ flux and its contributions for the zonal band of 50-60°S indicate circulation as the most variable contribution



Heat refers to net air-sea radiative heat flux:

Positive carbon sink trend: SST warming

Negative carbon sink trend: SST cooling

Lauderdale et al. 2016 input

• Text


Key Points:

- We have developed a quantitative framework for diagnosing regional drivers of air-sea CO₂ fluxes
- Components can be evaluated in a model or can be derived from operational data, climatologies, and ocean state estimates
- Model CO₂ fluxes result from a balance between air-sea heat fluxes, biological activity, and disequilibrium

Quantifying the drivers of ocean-atmosphere CO₂ fluxes

Jonathan M. Lauderdale¹, Stephanie Dutkiewicz^{1,2}, Richard G. Williams³, and Michael J. Follows¹

¹Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA, ²Center for Global Change Science, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA, ³Department of Earth, Ocean and Ecological Sciences, School of Environmental Science, University of Liverpool, Liverpool, Merseyside, UK



$$F_{\text{CO}_2} = \gamma_\theta \frac{F_{\text{heat}}}{\rho C_p} + \frac{F_W}{\rho_{\text{fw}}} \left(\gamma_S \bar{S} + \gamma_{A_T} \bar{A_T} - \bar{C_T} \right) - R_{C_T:P} \left(-\nabla \cdot (\bar{u}P) + \nabla \cdot (\kappa \nabla P) \right) h - \frac{1}{2} R_{\text{CaCO}_3} R_{C_T:P} \left(-\nabla \cdot (\bar{u}P) + \nabla \cdot (\kappa \nabla P) \right) h + \left(-\nabla \cdot (\bar{u}C_{\text{res}}) + \nabla \cdot (\kappa \nabla C_{\text{res}}) \right) h$$

CO₂ flux driver:

Heat flux

Freshwater flux

Organic matter sources and sinks

Calcium carbonate sources and sinks

Transport of disequilibrium component

Table 2. Values of the Linear Solubility Coefficients Used in the Attribution of Saturated Carbon Changes^a

Coefficient	Gradient	Units
γ_θ	-8.72	mmol C m ⁻³ °C ⁻¹
γ_S	-5.93	mmol C m ⁻³ psu ⁻¹
γ_{A_T}	0.81	mmol C (mmol eq) ⁻¹

^aCoefficients were empirically diagnosed by calculating C_{sat} over a range of values for temperature, salinity, or alkalinity while holding the others (including atmospheric CO₂) at surface mean values [Lewis and Wallace, 1998; Goodwin and Lenton, 2009; Ito and Follows, 2013] and finding the gradient by linear regression.