

CAAM 336 · DIFFERENTIAL EQUATIONS

Homework 36 · Solutions

Posted Friday 1 November 2013. Due 5pm Wednesday 13 November 2013.

36. [25 points] Let $H_D^1(0, 1) = \{w \in H^1(0, 1) : w(1) = 0\}$ and let the inner product $(\cdot, \cdot) : L^2(0, 1) \times L^2(0, 1) \rightarrow \mathbb{R}$ be defined by

$$(v, w) = \int_0^1 v(x)w(x) dx$$

and the inner product $a(\cdot, \cdot) : H_D^1(0, 1) \times H_D^1(0, 1) \rightarrow \mathbb{R}$ be defined by

$$a(v, w) = \int_0^1 v'(x)w'(x) dx.$$

Also, let $f \in L^2(0, 1)$, let N be a positive integer, and let V_N be a subspace of $H_D^1(0, 1)$. Moreover, let $u \in H_D^1(0, 1)$ be such that

$$a(u, v) = (f, v) \text{ for all } v \in H_D^1(0, 1)$$

and let $u_N \in V_N$ be such that

$$a(u_N, v) = (f, v) \text{ for all } v \in V_N.$$

- (a) Show that

$$a(u - u_N, u - u_N) = a(u, u) - a(u_N, u_N).$$

- (b) Let $\phi_0, \phi_1, \dots, \phi_N \in V_N$ and let $\mathbf{K} \in \mathbb{R}^{(N+1) \times (N+1)}$ be the matrix with entries $K_{jk} = a(\phi_{k-1}, \phi_{j-1})$ for $j, k = 1, \dots, N+1$. Also, let

$$u_N = \sum_{j=0}^N \alpha_j \phi_j$$

where, for $k = 1, \dots, N+1$, $\alpha_{k-1} \in \mathbb{R}$ is the k th entry of the vector $\mathbf{c} \in \mathbb{R}^{N+1}$. Show that

$$\mathbf{c}^T \mathbf{K} \mathbf{c} = a(u_N, u_N).$$

Solution.

- (a) [12 points] The properties satisfied by the inner product allow us to say that

$$\begin{aligned} a(u - u_N, u - u_N) &= a(u, u - u_N) - a(u_N, u - u_N) \\ &= a(u, u) - a(u, u_N) - a(u_N, u) + a(u_N, u_N) \\ &= a(u, u) - 2a(u, u_N) + a(u_N, u_N). \end{aligned}$$

Now, $u_N \in V_N$ and so the fact that

$$a(u_N, v) = (f, v) \text{ for all } v \in V_N$$

means that

$$a(u_N, u_N) = (f, u_N).$$

Moreover, $u_N \in H_D^1(0, 1)$, since V_N is a subspace of $H_D^1(0, 1)$ and $u_N \in V_N$, and so the fact that

$$a(u, v) = (f, v) \text{ for all } v \in H_D^1(0, 1)$$

means that

$$a(u, u_N) = (f, u_N).$$

So,

$$a(u, u) - 2a(u, u_N) + a(u_N, u_N) = a(u, u) - 2(f, u_N) + (f, u_N) = a(u, u) - (f, u_N).$$

Therefore,

$$a(u - u_N, u - u_N) = a(u, u) - (f, u_N) = a(u, u) - a(u_N, u_N)$$

because

$$a(u_N, u_N) = (f, u_N).$$

(b) [13 points] We first compute that

$$\mathbf{K}\mathbf{c} = \mathbf{d}$$

where $\mathbf{d} \in \mathbb{R}^N$ is the vector with entries

$$d_j = \sum_{k=1}^{N+1} a(\phi_{k-1}, \phi_{j-1}) c_k = \sum_{k=1}^{N+1} a(\phi_{k-1}, \phi_{j-1}) \alpha_{k-1} = \sum_{k=0}^N a(\phi_k, \phi_{j-1}) \alpha_k$$

for $j = 1, \dots, N+1$. Moreover, since

$$u_N = \sum_{j=0}^N \alpha_j \phi_j = \sum_{k=0}^N \alpha_k \phi_k,$$

the properties satisfied by the inner product yield that

$$\sum_{k=0}^N a(\phi_k, \phi_{j-1}) \alpha_k = a\left(\sum_{k=0}^N \alpha_k \phi_k, \phi_{j-1}\right) = a(u_N, \phi_{j-1})$$

and so

$$d_j = a(u_N, \phi_{j-1})$$

for $j = 1, \dots, N+1$. Therefore,

$$\begin{aligned} \mathbf{c}^T \mathbf{K} \mathbf{c} &= \mathbf{c}^T \mathbf{d} \\ &= \sum_{j=1}^{N+1} c_j a(u_N, \phi_{j-1}) \\ &= \sum_{j=1}^{N+1} \alpha_{j-1} a(u_N, \phi_{j-1}) \\ &= \sum_{j=0}^N \alpha_j a(u_N, \phi_j) \\ &= a\left(u_N, \sum_{j=0}^N \alpha_j \phi_j\right) \\ &= a(u_N, u_N) \end{aligned}$$

by the properties satisfied by the inner product and the fact that

$$u_N = \sum_{j=0}^N \alpha_j \phi_j.$$
