

# Worksheet 1: Practice with Joint Measures

Name: b Aaron Zoll

Due September 8, 2023

You will investigate the rule of succession which states that the probability of success for the  $(m+1)$  th trial in a sequence of independent bernoulli random variables  $x_1, \dots, x_{m+1}$ , given  $k \leq m$  successes in the first  $m$  trials is

$$\mathbb{P} \left( x_{m+1} = 1 \mid \sum_{j=1}^m x_j = k \right) = \frac{k+1}{m+2}.$$

In principle if  $x_1, \dots, x_{m+1}$  are truly independent, you might expect the right hand side to simply be  $\mathbb{P}(x_{m+1} = 1)$  which does not depend on  $k$  (i.e. the outcomes of  $x_1, \dots, x_m$ ).

In this worksheet, you will explore which space to make sense of eq. (1). While seemingly elementary, this example provides a clear illustration of Bayesian reasoning in machine learning, where model-defining parameters are defined by a measure instead of as point-values. We shall let  $\mathcal{X} = \{0, 1\}$ ,  $x \in \mathcal{X}$  denote an outcome, and let  $s_m : \mathcal{X}^m \rightarrow \mathbb{R}$  be defined by  $(x_1, \dots, x_m) \mapsto \sum_{j=1}^m x_j$ . Implicit in eq. (1) is that  $p := \mathbb{P}(x = 1)$  is unknown (i.e. there is some distribution on  $p$ ) and that given  $p$ , then the trials  $x_1, \dots, x_{m+1}$  are independent. To concretize this observation, we let  $I = [0, 1]$  and suppose that there is joint distribution  $\mathbb{P}_{\mathcal{X}^m \times I}$  according to which  $\mathbb{P}_{\mathcal{X}^m|I}$  is independent. In particular,

$$\mathbb{P}_{\mathcal{X}^m|I} \left( \sum_{j=1}^m x_j = k \mid i = p \right) = \binom{m}{k} p^k (1-p)^{m-k}.$$

Because  $p$  does not appear explicitly in eq. (1), it must be (being) marginalized away. That means that

$$\mathbb{P}(x_{m+1} = 1 \mid s_m(x_1, \dots, x_m) = k) = \mathbb{P}_{\mathcal{X} \times I | \mathcal{X}^m}(\{x_{m+1} = 1\} \times I \mid s_m(x_1, \dots, x_m) = k)$$

A reasonable measure to place on the marginal probability  $\mathbb{P}_I$  is the uniform one  $\mathbb{P}_I((a, b)) = b - a$  for  $0 \leq a \leq b \leq 1$ . With this preamble, you may now proceed.

1. Using the definition of conditional probability, rewrite eq. (2) as a ratio of two joint probabilities. You may use that  $\mathcal{X} \times I \times \mathcal{X}^m \cong \mathcal{X}^{m+1} \times I$  to simplify (but be careful to write the correct event in the numerator!). In the denominator, express the marginal probability  $\mathbb{P}_{\mathcal{X}}$  in terms of  $\mathbb{P}_{\mathcal{X}^m \times I}$ ; you will see why in the next problem.

## Solution

$$\begin{aligned} \mathbb{P}_{\mathcal{X} | \mathcal{X}^m}(x_{m+1} = 1 | s_m(x_1, \dots, x_m) = k) &= \mathbb{P}_{\mathcal{X} \times I | \mathcal{X}^m}(\{x_{m+1} = 1\} \times I | s_m(x_1, \dots, x_m) = k) \\ &= \frac{\mathbb{P}_{\mathcal{X} \times I \times \mathcal{X}^m}(\{x_{m+1} = 1\} \times I \times \{s_m(x_1, \dots, x_m) = k\})}{\mathbb{P}_{\mathcal{X}^m}(s_m = 1)} \\ &= \frac{\mathbb{P}_{\mathcal{X}^{m+1} \times I}(\{x_{m+1} = 1\} \times \{s_m(x_1, \dots, x_m) = k\} \times I)}{\mathbb{P}_{\mathcal{X}^m \times I}(\{s_m = 1\} \times I)} \end{aligned}$$

\* See Binomial Distribution for more information.

2. Conveniently, this expression (nearly) collapses the problem to a single computation (of either numerator or denominator). To compute either probability, you will want to condition again, this time as  $\mathcal{X}^m \mid I$ , and use the law of total probability. <sup>†</sup> Express each one:

**Solution**

Let  $A = \{x_{m+1} = 1\}$  and  $C = \{s_m = k\}$  Numerator (N):

$$\begin{aligned}
& \mathbb{P}_{\mathcal{X}^{m+1} \times I}(\{x_{m+1} = 1\} \times \{s_m(x_1, \dots, x_m) = k\} \times I) \\
&= \int_I \int_{A \times C|I} d\mathbb{P}_{\mathcal{X} \times \mathcal{X}^m|I}(x_1, \dots, x_m|i) d\mathbb{P}_I(i) \\
&= \int_I \int_{A|I} \int_{C|I} d\mathbb{P}_{\mathcal{X}|I}(x|i) d\mathbb{P}_{\mathcal{X}^m|I}(x_1, \dots, x_m|i) d\mathbb{P}_I(i) \text{ (by independence)} \\
&= \int_I \mathbb{P}_{\mathcal{X}|I}(A|I) \mathbb{P}_{\mathcal{X}^m|I}(C|I) d\mathbb{P}_I(i) \\
&= \int_0^1 \mathbb{P}_{\mathcal{X}|I}(A|i = p) \mathbb{P}_{\mathcal{X}^m|I}(C|i = p) dp \\
&= \int_0^1 p \binom{m}{k} p^k (1-p)^{m-k} dp \\
&= \int_0^1 \binom{m}{k} p^{k+1} (1-p)^{m-k} dp
\end{aligned}$$

Similarly, the denominator (D):

$$\begin{aligned}
& \mathbb{P}_{\mathcal{X}^m \times I}(\{s_m(x_1, \dots, x_m) = k\} \times I) \\
&= \int_I \int_{C|I} d\mathbb{P}_{\mathcal{X}^m|I}(x_1, \dots, x_m|i) d\mathbb{P}_I(i) \\
&= \int_I \mathbb{P}_{\mathcal{X}^m|I}(C|I) d\mathbb{P}_I(i) \\
&= \int_0^1 \mathbb{P}_{\mathcal{X}^m|I}(C|i = p) dp \\
&= \int_0^1 \binom{m}{k} p^k (1-p)^{m-k} dp
\end{aligned}$$

3. Explicitly compute the inner (conditional) probability (or expectation) for both:

**Solution** [see above]

4. Show by induction on  $k$  that  $\int_0^1 p^k(1-p)^{m-k} dp = \frac{k!(m-k)!}{(m+1)!}$ .

**Solution**

For  $k = 0$

$$\begin{aligned} &= \int_0^1 p^0(1-p)^m dp \\ &= -\frac{1}{m+1}(1-p)^{m+1} \Big|_0^1 \\ &= \frac{1}{m+1} \\ &= \frac{0!(m-0)!}{(m+1)!} \end{aligned}$$

Suppose this holds for  $k < m$

For  $j = k + 1$

$$\begin{aligned} &= \int_0^1 p^j(1-p)^{m-j} dp \\ &= \int_0^1 p^{k+1}(1-p)^{m-k-1} dp \\ \text{Let } u &= p^{k+1}, \quad dv = (1-p)^{m-k-1} \\ du &= (k+1)p^k, \quad v = -\frac{1}{m-k}(1-p)^{m-k} \\ &= uv \Big|_0^1 + \frac{k+1}{m-k} \int_0^1 p^k(1-p)^{m-k} dp \\ &= 0 + \frac{(k+1)k!(m-k)!}{(m-k)(m+1)!} \\ &= \frac{(k+1)!(m-k-1)!}{(m+1)!} \\ &= \frac{j!(m-j)!}{(m+1)!} \end{aligned}$$

5. Show also, using part 4. and judicious integration by parts that  $\int_0^1 p^{k+1}(1-p)^{m-k} dp = \frac{(k+1)!(m-k)!}{(m+2)!}$ .

**Solution** (no by parts needed!)

$$\begin{aligned} &\int_0^1 p^{k+1}(1-p)^{m-k} dp \\ &= \int_0^1 p^{k+1}(1-p)^{m+1-(k+1)} dp \\ &= \frac{(k+1)!((m+1)-(k+1))!}{((m+1)+1)!} \\ &= \frac{(k+1)!(m-k)!}{(m+2)!} \end{aligned}$$

6. Use results in parts 3., 4., and 5. to compute the expression (4) and simplifying, conclude (1).

**Solution** Thus, we get

$$\begin{aligned} (1) &= \frac{N}{D} \\ &= \frac{\binom{m}{k} \frac{(k+1)!(m-k)!}{(m+2)!}}{\binom{m}{k} \frac{(k)!(m-k)!}{(m+1)!}} \\ &= \frac{(k+1)!(m+1)!}{k!(m+2)!} \\ &= \frac{k+1}{m+2} \end{aligned}$$