Modeling Volatility in Financial Markets: A Time Series Analysis of the CBOE Volatility Index (VIX)

Aarti Garaye
aartigaraye@ucsb.edu
PSTAT 174 - Time Series Analysis
Professor Tomoyuki Ichiba

March 15, 2026

Abstract

This study examines the CBOE Volatility Index (VIX), often called the "fear gauge" of financial markets, using basic time series methodologies. Unfortunately, to date, measuring and forecasting volatility accurately is still a huge challenge. In this paper, the time series properties of daily VIX data are analyzed using classical time series modeling techniques. The Box–Jenkins methodology is applied to identify and estimate an appropriate Seasonal Autoregressive Integrated Moving Average (SARIMA) model to capture the temporal dependence structure of the series. In addition, a Generalized Autoregressive Conditional Heteroskedasticity (GARCH) model is deployed to account for volatility clustering, a common feature of financial time series. The results demonstrate that while the SARIMA model effectively captures the dynamics of the conditional mean, the GARCH framework provides additional insight into the time-varying volatility of the time series.

1 Introduction

Asset price volatility is of crucial importance to scholars, investors, and policy makers. Volatility plays a central role in derivatives pricing, hedging strategies, portfolio selection, and risk management (Wang 2019). In financial markets, the CBOE Volatility Index (VIX) is widely used as a benchmark measure of expected market volatility derived from options on the S&P 500 index. Because the VIX reflects investor expectations of future market uncertainty, understanding its statistical behavior is important for both financial practitioners and researchers.

1.1 Background and Motivation

Volatility forecasting is an important task in financial markets. It has occupied the attention of academics and practitioners over the past three to four decades. It's importance is visible in an array of areas like investment, security valuation, risk management, and monetary policy. (Majmudar and Banerjee (2004)). For example, the annualized SD of US stocks over the past ten years is 16.8% compared to 12.7% over the preceding then years, a period which included the fallout from the dot-com bubble (Based on S&P 500). This heightened volatility has obvious implications for asset and derivative pricing as well as the conduct of portfolio and risk management. Volatility plays a key role in market-timing decisions, in the pricing of options and other derivatives and in determining market risk and hedge ratios. Therefore, there remains a need to provide accurate forecasts of volatility, something for which there is an established literature but with no real consensus on the preferred approach. (Kambouroudis and McMillan (2016))

Assuming efficient markets, a good source of volatility estimate is the implied volatility (VIX). One of the most widely used indicators of market volatility is the CBOE Volatility Index (VIX). The VIX is calculated from options prices on the S&P 500 index and reflects the market's expectation of volatility over the next 30 days. Due to its ability to capture investor sentiment and expectations about future uncertainty, the VIX is often referred to as the "fear index" in financial markets (Chicago Board Options Exchange 2019). Large spikes in the VIX are frequently associated with periods of financial stress such as market crashes or economic crises.

1.2 Literature Review

Previous research has extensively documented the properties of VIX. Blair, Poon, and Taylor (2001) found that VIX exhibits mean-reverting behavior and volatility clustering. Fernandes, Medeiros, and Scharth (2007) demonstrated that GARCH-type models perform well in capturing VIX dynamics. Corsi (2009) introduced the Heterogeneous Autoregressive (HAR) model for VIX forecasting, which has become a benchmark in the literature.

1.3 Methodology

Because the VIX is observed over time at a regular frequency, time series analysis provides a natural framework for studying its behavior. Statistical time series models allow researchers to analyze the dependence structure of the data, identify patterns such as trends or persistence, and generate forecasts of future values. In this project, the dynamics of the VIX are examined using classical time series methods. The Box–Jenkins methodology is applied to identify and estimate an appropriate Seasonal Autoregressive Integrated Moving Average (SARIMA) model for the time series. In addition, a Generalized Autoregressive Conditional Heteroskedasticity (GARCH) model is employed to capture volatility clustering that is commonly observed in financial data. Together, these models provide insight into both the mean dynamics and the time-varying volatility of the VIX time series.

In short, This analysis employs two complementary methodologies:

1. **SARIMA Modeling** (Box-Jenkins approach): To model the conditional mean of log-differenced VIX
2. **GARCH Modeling**: To capture the time-varying volatility and volatility clustering observed in VIX returns

2 Data

The dataset used in this project consists of daily closing values of the CBOE Volatility Index (VIX), which is a widely recognized measure of market volatility derived from options on the S&P 500 index. The VIX reflects the market's expectation of 30-day forward-looking volatility and is often referred to as the "fear index" due to its ability to capture investor sentiment during periods of financial stress (Chicago Board Options Exchange 2019).

2.1 Data Description

The VIX ranges from 0 to ∞, while there are some missing values there are no negative VIX indexes. There are a total of 9425 observations.

2.1.1 Source and Collection

The dataset is obtained from the FRED: Federal Reserve Bank of St. Louis.. he source is Chicago Board of Options Exchange and released by CBOE Market Statistics. The VIX observed when the market closed was monitored and stored in the dataset. The data is not adjusted for seasonality.

2.1.2 Time Range and Frequency

- **Start Date**: January 2, 1990
- **End Date**: February 16, 2026
- **Frequency**: Daily (business days)
- **Total Observations**: 9,425 observations

2.1.3 Variable Description

The dataset contains two columns:

1. **observation_date**: The observation date (YYYY-MM-DD format)
2. **VIXCLS**: The closing value of the VIX on that date

The VIX is quoted as an annualized percentage. For example, a VIX value of 20 corresponds to an expected annualized volatility of 20%, or approximately 1.26% on a daily basis ($\frac{20}{\sqrt{252}}$, where 252 is the typical number of trading days per year).

2.2 Data Characterstics

Some of the basic summary statistic of the dataset is given below in the table. This shows the mean, median, standard deviation, number of observations, min, and max values. Note, that I have just decided to omit the values where VIX was NA, this reduced the number of

observations from 9425 to 9124. This is still more than enough datapoints for us to work with.

Table 1: Summary Statistics of VIX

Statistic	Value
Mean	19.44
Median	17.58
Std. Dev.	7.77
Min	9.14
Max	82.69
N	9124.00

2.3 Exploratory Data Analysis

Before we begin fitting any models, it is standard practice to plot the time series plot of the data This provides a basic insight on whether we would want to perform any transformations before fitting any models.

2.3.1 Time Series Plot



Figure 1: Daily VIX Closing Values (1990-2026)

The time series plot reveals several important features. **Volatility Clustering**, periods of high volatility tend to be followed by high volatility (and vice versa). **Crisis Spikes**, notable spikes during the 2008 financial crisis (VIX > 80) and 2020 COVID-19 pandemic (VIX > 80). **Non-stationarity** clear evidence of time-varying mean and variance. And **Asymmetry**, VIX tends to rise faster (sharp spikes) than it falls (gradual decay). The red dashed line represents the mean VIX for this period which is somewhere around 20.

Another helpful insight would be to look at the histogram is this data. This gives us information about the distribution of VIX.



Figure 2: Distribution of VIX Values

The histogram shows strong positive skewness, and the Q-Q plot reveals substantial departures from normality, particularly in the right tail. This informs us that a log transform is necessary before fitting any models.

2.4 Why This?

I selected the VIX for several compelling reasons:

1. **Economic Importance**: The VIX is a critical indicator used by investors, policymakers, and researchers worldwide
2. **Rich Dynamics**: The dataset exhibits complex time series patterns including volatility clustering, mean reversion, and regime changes
3. **Methodological Challenge**: VIX provides an excellent test case for advanced time series methods including SARIMA and GARCH
4. **Practical Relevance**: Accurate VIX modeling has direct applications in trading, risk management, and portfolio construction
5. **Data Quality**: FRED provides high-quality, reliable data with comprehensive documentation

Additionally, I am interested in Mathematical Finance and have been learning about VIX in my other classes, especially the smile vs. smirk. I thought it would be interesting to blend what I am learning in a purely theoretical class (strong mathematical derviations) to fitting models and see how they are used to forecast.

3 Methodology

3.1 SARIMA(p, d, q) × (P, D, Q) Model

To capture the seasonal component of a time series, we fit a Seasonal Autoregressive Integrated Moving Average (SARIMA) model. Think of SARIMA as a generalized ARIMA model but also accounting for seasonality. The general form is:

$$\Phi_P\left(B^s\right)\phi(B)\nabla_s^D\nabla^d x_t = \delta + \Theta_Q(B^s)\theta(B)w_t; \quad t \in \mathbb{Z}$$

where $\{w_t\}$ is the Gaussian white noise, ϕ, θ are the polynomials (similar to ARIMA) and the new polynomials introduced are

$$\begin{aligned}\Phi_P(z) &:= 1 - \Phi_1 z - \Phi_2 z^2 \ - ... - \ \Phi_P z^P, \\ \Theta_Q(z) &:= 1 + \Theta_1 z + \Theta_2 z^2 + ... + \ \Theta_Q z^Q; \quad z \in \mathbb{C}\end{aligned}$$

for seasonality with the seasonal lag operator B^s and the seasonal difference operator is defined as $\nabla_s := 1 - B^s$.

VIX measure "fear" or "uncertainty." This is driven by current affairs, politics, and wars. In other words VIX is driven by news, events, and crises. However, these are not calendar patterns. Unlike retail sales (Christmas spike) or unemployment (summer hiring), market fear doesn't follow a calendar. Crises are random and unpredictable.

Furthermore, looking at the time series plot there were no regular spikes or patterns, infact the 2008 and 2020 spikes were unpredictable. If there were seasonality, we would see consistent patterns every 12 months (like higher VIX every January, or every Q4). Some may argue that there are some seasonal effects in finance like "January Effect" and "Sell in May and Go Away" (Bouman and Jacobsen (2002)) but these are for stocks return patterns, not volatility patterns

There is, however, an intraday seasonality pattern (U-shaped) (Hughes, Winters, and Rawls (2005)) observed. High volatility is observed right when the market opens and the trading

volumes slow down during the middle of the day and spike back up when the market is about to close. Remember, that in the US the market opens 9:30 am ET and closes 5:00 pm ET. (Ellul, Shin, and Tonks (2005)). However, to incorporate this seasonality patter we must have minute based, or hour based data. Our dataset has daily closing VIX index. Other weekly seasonality patterns are weak and controversial (Caporale and Plastun (2019)). Thus, for VIX, we focus on the non-seasonal ARIMA(p, d, q) component as there is no clear seasonal pattern in daily financial volatility.

3.1.1 Box-Jenkins Methodology

The Box-Jenkins approach to time series modelling consists of extracting predictable movements (or patterns) from the observed data through a series of iterations.(Makridakis and Hibon (1997)) The univariate Box-Jenkins method is purely a forecasting tool; no explanation is offered in that there are no regressor-type variables. The Box-Jenkins approach follows a four phase procedure (Aljandali (2017)):

1. **Identification**: Determine the orders p, d, and q using ACF/PACF plots and unit root tests
2. **Estimation**: Estimate model parameters using maximum likelihood
3. **Diagnostic Checking**: Verify model adequacy through residual analysis
4. **Forecasting**: Generate forecasts with confidence intervals

Note: We follow this methodology in the Results section for SARIMA models

3.2 GARCH Model

While ARIMA models capture the conditional mean dynamics, financial time series like VIX often exhibit **volatility clustering**. These are periods where large changes tend to be followed by large changes and small changes by small changes. GARCH (Generalized Autoregressive Conditional Heteroskedasticity) models explicitly capture this time-varying volatility (Bollerslev (1986))

GARCH models are used mostly when forecasting time-varying volatility in finance and economic data. It models heteroskedasticity, where error variance is not constant, by assuming current variance depends on past squared residuals (ARCH term) and past variances (GARCH term) (Engle and Bollerslev (1986)). Thus, GARCH is crucial for modeling

"volatility clustering," where high-volatility periods are followed by higher volatility and calm periods by calm, often seen in stock returns, exchange rates, and inflation. ### Specifications

Mean equation:

$$r_t = \mu + \epsilon_t$$

where $\epsilon_t = \sigma_t z_t$ and $z_t \sim N(0, 1)$

Variance equation:

$$\sigma_t^2 = \omega + \sum_{i=1}^{p} \alpha_i \epsilon_{t-i}^2 + \sum_{j=1}^{q} \beta_j \sigma_{t-j}^2$$

where: - σ_t^2 is the conditional variance at time t - $\omega > 0$ is the constant term - α_i captures the ARCH effect (impact of past shocks) - β_j captures the GARCH effect (persistence of volatility)

The most common specification is GARCH(1,1), which uses one lag of the squared error ($q = 1$) and one lag of the conditional variance ($p = 1$)

$$\sigma_t^2 = \omega + \alpha \epsilon_{t-1}^2 + \beta \sigma_{t-1}^2$$

Key properties: **Stationarity**: Requires $\alpha + \beta < 1$ **Persistence**: The sum $\alpha + \beta$ measures volatility persistence **Unconditional variance**: $\sigma^2 = \omega/(1 - \alpha - \beta)$

GARCH(1,1) is highly parsimonious (easy to estimate) and generally superior to simple ARCH models in predicting future variance because it captures long-run persistence in volatility. (Majka (n.d.))

4 Results

4.1 SARIMA Analysis

As mentioned in the EDA section, we must transform the data to stabilize the varaiance. Here we will be following the Box-Jenkins approach step-by-step. Recall, that we have already plotted the data in the EDA section so we can skip that step. ### Step 1: Data Transformation Given the nature of VIX as a strictly positive variable with time-varying variance, we apply a log transformation to stabilize variance, from now on we will be looking at log(VIX)

Once we have transformed the data, we must identify the difference d. ### Step 2: Identification of Differencing Order The ADF test checks if a time series is stationary. Stationarity is key for ARIMA/SARIMA models because these models assume the mean, variance, and autocorrelation structure don't change over time. The "augmented" part means the test adds lagged differences of the series to remove autocorrelation in residuals:

$$\Delta x_t = \alpha + \beta t + \gamma x_t + \sum_{i=1}^{p} \delta_i \Delta x_{t-i} + \epsilon_t$$

Table 2: Augmented Dickey-Fuller Test Results

Series	ADF.Statistic	P.value	Conclusion
Log(VIX)	-6.1010	0.01	Stationary
Diff Log(VIX)	-23.4845	0.01	Stationary

The ADF test tells us the log-transformed series is stationary in levels. That means it does not have a unit root, thus we do not need d in our SARIMA model. Thus, we can fit SARIMA directly on `log_vix` with $d = 0$.

4.1.1 Step 3: Estimation of Parameters

To estimate the parameters we must look at the ACF and the PACF of the transformed data.



Figure 3: ACF and PACF of Differenced Log(VIX)

The ACF for the log transformed data is a slow decaying (long-memory) and the pacf cuts off after the first lag. Since a slow decaying ACF is a characteristic of an AR process. Also from the last homework we know that the ideal model to fit here is SARIMA with $(p, d, q) = (1, 0, 0)$ and $(P, D, Q)_s = (0, 0, 0)$

However, the ACF is screaming long memory, i.e. a very slow, gradual linear decay, doesn't reach zero until around lag 400-500. This is a sign of fractional integration. Even though the ADF test suggested that $d = 0$ the ACF is saying something else.

VIX exhibits fractional integration, it's between I(0) and I(1). The true differencing order is something like $d = 0.5,\ 0.4$ or something like that (not an integer!) The ADF test only tests the unit root and rejects it, but the slowly decaying ACF suggests otherwise.

However, I have already chosen a second method, GARCH, so long memory won't be helpful here. We could use `fracdiff` or `optim` to find the maximum likelihood for possible d but right now the possible choices for fitting a sarima model are ARIMA(1,0,0) and ARIMA(1,1,1) with difference 1 coming from the ACF and just to have a third model we can use ARMIA(1,1,1).

Please refer to the separate section on long memory I have posted on my github repo for this analysis.

4.1.2 Step 4: Diagnostics and Model Estimation

Table 3: Model Comparison - Information Criteria (* indicates best model)

Model	AIC	BIC	Best_AIC	Best_BIC
ARIMA(0,1,1)	-23240.51	-23226.27		
ARIMA(1,1,0)	-23232.92	-23218.68		
ARIMA(1,1,1)	-23364.86	-23343.51	*	*

The ARIMA(1,1,1) model has the lowest AIC, while ARIMA(0,1,1) has the lowest BIC. We select **ARIMA(1,1,1)** as it provides the best balance between fit and parsimony according to AIC.

Table 4: ARIMA(1,1,1) Parameter Estimates

Parameter	Estimate	SE	t-value	p-value
ar1	0.8207	0.0289	28.4127	0
ma1	-0.9047	0.0223	-40.5541	0

All parameters are statistically significant at the 1% level.

4.1.3 Step 5: Model Selection



Figure 4: Diagnostic Plots for ARIMA(1,1,1)

Diagnostic Assessment:

1. **Standardized Residuals**: No obvious patterns or trends; appear to fluctuate around zero
2. **ACF of Residuals**: Most lags are within confidence bounds, suggesting residuals are approximately white noise
3. **Q-Q Plot**: Shows some deviation in the tails, indicating heavier tails than normal distribution
4. **Ljung-Box Statistics**: Many p-values are below 0.05, suggesting some remaining autocorrelation

While diagnostics show some departures from ideal white noise residuals (particularly in the

tails and Ljung-Box test), this is common in financial data due to volatility clustering, which we address in the GARCH section.

4.1.4 Step 6: Forecasting



Figure 5: 12-Step Ahead Forecast from ARIMA(1,1,1)

The graph above shows log vix forcasted over the next 12 days. The forecast shows mean reversion toward the long-run average, with expanding confidence intervals reflecting increasing uncertainty. The 80% and 95% prediction intervals widen rapidly with the forecast horizon, indicating increasing uncertainty about future VIX levels. This is characteristic of financial volatility forecasts and reflects both model uncertainty and the inherent unpredictability of market fear. The constant mean forecast after a few steps is typical of ARIMA models. While useful for short-term prediction, the model cannot anticipate future crises or regime changes that drive major VIX spikes. The forecast assumes the historical data-generating process continues unchanged. This forecast represents expected market volatility under normal conditions. The model cannot predict extraordinary events (market crashes, pandemics)

that cause extreme VIX spikes.

4.2 GARCH

The ARIMA model diagnostics revealed heavy-tailed residuals, volatility clustering in residuals, and significant Ljung-Box statistics. These features suggest the presence of conditional heteroskedasticity, which GARCH models are designed to capture. In this project, a GARCH(1,1) model is applied to the VIX return series in order to capture the persistence of volatility over time.

4.2.1 Testing for ARCH Effects

Table 5: ARCH-LM Test for Conditional Heteroskedasticity

Test Statistic	Value
Chi-squared	412.0959
Degrees of Freedom	12
P-value	$< 2.2e\text{-}16$
Conclusion	ARCH effects present

The highly significant ARCH-LM test confirms the presence of conditional heteroskedasticity, justifying the use of GARCH modeling.

4.2.2 GARCH Model Specification

We employ a GARCH(1,1) specification, which has been shown to adequately capture volatility dynamics in most financial applications (Hansen and Lunde (2005)). The model balances parsimony with the ability to capture volatility clustering and persistence, the key features of VIX dynamics. Furthermore, this will give us parameters that have clear economic meaning making it easier to interpret.

While we could test higher-order specifications (GARCH(2,1), GARCH(1,2)), the literature strongly suggests that additional parameters rarely improve out-of-sample forecast performance for daily financial data.

Table 6: GARCH(1,1) Parameter Estimates with Student-t Errors

	Parameter	Estimate	Std. Error	t-statistic	p-value	Significance
Mean Equation						
mu	mu (mean)	-0.001986	0.000196	-10.121	< 0.001	***
ar1	AR(1)	0.880179	0.004956	177.601	< 0.001	***
ma1	MA(1)	-0.957919	0.000564	-1698.083	< 0.001	***
Variance Equation						
omega	omega	0.000417	0.000057	7.297	< 0.001	***
alpha1	alpha1	0.133904	0.013767	9.726	< 0.001	***
beta1	beta1	0.776131	0.022048	35.202	< 0.001	***
Distribution						
shape	nu (shape)	4.718420	0.228806	20.622	< 0.001	***

Note: Significance: *** p<0.001, ** p<0.01, * p<0.05

Table 7: GARCH(1,1) Model Fit Statistics

Statistic	Value
Log-Likelihood	12806.70
AIC	-2.81
BIC	-2.80
Shibata	-2.81
Hannan-Quinn	-2.80

Table 8: GARCH Persistence Measures

Measure	Value
alpha1 + beta1 (Persistence)	0.910000
Unconditional Variance	0.004632
Half-life (days)	7.400000

Note: Half-life: time for volatility shock to decay to 50% of initial impact

Key Findings:

- $\alpha + \beta$ (ARCH + GARCH) $\approx$ close to 1, indicating high persistence in volatility
- **Student-t shape parameter** < 30 confirms heavy tails
- Both ARCH and GARCH effects are highly significant

Table 6 presents the estimated GARCH(1,1) parameters with Student-t distributed errors. All parameters are highly significant ($p < 0.001$), confirming the model's validity. The mean equation reveals strong ARMA dynamics in VIX log returns. The AR(1) coefficient is large and positive, indicating substantial positive autocorrelation in returns. This suggests that VIX movements tend to persist in the short term. The MA(1) coefficient is large and negative, nearly offsetting the AR component. This near-cancellation is characteristic of overdifferenced series and confirms our earlier finding of long memory in VIX. The mean return is slightly negative but economically negligible. On an annualized basis, this represents approximately -0.5% drift in log(VIX), consistent with VIX's mean-reverting nature.

Table 7 reports goodness-of-fit statistics. The log-likelihood of 12,806.70 is substantially higher than comparable ARIMA-only models, confirming that GARCH significantly improves fit. The information criteria (AIC, BIC, Hannan-Quinn) are all negative, which is typical for well-fitting GARCH models on financial returns data.

Table 8 presents derived measures of volatility persistence, which are crucial for understanding VIX dynamics.

4.2.3 Forecasting



Figure 6: Conditional Volatility Forecast

Table 9: 12-Day GARCH Forecast

Day	Conditional Mean	Conditional Volatility	Annualized Vol (%)
1	-0.022147	0.08130	129.06
2	-0.019732	0.08020	127.31
3	-0.017605	0.07918	125.70
4	-0.015734	0.07825	124.21
5	-0.014087	0.07739	122.85
6	-0.012637	0.07659	121.59
7	-0.011361	0.07587	120.43
8	-0.010237	0.07520	119.37
9	-0.009249	0.07458	118.40
10	-0.008378	0.07402	117.50
11	-0.007613	0.07350	116.68
12	-0.006938	0.07303	115.93

The GARCH(1,1) volatility forecast exhibits **mean reversion** toward the unconditional volatility level. Key observations:

1. **Volatility Decay**: Forecast volatility gradually converges to the long-run unconditional volatility. This reflects the mean-reverting property implied by $\alpha_1 + \beta_1 < 1$

2. **Half-Life Effect**: With a half-life of 7.4 days, the forecast shows that if current volatility is elevated, it will take approximately 1-2 weeks to decay halfway to the unconditional level.

3. **No Regime Prediction**: Like the ARIMA mean forecast, GARCH cannot predict future crises or regime changes. The forecast assumes the current volatility regime continues and gradually reverts to the long-run average.

Current vs. Long-run: If the current conditional volatility is 0.08695 and unconditional is 0.06806, the forecast shows mean reversion downward as volatility returns to its equilibrium level.

5 Conclusion and Future Study

5.1 Acknowledgements

6 Appendix

Follow the code at this github repo

```
# Load required libraries
library(readr)
library(forecast)
library(astsa)
library(tseries)
library(rugarch)
library(ggplot2)
library(gridExtra)
library(knitr)
library(kableExtra)
library(moments)
```

6.1 Data

6.1.1 Summary Statistics Table

```
# Load the VIX data
vix_data <- read_csv("Data/VIXCLS.csv", show_col_types = FALSE)
vix_data <- na.omit(vix_data)

# Convert to time series object
vix_ts <- ts(vix_data$VIXCLS,
             start = c(1990, 1),
             frequency = 252)  # Trading days per year

# Basic statistics
vix_stats <- data.frame(
  Statistic = c("Mean", "Median", "Std. Dev.", "Min", "Max", "N"),
  Value = c(
    mean(vix_data$VIXCLS, na.rm = TRUE),
    median(vix_data$VIXCLS, na.rm = TRUE),
```

```r
    sd(vix_data$VIXCLS, na.rm = TRUE),
    min(vix_data$VIXCLS, na.rm = TRUE),
    max(vix_data$VIXCLS, na.rm = TRUE),
    nrow(vix_data)
  )
)

kable(vix_stats,
      digits = 2,
      caption = "Summary Statistics of VIX",
      booktabs = TRUE) %>%
  kable_styling(latex_options = c("hold_position"))
```

6.1.2 EDA

```r
# Time series plot
plot(vix_ts,
     main = "CBOE Volatility Index (VIX) - Daily Close",
     ylab = "VIX Level",
     xlab = "Year",
     lwd = 0.5)

# Add horizontal line at mean
abline(h = mean(vix_data$VIXCLS, na.rm = TRUE),
       col = "red",
       lty = 2,
       lwd = 2)

# Highlight major crisis periods
rect(2008, 0, 2009, 100, col = rgb(1, 0, 0, 0.1), border = NA)
rect(2020, 0, 2021, 100, col = rgb(1, 0, 0, 0.1), border = NA)

text(2008.5, 75, "2008 Financial\nCrisis", cex = 0.8)
text(2020.5, 75, "COVID-19\nPandemic", cex = 0.8)
```

```r
# Histogram and QQ
par(mfrow = c(1, 2))

# Histogram
hist(vix_data$VIXCLS,
     breaks = 50,
     main = "Histogram of VIX",
     xlab = "VIX Level",
     col = "lightblue",
     border = "white")

# Q-Q plot
qqnorm(vix_data$VIXCLS,
       main = "Q-Q Plot: VIX vs Normal",
       pch = 20,
       col = alpha("darkblue", 0.5))
qqline(vix_data$VIXCLS,
       col = "red",
       lwd = 2)
```

```r
par(mfrow = c(1, 1))
```

6.2 Results

6.2.1 SARMIA

```r
log_vix <- log(vix_data$VIXCLS)

# ADF test on levels
adf_levels <- adf.test(log_vix)

# ADF test on first differences
dlog_vix <- diff(log_vix)
adf_diff <- adf.test(dlog_vix)
```

```r
# Create results table
adf_results <- data.frame(
  Series = c("Log(VIX)", " Diff Log(VIX)"),
  `ADF Statistic` = c(adf_levels$statistic, adf_diff$statistic),
  `P-value` = c(adf_levels$p.value, adf_levels$p.value),
  Conclusion = c("Stationary", "Stationary")
)

kable(adf_results,
      digits = 4,
      caption = "Augmented Dickey-Fuller Test Results",
      booktabs = TRUE) %>%
  kable_styling(latex_options = "hold_position")

# ACF PACF
par(mfrow = c(2, 1))

acf(log_vix,
    lag.max = 100,
    main = "ACF of Log(VIX)",
    na.action = na.pass)

pacf(log_vix,
     lag.max = 100,
     main = "PACF of Log(VIX)",
     na.action = na.pass)
```

```r
par(mfrow = c(1, 1))

# model comparison
# Fit candidate models
fit011 <- arima(log_vix, order = c(0, 1, 1))
fit110 <- arima(log_vix, order = c(1, 1, 0))
fit111 <- arima(log_vix, order = c(1, 1, 1))
```

```r
# Compare models
model_comparison <- data.frame(
  Model = c("ARIMA(0,1,1)", "ARIMA(1,1,0)", "ARIMA(1,1,1)"),
  AIC = c(AIC(fit011), AIC(fit110), AIC(fit111)),
  BIC = c(BIC(fit011), BIC(fit110), BIC(fit111))
)

# Highlight best model
model_comparison$Best_AIC <- ifelse(model_comparison$AIC == min(model_comparison$AIC), '
model_comparison$Best_BIC <- ifelse(model_comparison$BIC == min(model_comparison$BIC), '

kable(model_comparison,
      digits = 2,
      caption = "Model Comparison - Information Criteria (* indicates best model)",
      booktabs = TRUE) %>%
  kable_styling(latex_options = "hold_position")

# Best model
best_arima <- fit111

# Display coefficients
coef_table <- data.frame(
  Parameter = names(coef(best_arima)),
  Estimate = coef(best_arima),
  SE = sqrt(diag(vcov(best_arima)))
)
coef_table$`t-value` <- coef_table$Estimate / coef_table$SE
coef_table$`p-value` <- 2 * (1 - pnorm(abs(coef_table$`t-value`)))

kable(coef_table,
      digits = 4,
      caption = "ARIMA(1,1,1) Parameter Estimates",
      booktabs = TRUE,
      row.names = FALSE) %>%
  kable_styling(latex_options = "hold_position")
```

```r
# fitting Sarima
# Use sarima for nice diagnostic plots
sarima_fit <- sarima(log_vix, 1, 1, 1)
```

```r
# Forecasting
# You already have this from the data section
vix_ts <- ts(vix_data$VIXCLS,
             start = c(1990, 1),
             frequency = 252)

# Take log of the ts object
log_vix_ts <- log(vix_ts)

# Now fit ARIMA
best_arima <- arima(log_vix_ts, order = c(1, 1, 1))

# Forecast
fc_arima <- forecast(best_arima, h = 12)

# Plot - will show years on x-axis
plot(fc_arima,
     main = "ARIMA(1,1,1) Forecast: Next 12 Days",
     xlab = "Time",
     ylab = "Log(VIX)",
     xlim = c(2025.5, 2026.3))  # Adjust to show end of series
```

6.2.2 GARCH

```r
# Compute squared residuals
resid_arima <- residuals(best_arima)

# ARCH-LM test
arch_test <- ArchTest(resid_arima, lags = 12)
```

```r
# Create formatted table
arch_results <- data.frame(
  Statistic = c("Chi-squared", "Degrees of Freedom", "P-value", "Conclusion"),
  Value = c(
    round(arch_test$statistic, 4),
    12,
    format.pval(arch_test$p.value, digits = 4),
    ifelse(arch_test$p.value < 0.05,
           "ARCH effects present",
           "No ARCH effects")
  )
)

kable(arch_results,
      caption = "ARCH-LM Test for Conditional Heteroskedasticity",
      booktabs = TRUE,
      col.names = c("Test Statistic", "Value")) %>%
  kable_styling(latex_options = "hold_position")

# Prepare log returns
log_returns <- diff(log(vix_data$VIXCLS))

# Specify GARCH(1,1) model
garch_spec <- ugarchspec(
  variance.model = list(model = "sGARCH", garchOrder = c(1, 1)),
  mean.model = list(armaOrder = c(1, 1), include.mean = TRUE),
  distribution.model = "std"  # Student-t for heavy tails
)

# Fit the model
garch_fit <- ugarchfit(spec = garch_spec, data = log_returns)

# Extract coefficients and standard errors
garch_coef <- coef(garch_fit)
garch_se <- garch_fit@fit$matcoef[, 2]  # Standard errors
```

```r
garch_tstat <- garch_fit@fit$matcoef[, 3]  # t-statistics
garch_pval <- garch_fit@fit$matcoef[, 4]  # p-values

# Create beautiful parameter table
param_table <- data.frame(
  Parameter = c("mu (mean)", "AR(1)", "MA(1)",
                "omega", "alpha1", "beta1",
                "nu (shape)"),
  Estimate = round(garch_coef, 6),
  `Std. Error` = round(garch_se, 6),
  `t-statistic` = round(garch_tstat, 3),
  `p-value` = ifelse(garch_pval < 0.001, "< 0.001",
                     round(garch_pval, 4)),
  Significance = ifelse(garch_pval < 0.001, "***",
                 ifelse(garch_pval < 0.01, "**",
                 ifelse(garch_pval < 0.05, "*", ""))),
  check.names = FALSE
)

kable(param_table,
      caption = "GARCH(1,1) Parameter Estimates with Student-t Errors",
      booktabs = TRUE,
      align = c('l', 'r', 'r', 'r', 'r', 'c')) %>%
  kable_styling(latex_options = c("hold_position")) %>%
  pack_rows("Mean Equation", 1, 3) %>%
  pack_rows("Variance Equation", 4, 6) %>%
  pack_rows("Distribution", 7, 7) %>%
  footnote(general = "Significance: *** p<0.001, ** p<0.01, * p<0.05",
           general_title = "Note:",
           footnote_as_chunk = TRUE)

# Model fit statistics
fit_stats <- data.frame(
  Statistic = c("Log-Likelihood", "AIC", "BIC", "Shibata", "Hannan-Quinn"),
  Value = c(
```

```r
    likelihood(garch_fit),
    infocriteria(garch_fit)[1],
    infocriteria(garch_fit)[2],
    infocriteria(garch_fit)[3],
    infocriteria(garch_fit)[4]
  )
)

kable(fit_stats,
      caption = "GARCH(1,1) Model Fit Statistics",
      booktabs = TRUE,
      digits = 2,
      align = c('l', 'r')) %>%
  kable_styling(latex_options = "hold_position")

# Calculate persistence and half-life
alpha <- garch_coef["alpha1"]
beta <- garch_coef["beta1"]
persistence <- alpha + beta
half_life <- log(0.5) / log(persistence)

# Persistence table
persistence_table <- data.frame(
  Measure = c("alpha1 + beta1 (Persistence)",
              "Unconditional Variance",
              "Half-life (days)"),
  Value = c(
    round(persistence, 4),
    ifelse(persistence < 1,
           round(garch_coef["omega"] / (1 - persistence), 6),
           "Undefined (alpha+beta >= 1)"),
    ifelse(persistence < 1,
           round(half_life, 1),
           "infinity (non-stationary)")
  )
```

```
)

kable(persistence_table,
      caption = "GARCH Persistence Measures",
      booktabs = TRUE,
      align = c('l', 'r')) %>%
  kable_styling(latex_options = "hold_position") %>%
  footnote(general = "Half-life: time for volatility shock to decay to 50% of initial im
           general_title = "Note:",
           footnote_as_chunk = TRUE)

# Forecasting
# Forecast 12 steps ahead
garch_fc <- ugarchforecast(garch_fit, n.ahead = 12)

# Extract forecasts
fc_mean  <- as.numeric(fitted(garch_fc))
fc_sigma <- as.numeric(sigma(garch_fc))  # Conditional volatility forecast

hist_sigma <- as.numeric(sigma(garch_fit))
n <- length(hist_sigma)
h <- length(fc_sigma)

# Plot historical + forecast volatility
plot(c((n-100):n, (n+1):(n+h)),
     c(tail(hist_sigma,101), fc_sigma),
     type = "n",
     main = "GARCH(1,1) Conditional Volatility Forecast",
     xlab = "Time Index",
     ylab = "Conditional Volatility",
     xlim = c(n-100, n+h+2))

# Historical volatility
lines((n-100):n, tail(hist_sigma,101), col = "darkblue", lwd = 1.5)
```

```r
# Forecast volatility
lines((n+1):(n+12), as.numeric(fc_sigma), col = "red", lwd = 2, lty = 1)
points((n+1):(n+12), as.numeric(fc_sigma), col = "red", pch = 19)

# Add vertical line at forecast start
abline(v = n, col = "gray50", lty = 2, lwd = 2)

# Legend
legend("topright",
       legend = c("Historical sigma", "Forecast sigma"),
       col = c("darkblue", "red"),
       lwd = c(1.5, 2),
       bty = "n")

# Add unconditional volatility line
uncond_vol <- sqrt(garch_coef["omega"] / (1 - persistence))
abline(h = uncond_vol, col = "darkgreen", lty = 2, lwd = 1.5)
text(n-50, uncond_vol + 0.005,
     "Unconditional sigma", col = "darkgreen", cex = 0.8)
```

```r
fc_table <- data.frame(
  Day = 1:12,
  `Conditional Mean` = round(as.numeric(fc_mean), 6),
  `Conditional Volatility` = round(as.numeric(fc_sigma), 5),
  `Annualized Vol (%)` = round(as.numeric(fc_sigma) * sqrt(252) * 100, 2),
  check.names = FALSE
)

kable(fc_table,
      caption = "12-Day GARCH Forecast",
      booktabs = TRUE,
      align = c('c','r','r','r')) %>%
  kable_styling(latex_options = c("hold_position","scale_down"))
```

References

Aljandali, Abdulkader. 2017. "The Box-Jenkins Methodology." In *Multivariate Methods and Forecasting with IBM® SPSS® Statistics*, 59–79. Springer.

Blair, Bevan J., Ser-Huang Poon, and Stephen J. Taylor. 2001. "Forecasting s&p 100 Volatility: The Incremental Information Content of Implied Volatilities and High-Frequency Index Returns." *Journal of Econometrics* 105 (1): 5–26. https://doi.org/https://doi.org/10.1016/S0304-4076(01)00068-9.

Bollerslev, Tim. 1986. "Generalized Autoregressive Conditional Heteroskedasticity." *Journal of Econometrics* 31 (3): 307–27. https://doi.org/https://doi.org/10.1016/0304-4076(86)90063-1.

Bouman, Sven, and Ben Jacobsen. 2002. "The Halloween Indicator, "Sell in May and Go Away": Another Puzzle." *American Economic Review* 92 (5): 1618–35. https://doi.org/10.1257/000282802762024683.

Caporale, Guglielmo Maria, and Alex Plastun. 2019. "On Stock Price Overreactions: Frequency, Seasonality and Information Content." *Journal of Applied Economics* 22 (1): 602–21.

Chicago Board Options Exchange. 2019. "VIX White Paper." https://www.cboe.com/tradable_products/vix/vix_white_paper/.

Corsi, Fulvio. 2009. "A Simple Approximate Long-Memory Model of Realized Volatility." *Journal of Financial Econometrics* 7: 174–96.

Ellul, Andrew, Hyun Song Shin, and Ian Tonks. 2005. "Opening and Closing the Market: Evidence from the London Stock Exchange." *Journal of Financial and Quantitative Analysis* 40 (4): 779–801.

Engle, Robert F, and Tim Bollerslev. 1986. "Modelling the Persistence of Conditional Variances." *Econometric Reviews* 5 (1): 1–50.

Fernandes, Marcelo, Marcelo Cunha Medeiros, and MArcelo Scharth. 2007. "Modeling and Predicting the CBOE Market Volatility Index." Textos para discussÃ£o 548. Department of Economics PUC-Rio (Brazil). https://doi.org/None.

Hansen, Peter R, and Asger Lunde. 2005. "A Forecast Comparison of Volatility Models: Does Anything Beat a GARCH (1, 1)?" *Journal of Applied Econometrics* 20 (7): 873–89.

Hughes, Michael P, Drew B Winters, and Jerry S Rawls. 2005. "What Is the Source of Different Levels of Time-Series Return Volatility? The Intraday u-Shaped Pattern or Time-Series Persistence." *Journal of Economics and Finance* 29 (3): 300–312.

Kambouroudis, Dimos S., and David G. McMillan. 2016. "Does VIX or Volume Improve GARCH Volatility Forecasts?" *Applied Economics* 48 (13): 1210–28. https://doi.org/

10.1080/00036846.2015.1096004.

Majka, Marcin. n.d. "GARCH Model in Finance."

Majmudar, Utkarsh, and Arnab Banerjee. 2004. "Vix Forecasting."

Makridakis, Spyros, and Michele Hibon. 1997. "ARMA Models and the Box–Jenkins Methodology." *Journal of Forecasting* 16 (3): 147–63.

Wang, Hui. 2019. "VIX and Volatility Forecasting: A New Insight." *Physica A: Statistical Mechanics and Its Applications* 533: 121951. https://doi.org/https://doi.org/10.1016/j.physa.2019.121951.